arXiv:hep-th/9908171v1 26 Aug 1999

ROM2F/99/19

$\mathcal{N}$=4 supersymmetric Yang–Mills theory and the AdS/SCFT correspondence

Stefano Kovacs

Dipartimento di Fisica, Università di Roma "Tor Vergata"
I.N.F.N. - Sezione di Roma "Tor Vergata"
Via della Ricerca Scientifica, 1
00173 Roma, ITALY

Abstract

This dissertation reviews various aspects of the $\mathcal{N}$=4 supersymmetric Yang–Mills theory in particular in relation with the AdS/CFT correspondence.

The first two chapters are introductory. The first one contains a description of the general properties of rigid supersymmetric theories in four dimensions both at the classical and at the quantum level. The second chapter is a review of the main properties of the $\mathcal{N}$=4 SYM theory under consideration.

Original results are reported in chapters 3, 4 and 5. A systematic re-analysis of the perturbative properties of the theory is presented in the third chapter. Two-, three- and four-point Green functions of elementary fields are computed using the component formulation and/or the superfield approach and subtleties related to the gauge-fixing are pointed out. In the fourth chapter, after an introduction to instanton calculus in supersymmetric gauge theories, the computation of the one-instanton contributions to Green functions of gauge invariant composite operators in the semiclassical approximation is reported. The calculations of four-, eight- and sixteen-point Green functions of operators in the supercurrent multiplet are reviewed in detail. The final chapter is devoted to the AdS/SCFT correspondence. Some general aspects are discussed. Then the attention is focused on the relation between instantons in $\mathcal{N}$=4 SYM and D-instanton effects in type IIB string theory. The comparison between instanton contributions to Green functions of composite operators in the boundary field theory and D-instanton generated terms in the amplitudes computed in type IIB string theory is performed and agreement between these two sources of non-perturbative effects is shown.

UNIVERSITÀ DEGLI STUDI DI ROMA
"TOR VERGATA"

FACOLTÀ DI SCIENZE MATEMATICHE, FISICHE E NATURALI
Dipartimento di Fisica

$\mathcal{N}$=4 supersymmetric Yang–Mills theory and the AdS/SCFT correspondence

Tesi di dottorato di ricerca in Fisica

presentata da

Stefano Kovacs

Relatori

Professor *Giancarlo Rossi*

Professor *Massimo Bianchi*

Coordinatore del dottorato

Professor *Piergiorgio Picozza*

Ciclo XI

Anno Accademico 1997-1998

Contents

Acknowledgments **v**

Note added **v**

1 Rigid supersymmetric theories in four dimensions **1**

1.1 General supersymmetry algebra in four dimensions 2

1.2 Representations of the supersymmetry algebra 5

1.3 $\mathcal{N}$=1 superspace 10

1.4 $\mathcal{N}$=1 supersymmetric models 17

1.4.1 Supersymmetric chiral theories: Wess–Zumino model 18

1.4.2 Supersymmetric gauge theories 19

1.5 Theories of extended supersymmetry 22

1.5.1 Extended superspace: a brief survey 23

1.5.2 $\mathcal{N} > 1$ supersymmetric models 27

1.6 Quantization of supersymmetric theories 29

1.6.1 General formalism 30

1.6.2 Superfield propagators 33

1.6.3 Feynman rules 35

1.7 General properties of supersymmetric theories 36

2 $\mathcal{N} = 4$ supersymmetric Yang–Mills theory **40**

2.1 Diverse formulations of $\mathcal{N}$=4 supersymmetric Yang–Mills theory . . . 41

2.2 Classical symmetries and conserved currents 46

2.3 Finiteness of $\mathcal{N}$=4 supersymmetric Yang–Mills theory 50

2.4 S-duality 52

2.4.1 Montonen–Olive and SL(2,$\mathbb{Z}$) duality 54

2.4.2 S-duality in $\mathcal{N}$=4 supersymmetric Yang–Mills theory 57

2.4.3 Semiclassical quantization 60

3 Perturbative analysis of $\mathcal{N} = 4$ supersymmetric Yang–Mills theory 66

3.1 Perturbation theory in components: problems with the Wess–Zumino gauge 68

3.1.1 One loop corrections to the propagator of the fermions belonging to the chiral multiplet 72

3.1.2 One loop corrections to the propagator of the scalars belonging to the chiral multiplet 74

3.2 Perturbation theory in $\mathcal{N}$=1 superspace: propagators 79

3.2.1 Propagator of the chiral superfield 81

3.2.2 Propagator of the vector superfield 86

3.2.3 Discussion 93

3.3 Three- and four-point functions of $\mathcal{N}$=1 superfields 96

3.3.1 Three-point functions 96

3.3.2 Four-point functions 99

4 Nonperturbative effects in $\mathcal{N} = 4$ super Yang–Mills theory 108

4.1 Instanton calculus in non Abelian gauge theories 110

4.1.1 Yang–Mills instantons 111

4.1.2 Semiclassical quantization 114

4.1.3 Bosonic zero-modes 116

4.2 Instanton calculus in the presence of fermionic fields 118

4.2.1 Semiclassical approximation 119

4.2.2 Fermionic zero-modes 121

4.3 Instanton calculus in supersymmetric gauge theories 123

4.4 Instanton calculations in $\mathcal{N}$=4 supersymmetric Yang–Mills theory . . 125

4.5 A correlation function of four scalar supercurrents 126

4.6 Eight- and sixteen-point correlation functions of current bilinears . . 136

4.6.1 The correlation function of sixteen fermionic currents 136

4.6.2 The correlation function of eight gaugino bilinears 137

4.7 Generalization to arbitrary N and to any K for large N 138

5 AdS/SCFT correspondence **141**
5.1 Type IIB superstring theory: a bird's eye view 142
5.2 D-branes 147
5.3 The AdS/SCFT correspondence conjecture 151
5.4 Calculations of two- and three-point functions 160
5.5 Four-point functions 170
5.6 D-instanton effects in d=10 type IIB superstring theory 177
5.6.1 Testing the AdS/CFT correspondence: type IIB D-instantons vs. Yang–Mills instantons 181
5.7 The D-instanton solution in $AdS_5 \times S^5$ 184

Conclusions **189**

A Notations and spinor algebra **193**

B Electric-magnetic duality **198**
B.1 Electric-magnetic duality and the Dirac monopole 198
B.2 Non Abelian case: the 't Hooft–Polyakov monopole 200
B.3 Monopoles and fermions 204

Bibliography **207**

Acknowledgments

This dissertation is based on research done at the Department of Physics of the Università di Roma "Tor Vergata"*, during the period November 1995 - October 1998, under the supervision of Prof. Massimo Bianchi and Prof. Giancarlo Rossi. I would like to thank them for introducing me to the arguments covered in this thesis, for all what they have taught me and for their willingness and constant encouragement. I also wish to thank Prof. Michael B. Green for the very enjoyable and fruitful collaboration on the subjects reported in the last two chapters. I am grateful to Yassen Stanev for many useful and interesting discussions. Finally I thank the whole theoretical physics group at the Department of Physics of the* Università di Roma "Tor Vergata" *for the very stimulating environment I have found there.*

Note added

This thesis was presented to the "Università di Roma *Tor Vergata*" in candidacy for the degree of "Dottore di Ricerca in Fisica" and it was defended in April 1999. Hence more recent developments are not discussed here and moreover the references are not updated. However I cannot help mentioning at least the very detailed and comprehensive review by O. Arhony, S. Gubser, J. Maldacena, H. Ooguri and Y. Oz on "Large N Field Theories, String Theory and Gravity" (hep-th/9905111), that is not cited in the bibliography. Actually this paper would have made my writing of the last chapter much simpler if it came out earlier!

There are some direct developments of this thesis that have been carried on during the last few months. In particular the analysis of logarithmic singularities in four-point Green functions and their relation to anomalous dimensions of un-protected operators in $\mathcal{N}$=4 SYM has been finalised in collaboration with Massimo Bianchi, Giancarlo Rossi and Yassen Stanev. This issue is only briefly addressed in the concluding remarks. The interested reader is referred to the paper "On the logarithmic behaviour in $\mathcal{N}$=4 SYM theory", hep-th/9906188.

Chapter 1

Rigid supersymmetric theories in four dimensions

Supersymmetry was introduced in [1] as a generalization of Poincaré invariance and then implemented in the context of four dimensional quantum field theory in [2]. The idea of a symmetry transformation exchanging bosonic and fermionic degrees of freedom has proved extremely rich of consequences. Although up to now there exists no experimental evidence for supersymmetry, it plays a central rôle in theoretical physics.

At the present day the interest in the study of supersymmetric theories relies on one side on the rôle it plays in the construction of a more fundamental model beyond the standard model of particle physics (SM) and on the other on the possibility of deriving results that can hopefully be extended to phenomenologically more relevant models.

The standard model describes the strong and electro-weak interactions within the framework of a SU(3)×SU(2)×U(1) non-Abelian gauge theory; despite the full agreement with experimental data there are various motivations for believing that it cannot be a truly fundamental theory. A first problem arises in relation with the grand unification of particle interactions. The gauge couplings are believed to be unified at an energy scale of the order of 10^{16} GeV. The standard model contains many free parameters (masses and mixing angles) that must be "fine-tuned" in order to achieve the grand unification (*hierarchy problem*): the problem follows from the

existence of ultra-violet quadratic divergences and is solved in supersymmetric models in which the divergences, as will be discussed later, are milder.

Moreover the standard model does not include gravitational interactions: at present the only consistent description of gravity at the quantum level is provided by (super)string theories and supersymmetry appears to be a fundamental ingredient of such theories.

In a more immediate perspective supersymmetric theories can be of help in understanding aspects of phenomenologically interesting models. For example a description of confinement in asymptotically free theories as dual superconductivity has been proposed and various realizations of this mechanism have been observed in supersymmetric models.

More recently a correspondence between four dimensional quantum field theories and supergravity has been suggested; this particular subject appears very promising and will be extensively studied in chapter 5.

This first chapter contains a review of rigid supersymmetry in four dimensions. The material presented is by no means original, the aim is to establish the notations and to recall ideas and results, mainly about the quantization of supersymmetric theories, that will be extensively used in the following.

In the first part of the chapter the supersymmetry algebra and its realizations are described. Then the formalism of superspace is introduced in somewhat more detail both at the classical and at the quantum level. The last section reviews general properties of supersymmetric models.

An introduction to supersymmetry is provided by the books by P. West [3] and J. Wess and J. Bagger [4] (whose notations will be followed in this thesis) and by the review papers [5, 6]; superspace techniques are treated in more detail in [7]; for phenomenological aspects of supersymmetry see [8]; an extensive introduction to superstring theory is given in the book by M.B. Green, J.H. Schwarz and E. Witten [9].

1.1 General supersymmetry algebra in four dimensions

Quantum field theory models of interest for the description of fundamental interactions combine the Poincaré invariance with the invariance under a global internal symmetry group G. In [10] Coleman and Mandula proved, under general assumptions

following from the axioms of quantum field theory, a "no-go" theorem that, states that the most general symmetry group for a theory with non trivial S-matrix is the direct product of the Poincaré group and an internal group G, which must be of the form of a semisimple Lie group times possible U(1) factors.

Historically supersymmetry was discovered as an attempt to evade the constraints of the Coleman–Mandula no-go theorem .

The Poincaré group P=ISO(3,1) is the semidirect product of the group T_4 of four dimensional translations and the Lorentz group SO(3,1); it is generated by P_μ, $M_{\mu\nu}$ satisfying:

$$\begin{aligned}
&[P_\mu, P_\nu] = 0 \\
&[P_\lambda, M_{\mu\nu}] = (\eta_{\lambda\mu} P_\nu - \eta_{\lambda\nu} P_\mu) \\
&[M_{\mu\nu}, M_{\rho\sigma}] = -(\eta_{\mu\rho} M_{\nu\sigma} + \eta_{\nu\sigma} M_{\mu\rho} - \eta_{\mu\sigma} M_{\nu\rho} - \eta_{\nu\rho} M_{\mu\sigma})\,.
\end{aligned} \tag{1.1}$$

(See appendix A for the conventions and a collection of useful formulas).

The generic internal symmetry group is a Lie group defined by the commutation relations:

$$[T^a, T^b] = i f^{ab}{}_c T^c\,. \tag{1.2}$$

Internal symmetries are supplemented by the discrete ones C, P and T.

The supersymmetry algebra is a graded extension of (1.1): the theorem of Coleman and Mandula is bypassed by allowing anticommuting as well as commuting generators. In [11] the most general algebra allowed by this weakening of the hypothesis of [10] was constructed: it contains, beyond (1.1) and (1.2), anticommuting generators Q^i_α, $\overline{Q}_{\dot\alpha j}$ that are Weyl spinors transforming in the $(\frac{1}{2}, 0)$ and $(0, \frac{1}{2})$ representations of the Lorentz group respectively, so that they do not commute with the generators P_μ and $M_{\mu\nu}$. Here $\alpha, \dot\alpha = 1, 2$ are spinor indices and i, j ranging from 1 to $\mathcal{N} \geq 1$ label the various of supersymmetries.

The most general symmetry algebra of a supersymmetric quantum field theory is therefore given by (1.1) and (1.2) supplemented by

$$\begin{aligned}
&\{Q^i_\alpha, \overline{Q}^j_\beta\} = 2\sigma^\mu_{\alpha\dot\alpha} P_\mu \delta^{ij} \\
&[P_\mu, Q^i_\alpha] = [P_\mu, \overline{Q}_{\dot\alpha j}] = 0 \\
&[Q^i_\alpha, M_{\mu\nu}] = (\sigma_{\mu\nu})_\alpha{}^\beta Q^i_\beta
\end{aligned}$$

$$
\begin{aligned}
&[\overline{Q}_{\dot\alpha j}, M_{\mu\nu}] = (\overline{\sigma}_{\mu\nu})^{\dot\alpha}{}_{\dot\beta}\overline{Q}^{\dot\beta}_{j} \\
&\{Q^i_\alpha, Q^j_\beta\} = \varepsilon_{\alpha\beta} Z^{ij} \\
&\{\overline{Q}_{\dot\alpha i}, \overline{Q}_{\dot\beta j}\} = \varepsilon_{\dot\alpha\dot\beta} Z^\dagger_{ij} \\
&[Q^i_\alpha, T^a] = B^{ai}{}_j Q^j_\alpha \\
&[T^a, \overline{Q}_{\dot\alpha i}] = B^{\dagger ai}_j \overline{Q}_{\dot\alpha i} \\
&[Z^{ij}, X] = [Z^\dagger_{lk}, X] = 0\,,
\end{aligned}
\tag{1.3}
$$

where $B^{\dagger aj}_i = B^{aj}{}_i$. In (1.3) X denotes an arbitrary generator in the algebra so that $Z^{ij} = -Z^{ji}$ (central charges) generate the center of the algebra [1]. Central charges play a major rôle in the quantization of $\mathcal{N}$-extended supersymmetric theories. They generate an Abelian subalgebra so that

$$
Z^{ij} = a^{ij}_b T^b\,,
$$

with $B^i_{aj} a^{b\,jk} = -a^{b\,ij} B_{a\,j}{}^k$.

The supersymmetry algebra (1.3) considered as a graded Lie algebra possesses a group of automorphisms which is U($\mathcal{N}$) in the general case . The U($\mathcal{N}$) group of automorphisms acts on the charges Q^i_α and $\overline{Q}_{\dot\alpha j}$ as

$$
Q^i_\alpha \longrightarrow U^i{}_k Q^k_\alpha \qquad \overline{Q}_{\dot\alpha j} \longrightarrow \overline{Q}_{\dot\alpha k} U^{\dagger k}_j\,.
$$

In the particular case of $\mathcal{N}$=1 supersymmetry the U(1) group of automorphisms is generated by R such that

$$
[Q_\alpha, R] = Q_\alpha\,, \qquad [\overline{Q}_{\dot\alpha}, R] = -\overline{Q}_{\dot\alpha}\,.
$$

The supersymmetry charges Q^i_α and $\overline{Q}_{\dot\alpha j}$ transform in spin $\frac{1}{2}$ representations of the Lorentz group; this implies that acting with Q^i_α or $\overline{Q}_{\dot\alpha j}$ on a state of spin j produces a state of spin $j \pm \frac{1}{2}$: supersymmetry generators exchange bosonic and fermionic states. Reversing the line of reasoning and starting with a symmetry transforming the fields among themselves in such a way as to mix bosons and fermions, one concludes that the different physical dimension of bosonic and fermionic fields implies that the transformation must involve derivatives, *i.e.* space-time translations. This gives an

[1] In principle the antisymmetry of Z^{ij} is expected to imply that no central extension can exist in non extended ($\mathcal{N}$=1) theories. It can be proved [12] that a central extension can actually be present in $\mathcal{N}$=1 supersymmetric Yang–Mills theory. See [13] for a recent discussion in the context of the AdS/CFT correspondence.

intuitive explanation of how the requirement of a non trivial mixing of internal and space-time symmetries naturally leads to supersymmetry.

An immediate consequence of the supersymmetry algebra is that in supersymmetric theories the energy is positive definite and vanishes only on supersymmetric (ground) states. The first equation in (1.3) implies

$$0 \leq \sum_i \left(\{Q_1^i, (Q_{1i})^\dagger\} + \{Q_2^i, (Q_{2i})^\dagger\} \right) = -4\mathcal{N} P_0 = 4\mathcal{N} H \,,$$

so that

$$\langle \psi | H | \psi \rangle \geq 0$$

for every state $|\psi\rangle$ and $H|\Psi\rangle = 0$ if and only if $Q|\psi\rangle = 0$, *i.e.* if $|\psi\rangle$ is supersymmetric.

1.2 Representations of the supersymmetry algebra

The irreducible representations of the general supersymmetry algebra can be constructed starting from the (anti)commutation relations of the preceding section and the definition of the Casimir operators. For the Poincaré group the quadratic Casimir operators are $P^2 = P_\mu P^\mu$ and $W^2 = W_\mu W^\mu$, where W_μ is the Pauli–Lubanski vector

$$W_\mu = \frac{1}{2} \varepsilon_{\mu\nu\rho\sigma} P^\nu M^{\rho\sigma} \,.$$

Here W^2 is not a Casimir anymore, because $M_{\mu\nu}$ does not commute with the supersymmetry generators Q_α^i and $\overline{Q}_{\dot{\alpha} j}$, and is substituted by C^2 defined (for the case $\mathcal{N}$=1) by

$$\begin{aligned} C^2 &= C_{\mu\nu} C^{\mu\nu} \\ C_{\mu\nu} &= B_\mu P_\nu - B_\nu P_\mu \\ B_\mu &= W_\mu - \frac{1}{4} \overline{Q}_{\dot{\alpha}} \overline{\sigma}_\mu^{\dot{\alpha}\alpha} Q_\alpha \,. \end{aligned}$$

The irreducible representations can be constructed by Wigner's technique of induced representations.

A general result following from the supersymmetry algebra is that every irreducible representation contains an equal number of bosonic and fermionic states. Defining the fermion number operator $(-)^{N_f}$ such that

$$(-)^{N_f} |B\rangle = +|B\rangle \,, \qquad (-)^{N_f} |F\rangle = -|F\rangle \,,$$

where $|B\rangle$ denotes a bosonic state and $|F\rangle$ a fermionic one, it follows that

$$(-)^{N_f} Q_\alpha = -Q_\alpha (-)^{N_f} \,, \qquad (-)^{N_f} \overline{Q}_{\dot\alpha} = -\overline{Q}_{\dot\alpha} (-)^{N_f} \,.$$

These relations imply

$$0 = \mathrm{tr}\left[(-)^{N_f} \{Q_\alpha, \overline{Q}_{\dot\alpha}\}\right] = 2\sigma^\mu_{\alpha\dot\alpha} \delta^{ij} P_\mu \mathrm{tr}(-)^{N_f} \,,$$

so that in conclusion for non vanishing P_μ

$$\mathrm{tr}(-)^{N_f} = 0$$

from which the statement follows.

Massive representations without central charges.

For particles of mass M in the rest frame $P_\mu = (-M, 0, 0, 0)$ the supersymmetry algebra reduces to

$$\begin{aligned} \{Q^i_\alpha, \overline{Q}_{\dot\beta j}\} &= 2M \delta_{\alpha\dot\alpha} \delta^i_j \\ \{Q_\alpha, Q_\beta\} = \{\overline{Q}_{\dot\alpha i} \overline{Q}_{\dot\beta j}\} &= 0 \,, \end{aligned} \tag{1.4}$$

and creation and annihilation operators can be defined as follows

$$\begin{aligned} a^i_\alpha &= \frac{1}{\sqrt{2M}} Q^i_\alpha \\ \left(a^i_\alpha\right)^\dagger &= \frac{1}{\sqrt{2M}} \overline{Q}_{\dot\alpha i} \,. \end{aligned} \tag{1.5}$$

a^i_α and $(a^i_\alpha)^\dagger$ satisfy the algebra of fermionic creation and annihilation operators. In fact by direct calculation one finds

$$\begin{aligned} \{a^i_\alpha, \left(a^j_\beta\right)^\dagger\} &= \delta^\beta_\alpha \delta^i_j \\ \{a^i_\alpha, a^j_\beta\} &= \{\left(a^i_\alpha\right)^\dagger, \left(a^j_\beta\right)^\dagger\} = 0 \,. \end{aligned} \tag{1.6}$$

Given a Clifford vacuum $|\Omega\rangle$ defined by

$$a^i_\alpha |\Omega\rangle = 0 \qquad \forall\, i, \alpha \,,$$

with $P^2|\Omega\rangle = -M^2|\Omega\rangle$, the states are constructed by acting with the operators $(a^i_\alpha)^\dagger$ on $|\Omega\rangle$:

$$|\Omega^{(n)\,\alpha_1 \ldots \alpha_n}_{i_1 \ldots i_n}\rangle = \frac{1}{\sqrt{n!}} \left(a^{i_n}_{\alpha_n}\right)^\dagger \ldots \left(a^{i_1}_{\alpha_1}\right)^\dagger |\Omega\rangle \,. \tag{1.7}$$

Antisymmetry under the exchange of pairs (α_k, i_k), (α_l, i_l) implies $n \leq 2\mathcal{N}$. For each n the number of states (degeneracy) is $\begin{pmatrix} 2\mathcal{N} \\ n \end{pmatrix}$, so that the total number of states (*i.e.* the dimension of the representation) is

$$d = \sum_{n=0}^{2\mathcal{N}} \begin{pmatrix} 2\mathcal{N} \\ n \end{pmatrix} = 2^{2\mathcal{N}} .$$

The representation contains $2^{2\mathcal{N}-1}$ bosonic and $2^{2\mathcal{N}-1}$ fermionic states. The highest spin state is obtained symmetrizing the maximum number of spinor indices, namely $\mathcal{N}$, leading to the value $\frac{1}{2}\mathcal{N}$ for the highest spin in the multiplet.

Massless representations without central charges.

In the "light-like" reference frame $P_\mu = (-E, 0, 0, E)$, with $P^2 = 0$, the supersymmetry algebra reads

$$\begin{aligned} \{Q^i_\alpha, \overline{Q}_{\dot\beta j}\} &= 2 \begin{pmatrix} 2E & 0 \\ 0 & 0 \end{pmatrix} \delta^i_j \\ \{Q^i_\alpha, Q^j_\beta\} &= \{\overline{Q}_{\dot\alpha i}, \overline{Q}_{\dot\beta j}\} = 0 . \end{aligned}$$

The natural definition of creation and annihilation operators is then

$$\begin{aligned} a^i &= \frac{1}{2\sqrt{E}} Q^i_1 \\ a^\dagger_i &= (a^i)^\dagger = \frac{1}{2\sqrt{E}} \overline{Q}^i_{\dot 1} . \end{aligned} \tag{1.8}$$

They generate the Clifford algebra

$$\begin{aligned} \{a^i, a^\dagger_j\} &= \delta^i_j \\ \{a^i, a^j\} &= \{a^\dagger_i, a^\dagger_j\} = 0 \end{aligned} \tag{1.9}$$

Notice that in this case only half of the generators Q^i_α can be used to construct the fermionic oscillators because Q^i_2 and $\overline{Q}_{\dot 2 j}$ anticommute. Starting with a state of lowest helicity λ, $|\Omega_\lambda\rangle$, playing the rôle of a Clifford vacuum, defined by

$$a^i |\Omega_\lambda\rangle = 0$$

the states which constitute the multiplet are constructed as

$$|\Omega^{(n)}_{\lambda+\frac{n}{2}; i_1, \ldots, i_n}\rangle = \frac{1}{\sqrt{n!}} a^\dagger_{i_n} \ldots a^\dagger_{i_1} |\Omega_\lambda\rangle . \tag{1.10}$$

The state $|\Omega^{(n)}_{\lambda+\frac{n}{2};i_1,\dots,i_n}\rangle$ has helicity $\lambda+\frac{n}{2}$ and because of antisymmetry in the indices $i_1,\dots,i_n$ is $\begin{pmatrix}\mathcal{N}\\ n\end{pmatrix}$ times degenerate. The highest helicity in the multiplet is $\overline{\lambda}=\lambda+\frac{\mathcal{N}}{2}$. The dimension of the representation given by the total number of states is

$$d=\sum_{i=0}^{\mathcal{N}}\begin{pmatrix}\mathcal{N}\\ n\end{pmatrix}=2^{\mathcal{N}}\,.$$

CPT invariance implies in general a doubling of the number of states because it reverses the helicity. So for example the $\mathcal{N}$=3 multiplet in four dimensions coincides with the $\mathcal{N}$=4 multiplet if CPT invariance is required. Multiplets of particular relevance such as $\mathcal{N}$=2 with $\lambda=-\frac{1}{2}$, $\mathcal{N}$=4 with $\lambda=-1$ and $\mathcal{N}$=8 with $\lambda=-2$ are automatically CPT invariant.

Supersymmetry representations in the presence of central charges.

For $P^2=-M^2$ in the rest frame $P_\mu=(-M,0,0,0)$ the supersymmetry algebra with non vanishing central charges is given by

$$\begin{aligned}\{Q^i_\alpha,\overline{Q}_{\dot\alpha j}\} &= 2M\delta^i_j\delta_{\alpha\dot\alpha}\\ \{Q^i_\alpha,Q^j_\beta\} &= \varepsilon_{\alpha\beta}Z^{ij}\\ \{\overline{Q}_{\dot\alpha i},\overline{Q}_{\dot\beta j}\} &= -\varepsilon_{\dot\alpha\dot\beta}Z^\dagger_{ij}\,,\end{aligned}\tag{1.11}$$

with $Z^{ij}=-Z^{ji}$, $Z_{ij}=-Z^{ij}$. Since Z^{ij} commute with all the generators, by invoking Schuur's lemma there exists a basis in which they are diagonal with eigenvalues Z^{ij}. The antisymmetric $\mathcal{N}\times\mathcal{N}$ matrix with elements Z^{ij} can be put in the following form

$$Z^{ij}=U^i{}_k\tilde{Z}^{kl}(U^T)_l{}^j\,,$$

with U is a unitary matrix and

$$\begin{aligned}\tilde{Z} &= \varepsilon\otimes D && (\text{for } \mathcal{N} \text{ even})\\ \tilde{Z} &= \begin{pmatrix}\varepsilon\otimes D & 0\\ 0 & 0\end{pmatrix} && (\text{for } \mathcal{N} \text{ odd})\,,\end{aligned}$$

where $\varepsilon=\begin{pmatrix}0&1\\-1&0\end{pmatrix}$ and D is a diagonal $\frac{\mathcal{N}}{2}\times\frac{\mathcal{N}}{2}$ matrix with eigenvalues Z_R, such that $Z^*_R=Z_R$, $Z_R>0$. Considering the case with $\mathcal{N}$ even the algebra can be rewritten in a more suitable form by decomposing the indices i and j as

$$i=(a,R)\,,\qquad\qquad j=(b,S)\,,$$

with $a, b = 1, 2$ and $R, S = 1, \ldots, \frac{\mathcal{N}}{2}$. Now the algebra can be put in the form

$$\begin{aligned} \{Q_\alpha^{aR}, \overline{Q}_{\dot\beta}^{bS}\} &= 2M\delta_{\alpha\dot\alpha}\delta^{ab}\delta^{RS} \\ \{Q_\alpha^{aR}, Q_\beta^{bS}\} &= \varepsilon_{\alpha\beta}\varepsilon^{ab}\delta^{RS}Z_S \\ \{\overline{Q}_{\dot\alpha}^{aR}, \overline{Q}_{\dot\beta}^{bS}\} &= \varepsilon_{\dot\alpha\dot\beta}\varepsilon^{ab}\delta^{RS}Z_S\,, \end{aligned} \tag{1.12}$$

so that creation and annihilation operators can be defined in the following way:

$$\begin{aligned} a_\alpha^R &= \frac{1}{\sqrt{2}}\left[Q_\alpha^{1R} + \varepsilon_{\alpha\beta}(Q_\beta^{2R})^\dagger\right]\,, \qquad (a_\alpha^R)^\dagger = a_\alpha^{\dagger\,R} \\ b_\alpha^R &= \frac{1}{\sqrt{2}}\left[Q_\alpha^{1R} - \varepsilon_{\alpha\beta}(Q_\beta^{2R})^\dagger\right]\,, \qquad (b_\alpha^R)^\dagger = b_\alpha^{\dagger\,R} \end{aligned}$$

These operators satisfy the algebra of fermionic creation and annihilation operators

$$\begin{aligned} \{a_\alpha^R, a_\beta^S\} &= \{b_\alpha^R, b_\beta^S\} = \{a_\alpha^R, b_\beta^S\} = 0 \\ \{a_\alpha^R, (a_\beta^S)^\dagger\} &= \delta_{\alpha\beta}\delta^{RS}(2M + Z_S) \\ \{b_\alpha^R, (b_\beta^S)^\dagger\} &= \delta_{\alpha\beta}\delta^{RS}(2M - Z_S)\,. \end{aligned} \tag{1.13}$$

Since $\{A, A^\dagger\}$ is a positive definite operator $\forall A$, it immediately follows that $Z_R \leq 2M$. For $Z_R < 2M\ \forall R$ the structure of the multiplet is the same as in the case of no central charges. For $Z_R = 2M$ with $R = 1, \ldots, \overline{R}$ a subset of operators b cancels out and the algebra reduces to a $2(\mathcal{N} - \overline{R})$ dimensional Clifford algebra giving rise to so called *short multiplets*. States belonging to short multiplets, also referred to as *Bogomol'nyi Prasad Sommerfield (BPS) saturated states*, play a crucial rôle in the non perturbative dynamics of supersymmetric gauge theories as will be discussed later.

As an example of the previous construction consider the simplest case, namely $\mathcal{N}$=1 massive multiplet with a spin zero Clifford vacuum $|\Omega\rangle$: the states are

$$\begin{aligned} |\Omega\rangle &\qquad (\text{spin}\ \ 0) \\ (a_\alpha)^\dagger|\Omega\rangle &\qquad (\text{spin}\ \ \frac{1}{2}) \\ \tfrac{1}{\sqrt{2}}(a_\alpha)^\dagger(a_\beta)^\dagger|\Omega\rangle = -\tfrac{1}{2\sqrt{2}}\varepsilon_{\alpha\beta}(a^\gamma)^\dagger(a_\gamma)^\dagger|\Omega\rangle &\qquad (\text{spin}\ \ 0) \end{aligned}$$

In the following the discussion will focus mainly on the $\mathcal{N}$=4 massless case. For a rigid supersymmetric theory with fields of spin not larger than one the multiplet contains the following states with relative degeneracies

helicity	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1
degeneracy	1	4	6	4	1

Given the irreducible representations of the supersymmetry algebra the corresponding realizations in terms of fields can be immediately deduced. Although this was the historical pattern in the construction of supersymmetric quantum field theory models, it appears more natural to derive the structure of supermultiplets using the superfield formalism, so this will be the approach followed here.

Notice that Wigner's method here employed gives representations of "on-shell" states. It is always possible to construct sets of fields realizing such irreducible representations, but on the contrary it is usually not possible, for $\mathcal{N} > 1$, to achieve the closure of the supersymmetry algebra on a set of "off-shell" fields, *i.e.* without use of the equations of motion.

The irreducible representations of supersymmetry have automatically a well defined transformation under the group of automorphisms of the algebra (1.3). For massless representations the largest subgroup of the automorphisms group that respects helicity is the whole $\mathrm{U}(\mathcal{N})$, whereas for massive states it depends on the presence and number of central charges. In the absence of central charges, or if none of them saturates the BPS bound, the spin preserving automorphisms group is $\mathrm{USp}(2\mathcal{N})$. It reduces to $\mathrm{USp}(\mathcal{N})$ for $\mathcal{N}$ even and to $\mathrm{USp}(\mathcal{N}+1)$ for $\mathcal{N}$ odd if one central charge satisfies the condition $Z = 2M$.

1.3 $\mathcal{N}$=1 superspace

Ordinary quantum field theories are constructed in terms of field operators that are functions (actually distributions) defined on Minkowskian space-time, parametrized by the coordinates x_μ. The action is required to be invariant under Poincaré space-time symmetry as well as under transformations of an internal group G. The transformation of a generic field $f(x)$ is of the form

$$R(g)f(x) = f(\tau(g)x)\,, \tag{1.14}$$

with $R(g)$ and $\tau(g)$ suitable representations of the relevant symmetry group.

In supersymmetric models the space-time symmetry is extended by the introduction of the transformations generated by anticommuting charges Q^i_α, $\overline{Q}_{\dot{\alpha}j}$. The

natural way to implement the extended symmetry algebra is to construct the theory in terms of generalized fields, called *superfields* [14, 15], depending on the coordinates x_μ as well as on "fermionic" coordinates θ_α and $\overline{\theta}_{\dot\alpha}$

$$F = F(x_\mu, \theta_\alpha, \overline{\theta}_{\dot\alpha})\,. \tag{1.15}$$

Starting from the supersymmetry algebra a generic element of the corresponding group of transformations can be written in the form

$$g(x, \theta, \overline{\theta}) = e^{i(x^\mu P_\mu + \theta^\alpha Q_\alpha + \overline{\theta}_{\dot\alpha}\overline{Q}^{\dot\alpha})}\,.$$

The action on superfields is then defined through a differential representation of the operators P_μ, Q_α and $\overline{Q}^{\dot\alpha}$.

This construction can be carried out in a more general fashion, suitable for generalization to the case of $\mathcal{N}$-extended supersymmetry, defining the superspace parametrized by

$$z_m = (x_\mu, \theta_\alpha, \overline{\theta}_{\dot\alpha})$$

as a coset space. $\mathcal{N}$=1 superspace under consideration can be obtained as the coset SP/L, where SP is the super Poincaré group generated by (1.3) and L is the Lorentz group SO(3,1).

A coset group K=G/H can be parametrized by coordinates ξ_m, $m = 1, \ldots,$ (dim G− dim H). By writing an element of G through the exponential map

$$g = e^{i\xi^m L_m} e^{i\zeta^k H_k}\,,$$

where the generators of G split into the generators of H (H_k) and the remaining L_n, elements of the coset are obtained by taking $\zeta^k = 0$.

For the present case K=SP/L this implies that one should write

$$g = e^{i(a^\mu P_\mu + \eta^\alpha Q_\alpha + \overline{\eta}_{\dot\alpha}\overline{Q}^{\dot\alpha})} e^{\frac{1}{2} w^{\mu\nu} M_{\mu\nu}} \tag{1.16}$$

so that $w^{\mu\nu} = 0$ gives the elements of K

$$k = e^{i(x^\mu P_\mu + \theta^\alpha Q_\alpha + \overline{\theta}_{\dot\alpha}\overline{Q}^{\dot\alpha})} = e^{z^m K_m}$$

which are parametrized by $z_m = (x_\mu, \theta_\alpha, \overline{\theta}_{\dot\alpha})$; as a result $\mathcal{N}$=1 superspace is an eight dimensional manifold. Under the action of a group element $g_0 = (a_\mu, \eta_\alpha, \overline{\eta}_{\dot\alpha})$

$$g_0 \circ e^{z^m K_m} = e^{z'^m K_m} e^{\frac{1}{2} w'^{\mu\nu} M_{\mu\nu}} \equiv g'\,.$$

An explicit calculation using Becker–Hausdorff's formula and anticommutation of Q and P gives the following transformations of the coordinates z_m

$$\begin{aligned} x'^{\mu} &= x^{\mu} + a^{\mu} + i\eta^{\alpha}\sigma^{\mu}_{\alpha\dot{\alpha}}\overline{\theta}^{\dot{\alpha}} - i\theta^{\alpha}\sigma^{\mu}_{\alpha\dot{\alpha}}\overline{\eta}^{\dot{\alpha}} + w^{\mu\nu}x_{\nu} \\ \theta'^{\alpha} &= \theta^{\alpha} + \eta^{\alpha} + \frac{1}{4}w_{\mu\nu}\sigma^{\mu\nu\,\alpha}{}_{\beta}\theta_{\beta} \\ \overline{\theta}'^{\dot{\alpha}} &= \overline{\theta}^{\dot{\alpha}} + \overline{\eta}^{\dot{\alpha}} - \frac{1}{4}\overline{\theta}^{\dot{\beta}}\overline{\sigma}^{\mu\nu\,\dot{\alpha}}_{\dot{\beta}}w_{\mu\nu}\,. \end{aligned} \tag{1.17}$$

Now comparison of $(g_1 \circ g_2) \circ e^{z^m K_m}$ with $g_1 \circ (g_2 \circ e^{z^m K_m})$ allows one to prove that the generators Q_α and $\overline{Q}_{\dot{\alpha}}$ satisfy the supersymmetry algebra (1.3) for $\mathcal{N}$=1.

As already mentioned a superfield is a function of z_m

$$F(z) = F(x, \theta, \overline{\theta})\,.$$

The transformation under the group action is the generalization of (1.14) that implements $z' = \tau(g)z$ with z' given by (1.17). For infinitesimal transformations

$$\delta_g F = F(z + \delta z) - F(z) = \delta g_m X^m F(z)\,,$$

which is achieved realizing the super Poincaré generators through the differential operators $X_m = (\ell_m, \ell_\alpha, \overline{\ell}_{\dot{\alpha}}, \ell_{\mu\nu})$

$$\begin{aligned} \ell_{\mu} &= \partial_{\mu} \\ \ell_{\alpha} &= \frac{\partial}{\partial\theta^{\alpha}} + i\sigma^{\mu}_{\alpha\dot{\alpha}}\overline{\theta}^{\dot{\alpha}}\partial_{\mu} \\ \overline{\ell}_{\dot{\alpha}} &= \frac{\partial}{\partial\overline{\theta}^{\dot{\alpha}}} + i\theta^{\alpha}\sigma^{\mu}_{\alpha\dot{\alpha}}\partial_{\mu} \\ \ell_{\mu\nu} &= -(x_{\mu}\partial_{\nu} - x_{\nu}\partial_{\mu}) - \frac{1}{2}\theta^{\beta}\sigma^{\mu\nu\alpha}_{\beta}\frac{\partial}{\partial\theta^{\alpha}} + \frac{1}{2}\overline{\theta}^{\dot{\beta}}\overline{\sigma}^{\mu\nu\,\dot{\alpha}}_{\dot{\beta}}\frac{\partial}{\partial\overline{\theta}^{\dot{\alpha}}}\,. \end{aligned} \tag{1.18}$$

A calculation shows that they satisfy the super Poincaré algebra

$$\begin{aligned} &\{\ell_\alpha, \ell_\beta\} = 0\,, \qquad &&\{\ell_\alpha, \overline{\ell}_{\dot{\alpha}}\} = 2i\sigma^{\mu}_{\alpha\dot{\alpha}}\ell_{\mu}\,, \\ &[\ell_\mu, \ell_\alpha] = 0\,, \qquad &&[\ell_\alpha, \ell_{\mu\nu}] = \frac{1}{2}\sigma_{\mu\nu\alpha}{}^{\beta}\ell_{\beta}\,. \end{aligned} \tag{1.19}$$

Superfields defined in this way form linear representations of supersymmetry. Since supersymmetry acts on superfields through linear differential operators it follows that the product of superfields is a new superfield.

The objects considered up to now are scalar superfields [2]; more general superfields transforming in a non trivial way under Lorentz transformations can be considered as well. In general a superfield carrying a set of Lorentz indices encoded in a single label p transforms as

$$R(g)f_p(z) = D_p{}^q \left(e^{-\frac{1}{2}w^{\mu\nu}M_{\mu\nu}}\right) F_q(\tau(g)z)\,,$$

where D^q_p is a suitable representation of the Lorentz group.

Physical fields directly related to the states considered in section 1.2 can be obtained from the superfields by expanding in the θ and $\overline{\theta}$ variables. Since θ_α and $\overline{\theta}_{\dot\alpha}$ are Weyl spinors the expansion reduces to a polynomial. For a generic superfield

$$\begin{aligned} F(x,\theta,\overline{\theta}) &= f(x) + \theta^\alpha\chi_\alpha(x) + \overline{\theta}_{\dot\alpha}\overline{\psi}^{\dot\alpha}(x) + \theta^\alpha\theta_\alpha g(x) + \overline{\theta}_{\dot\alpha}\overline{\theta}^{\dot\alpha}h(x) + \\ &+\theta^\alpha\sigma^\mu_{\alpha\dot\alpha}\overline{\theta}^{\dot\alpha}r_\mu(x) + \theta^\alpha\theta_\alpha\overline{\theta}_{\dot\alpha}\overline{\lambda}^{\dot\alpha}(x) + \overline{\theta}_{\dot\alpha}\overline{\theta}^{\dot\alpha}\theta^\alpha\xi_\alpha(x) + \theta^\alpha\theta_\alpha\overline{\theta}_{\dot\alpha}\overline{\theta}^{\dot\alpha}s(x)\,. \end{aligned} \tag{1.20}$$

The fields $f(x)$, $\chi(x)$, $\overline{\psi}(x)$, $g(x)$, $h(x)$, $r_\mu(x)$, $\overline{\lambda}(x)$, $\xi(x)$ and $s(x)$ constitute a supermultiplet and their transformation law is derived from the transformation of the superfield by defining

$$\begin{aligned} \delta_\eta F(x,\theta,\overline{\theta}) &= \delta_\eta f(x) + \theta^\alpha\delta_\eta\chi_\alpha(x) + \overline{\theta}_{\dot\alpha}\delta_\eta\overline{\psi}^{\dot\alpha}(x) + \theta^\alpha\theta_\alpha\delta_\eta g(x) + \overline{\theta}_{\dot\alpha}\overline{\theta}^{\dot\alpha}\delta_\eta h(x) + \\ &+\theta^\alpha\sigma^\mu_{\alpha\dot\alpha}\overline{\theta}^{\dot\alpha}\delta_\eta r^\mu(x) + \theta^\alpha\theta_\alpha\overline{\theta}_{\dot\alpha}\delta_\eta\overline{\lambda}^{\dot\alpha}(x) + \overline{\theta}_{\dot\alpha}\overline{\theta}^{\dot\alpha}\theta^\alpha\delta_\eta\xi_\alpha(x) + \theta^\alpha\theta_\alpha\overline{\theta}_{\dot\alpha}\overline{\theta}^{\dot\alpha}\delta_\eta s(x) \end{aligned} \tag{1.21}$$

and matching terms with the same powers of θ and $\overline{\theta}$.

Linear representations of supersymmetry constructed in this way are in general reducible; the irreducible representations studied in section 1.2 are in correspondence with constrained superfields. All the representations of rigid $\mathcal{N}$=1 supersymmetry in four dimensions are realized in terms of two basic superfields, namely *chiral superfields* and *vector superfields*.

From now on the differential operators ℓ_α and $\overline{\ell}_{\dot\alpha}$ will be denoted as Q_α and $\overline{Q}_{\dot\alpha}$ like the abstract generators since they satisfy the same algebra. The operators Q_α and $\overline{Q}_{\dot\alpha}$ realize the left regular representation of supersymmetry on superfields, it is useful to analogously define the right regular representation realized by the differential operators

$$D_\alpha = \frac{\partial}{\partial\theta^\alpha} + i\sigma^\mu_{\alpha\dot\alpha}\overline{\theta}^{\dot\alpha}\partial_\mu\,, \qquad \overline{D}_{\dot\alpha} = -\frac{\partial}{\partial\overline{\theta}^{\dot\alpha}} - i\theta^\alpha\sigma^\mu_{\alpha\dot\alpha}\partial_\mu \tag{1.22}$$

[2] From now on scalar superfields will be referred to simply as superfields.

satisfying

$$\begin{aligned} \{D_\alpha, \overline{D}_{\dot\alpha}\} &= -2i\sigma^\mu_{\alpha\dot\alpha}\partial_\mu \\ \{D_\alpha, D_\beta\} &= \{\overline{D}_{\dot\alpha}, \overline{D}_{\dot\beta}\} = 0\,. \end{aligned}$$

D_α and $\overline{D}_{\dot\alpha}$ are usually referred to as superspace covariant derivatives, actually they can be obtained from the general expression for the covariant derivative in the coset space SP/L: the fact that D_α and $\overline{D}_{\dot\alpha}$ do not reduce to ∂_α and $\overline{\partial}_{\dot\alpha}$ despite the flatness of the $\mathcal{N}$=1 superspace is a consequence of the presence of a non vanishing torsion.

Chiral superfields.

Chiral superfields are characterized by the constraint

$$\overline{D}_{\dot\alpha}\Phi(x,\theta,\overline{\theta}) = 0\,,$$

which takes a particularly simple form expressing $\Phi(x,\theta,\overline{\theta})$, D and $\overline{D}$ in terms of the variable $y^\mu = x^\mu + i\theta\sigma^\mu\overline{\theta}$. As functions of y, θ and $\overline{\theta}$ one gets

$$\begin{aligned} D_\alpha &= \frac{\partial}{\partial\theta^\alpha} + 2i\sigma^\mu_{\alpha\dot\alpha}\overline{\theta}^{\dot\alpha}\frac{\partial}{\partial y^\mu} \\ \overline{D}_{\dot\alpha} &= -\frac{\partial}{\partial\overline{\theta}^{\dot\alpha}}\,, \end{aligned}$$

so that the general chiral superfield assumes the form

$$\Phi(y,\theta) = \phi(y) + \sqrt{2}\theta\psi(y) + \theta\theta F(y)\,.$$

Substituting $y^\mu = x^\mu + i\theta\sigma^\mu\overline{\theta}$ and expanding in θ and $\overline{\theta}$ yields

$$\begin{aligned} \Phi(x,\theta,\overline{\theta}) &= \phi(x) + \sqrt{2}\theta\psi(x) + i\theta\sigma^\mu\overline{\theta}\partial_\mu\phi(x) + \\ &\quad +\theta\theta F(x) - \frac{i}{\sqrt{2}}\theta\theta\partial_\mu\psi(x)\sigma^\mu\overline{\theta} + \frac{1}{4}\theta\theta\overline{\theta}\overline{\theta}\Box\phi(x)\,. \end{aligned} \tag{1.23}$$

Analogously antichiral superfields are defined by

$$D_\alpha\Phi^\dagger(x,\theta,\overline{\theta}) = 0\,.$$

The natural variable is $y^{\dagger\,\mu} = x^\mu - i\theta\sigma^\mu\overline{\theta}$; in terms of $y^{\dagger\,\mu}$

$$\Phi^\dagger = \Phi^\dagger(y^\dagger,\overline{\theta}) = \phi^*(y^\dagger) + \sqrt{2}\overline{\theta\psi}(y^\dagger) + \overline{\theta\theta}F^*(y^\dagger)\,.$$

The field components associated to chiral and antichiral multiplets are of two complex scalars (ϕ, F) and one Weyl spinor (ψ) corresponding to twice the number of states in the massive on-shell $\mathcal{N}$=1 multiplet of section 1.2.

Products of (anti) chiral superfields are still (anti) chiral superfields. For example

$$\begin{aligned}\Phi_i\Phi_j &= \phi_i(y)\phi_j(y) + \sqrt{2}\theta[\psi_i(y)\phi_j(y) + \phi_i(y)\psi_j(y)] + \\ &\quad +\theta\theta[\phi_i(y)F_j(y) + \phi_j(y)F_i(y) - \psi_i(y)\psi_j(y)]\end{aligned}$$

The product $\Phi^\dagger\Phi$ is neither chiral nor antichiral. Moreover since $D^3 = \overline{D}^3 = 0$ it immediately follows that for a generic superfield F

$$\begin{aligned}&\overline{D}^2 F = \Phi && \text{is chiral} \\ &D^2 F = \Phi^\dagger && \text{is antichiral}\,.\end{aligned}$$

Notice that the double constraint $D_\alpha\Phi = \overline{D}_{\dot\alpha}\Phi = 0$ implies that Φ is constant.

Vector superfields.

Vector superfields obey the constraint

$$V^\dagger(x,\theta,\overline{\theta}) = V(x,\theta,\overline{\theta})\,,$$

which gives rise to the component expansion

$$\begin{aligned}&V(x,\theta,\overline{\theta}) = C(x) + i\theta\chi(x) - i\overline{\theta}\overline{\chi}(x) + \frac{i}{\sqrt{2}}\theta\theta S(x) + \\ &-\frac{i}{\sqrt{2}}\overline{\theta\theta}S^\dagger(x) - \theta\sigma^\mu\overline{\theta}A_\mu(x) + i\theta\theta\overline{\theta}\left[\overline{\lambda}(x) + \frac{i}{2}\overline{\sigma}^\mu\partial_\mu\chi(x)\right] + \\ &-i\overline{\theta\theta}\theta\left[\lambda(x) + \frac{i}{2}\sigma^\mu\partial_\mu\overline{\chi}(x)\right] + \frac{1}{2}\theta\theta\overline{\theta\theta}\left[D(x) + \frac{1}{2}\Box C(x)\right]. \qquad (1.24)\end{aligned}$$

The supermultiplet contains two real scalars (C, D), one complex scalar (S), four Weyl spinors (χ, $\overline{\chi}$, λ, $\overline{\lambda}$) and one real vector (A_μ). This is twice the content of the massive $\mathcal{N}$=1 multiplet obtained starting with a Clifford vacuum of spin $\frac{1}{2}$ and also of the massless CPT invariant multiplet with minimum helicity $\lambda = -1$.

Since the vector multiplet contains a vector field $A_\mu(x)$ it is the basic ingredient for the construction of supersymmetric gauge theories. The supersymmetric generalization of an Abelian gauge transformation [16, 17] is

$$V \longrightarrow V + \Phi + \Phi^\dagger\,,$$

with

$$\begin{aligned}\Phi+\Phi^\dagger &= \phi+\phi^*+\sqrt{2}(\theta\psi+\overline{\theta\psi})+\theta\theta F+\overline{\theta\theta}F^*+i\theta\sigma^\mu\overline{\theta}\partial_\mu(\phi-\phi^*)+\\ &\quad+\frac{1}{\sqrt{2}}\theta\theta\overline{\theta}\overline{\sigma}^\mu\partial_\mu\psi+\frac{1}{\sqrt{2}}\overline{\theta\theta}\sigma^\mu\theta\partial_\mu\overline{\psi}+\frac{1}{4}\theta\theta\overline{\theta\theta}\Box(\phi+\phi^*)\,.\end{aligned}$$

For the components one gets

$$\begin{aligned}C &\longrightarrow C+\phi+\phi^*\\ \chi &\longrightarrow \chi-i\sqrt{2}\psi\\ S &\longrightarrow S-i\sqrt{2}F\\ A_\mu &\longrightarrow A_\mu-i\partial_\mu(\phi-\phi^*)\\ \lambda &\longrightarrow \lambda\\ D &\longrightarrow D\,.\end{aligned}$$

Notice that the transformations for the lowest components C, χ and S are purely algebraic (no derivatives involved), so they can always be used to put these fields to zero. This particular choice of gauge is known as *Wess–Zumino* (WZ) *gauge*. Subtleties related to the choice of the Wess–Zumino gauge will be discussed in chapter 3. It is important to note that in this gauge one has

$$\begin{aligned}V(x,\theta,\overline{\theta}) &= -\theta\sigma^\mu\overline{\theta}A_\mu(x)+i\theta\theta\overline{\theta\lambda}(x)-i\overline{\theta\theta}\theta\lambda(x)+\frac{1}{2}\theta\theta\overline{\theta\theta}D(x)\\ V^2(x,\theta,\overline{\theta}) &= -\frac{1}{2}\theta\theta\overline{\theta\theta}A_\mu(x)A^\mu(x)\\ V^n(x,\theta,\overline{\theta}) &= 0\,,\qquad \forall n\geq 3\,.\end{aligned}$$

Fixing the Wess–Zumino gauge still leaves an arbitrary Abelian gauge freedom on A_μ.

From the vector superfield it is straightforward to construct a superfield containing the field strength for the field A_μ: it is defined as

$$W_\alpha(x,\theta,\overline{\theta})=-\frac{1}{4}\overline{DD}D_\alpha V(x,\theta,\overline{\theta}) \tag{1.25}$$

and is a spinor superfield. Besides W_α it is useful to introduce

$$\overline{W}_\alpha(x,\theta,\overline{\theta})=-\frac{1}{4}DD\overline{D}_{\dot\alpha}V(x,\theta,\overline{\theta})\,. \tag{1.26}$$

They are respectively chiral and antichiral:

$$\overline{D}_{\dot\alpha}W_\alpha=0\,,\qquad\qquad D_\alpha\overline{W}_{\dot\alpha}=0$$

and gauge invariant

$$W[V+\Phi+\Phi^\dagger] = W[V]\,.$$

Gauge invariance of W_α allows to compute its component expansion in the Wess–Zumino gauge; in terms of y^μ one gets

$$\begin{aligned} W_\alpha(y,\theta) &= -i\lambda(y) + \left[\delta_\alpha{}^\beta D - \frac{i}{2}(\sigma^\mu\sigma^\nu)_\alpha{}^\beta(\partial_\mu A_\nu(y) - \partial_\nu A_\mu(y))\right]\theta_\beta + \\ &\quad +\theta\theta\sigma^\mu_{\alpha\dot\alpha}\partial_\mu\overline{\lambda}^{\dot\alpha}(y) = \\ &= -i\lambda(y) + \left[\delta_\alpha{}^\beta D - \frac{i}{2}(\sigma^\mu\sigma^\nu)_\alpha{}^\beta F_{\mu\nu}(y)\right]\theta_\beta + \theta\theta\sigma^\mu_{\alpha\dot\alpha}\partial_\mu\overline{\lambda}^{\dot\alpha}(y) \end{aligned} \tag{1.27}$$

and analogously for $\overline{W}_\alpha$.

1.4 $\mathcal{N}$=1 supersymmetric models

The superfield formalism allows a straightforward construction of supersymmetric actions. The supersymmetry transformation of the $(\overline{\theta\theta})$ $\theta\theta$ component (F-component) of a (anti) chiral superfield is a total derivative, as can be deduced by specializing (1.21) to the case of a chiral superfield, so that an action of the form [3]

$$S = \int d^4x \left[\int d^2\theta\,\Phi + \int d^2\overline{\theta}\,\Phi^\dagger\right]$$

is automatically supersymmetric for an arbitrary chiral Φ.

The same situation holds for the $\theta\theta\overline{\theta\theta}$ component (D-component) of a vector superfield yielding the supersymmetric expression

$$S = \int d^4x \int d^4\theta\, V\,,$$

for all V's such that $V^\dagger = V$.

Requirement of invariance under the group of automorphisms allows to further restrict the possible couplings in the action for supersymmetric models; in this context the symmetry under transformations of the group of automorphisms is referred to as R-*symmetry*.

[3] The properties of Grassmannian integration and the conventions here employed are summarized in appendix A.

1.4.1 Supersymmetric chiral theories: Wess–Zumino model

The action for the Wess–Zumino model [18] is constructed in terms of only chiral superfields and reads

$$\begin{aligned} S &= \int d^4x \left\{ d^4\theta\, \Phi_i^\dagger \Phi_i + \left[\int d^2\theta \left(\frac{1}{2} m_{ij} \Phi_i \Phi_j + \right.\right.\right. \\ &\left.\left.\left. + \frac{1}{3} g_{ijk} \Phi_i \Phi_j \Phi_k + \lambda_i \Phi_i \right) + \text{h.c.} \right] \right\}. \end{aligned} \tag{1.28}$$

In terms of component fields it becomes

$$\begin{aligned} S &= \int d^4x \left\{ -\partial_\mu \phi_i^* \partial^\mu \phi_i - i\overline{\psi}_i \overline{\sigma}^\mu \partial_\mu \psi_i + F_i^* F_i + \left[m_{ij} (\phi_i F_j - \frac{1}{2} \psi_i \psi_j) + \right.\right. \\ &\left.\left. + g_{ijk} (\phi_i \phi_j F_k - \psi_i \psi_j \phi_k) + \lambda_i F_i + \text{h.c.} \right] \right\}. \end{aligned} \tag{1.29}$$

The action of the Wess–Zumino model is invariant under the supersymmetry transformations

$$\begin{aligned} \delta_\eta \phi_i &= \sqrt{2} \eta \psi_i \\ \delta_\eta \psi_i &= i\sqrt{2} \sigma^\mu \overline{\eta} \partial_\mu \phi_i + \sqrt{2} \eta F_i \\ \delta_\eta F_i &= i\sqrt{2} \overline{\eta \sigma}^\mu \partial_\mu \psi_i \end{aligned}$$

The action (1.29) does not contain derivatives of the fields F_i. They are *auxiliary fields*: their equations of motion are algebraic and can be used to eliminate them from the action. Solving the equations for F_i

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial F_k^*} &= F_k + \lambda_k^* + m_{ik}^* \phi_i^* + g_{ijk}^* \phi_i^* \phi_j^* = 0 \\ \frac{\partial \mathcal{L}}{\partial F_k} &= F_k^* + \lambda_k + m_{ik} \phi_i + g_{ijk} \phi_i \phi_j = 0\,, \end{aligned}$$

yields the Lagrangian

$$\begin{aligned} \mathcal{L} &= -i\overline{\psi}_i \overline{\sigma}^\mu \partial_\mu \psi_i - \partial^\mu \phi^* \partial_\mu \phi - \frac{1}{2} m_{ij} \psi_i \psi_j - \frac{1}{2} m_{ij}^* \overline{\psi}_i \overline{\psi}_j + \\ &\quad - g_{ijk} \psi_i \psi_j \phi_k - g_{ijk}^* \overline{\psi}_i \overline{\psi}_j \phi_k^* - \mathcal{V}(\phi_i, \phi_i^*)\,, \end{aligned}$$

where the potential $\mathcal{V}$ is given (for $\lambda_i = 0$) by the expression

$$\mathcal{V} = (F_k^* F_k) = (m_{ik} \phi_i + g_{ijk} \phi_i \phi_j)(m_{mk}^* \phi_m^* + g_{mnk}^* \phi_m^* \phi_n^*)\,. \tag{1.30}$$

This scalar potential in particular contains a ϕ^4 interaction and the mass terms for the scalars ϕ_k; actually the Wess–Zumino model is the most general renormalizable supersymmetric theory involving only chiral superfields.

The action (1.29) is the most general renormalizable supersymmetric action that can be constructed in terms of only chiral superfields. More generally one can consider an invariant action of the form

$$S = \int d^4x \left\{ \int d^4\theta \, \Phi_i^\dagger \Phi_i + \left[\int d^2\theta \, \mathrm{W}(\Phi) + \mathrm{h.c.} \right] \right\} ,$$

where $\mathrm{W}(\Phi)$ is a holomorphic function of Φ called *superpotential*. In this case the scalar potential becomes

$$\mathcal{V}(\Phi) = \left| \frac{\partial \mathrm{W}(\Phi)}{\partial \Phi} \right|^2 .$$

The Wess–Zumino model can be constructed in terms of unconstrained superfields using the property $D^3 = \overline{D}^3 = 0$ and defining

$$\Phi = \overline{D}^2 \Psi \qquad \Phi^\dagger = D^2 \Psi^\dagger ,$$

The unconstrained action that reduces to (1.28), for the case of one single field Φ, is

$$\begin{aligned} S \;\; = \;\; & \int d^4x d^4\theta \, \Big\{ (D^2\Psi^\dagger)(\overline{D}^2\Psi) - 2m(\Psi \overline{D}^2 \Psi + \Psi^\dagger D^2 \Psi + \\ & - \frac{4g}{3!} \left[\Psi (\overline{D}^2 \Psi)^2 + \Psi^\dagger (D^2 \Psi^\dagger)^2 \right] \Big\} . \end{aligned} \tag{1.31}$$

The equivalence follows using the identity

$$\int d^4x \int d^2\theta \, \Psi = \int d^4x \int d^4\theta \, \left(-\frac{1}{4} \overline{D}^2 \right) \Psi .$$

1.4.2 Supersymmetric gauge theories

Gauge invariant interactions can be constructed in terms of vector superfields and more precisely in terms of the superfield W_α defined in section 1.3, which contains the gauge field strength among its components.

Given a chiral superfield Φ and a vector one V an Abelian gauge theory is obtained as follows. The gauge transformation for the chiral superfield is

$$\begin{aligned} \Phi_k' &= e^{-iq_k \Lambda} \Phi_k , \qquad \text{with} \qquad \overline{D}_{\dot\alpha} \Lambda = 0 \\ \Phi_k^{\dagger\prime} &= e^{iq_k \Lambda^\dagger} \Phi_k^\dagger , \qquad \text{with} \qquad D_\alpha \Lambda^\dagger = 0 , \end{aligned}$$

where q_k denotes the charge of the superfield Φ_k. The gauge parameter must be a chiral superfield Λ. Supplementing these transformations with the following one for the vector superfield V

$$V' = V + i(\Lambda - \Lambda^\dagger)\,,$$

an invariant action can be written in the form

$$\begin{aligned} S \;=\; & \int d^4x \left\{ \int d^2\theta\, \frac{1}{4} W^\alpha W_\alpha + \int d^2\overline{\theta}\, \frac{1}{4} \overline{W}_{\dot\alpha} \overline{W}^{\dot\alpha} + \int d^4\theta\, \Phi_i^\dagger e^{q_i V} \Phi_i + \right. \\ & \left. + \left[\int d^2\theta\, \left(\frac{1}{2} m_{ij} \Phi_i \Phi_j + \frac{1}{3} g_{ijk} \Phi_i \Phi_j \Phi_k \right) + \text{ h.c.} \right] \right\} \end{aligned} \tag{1.32}$$

Invariance under the above defined transformations is immediately verified. Notice that the action (1.32) is non-polynomial: this a general feature of supersymmetric gauge theories and is a consequence of the adimensionality of the vector superfield V. Choosing the Wess–Zumino gauge reduces (1.32) to a polynomial form since it implies $V^3 = 0$. This choice allows one to prove the renormalizability of the model.

The vector superfield V contains the gauge dependent components C, χ and S that can be put to zero by choosing the Wess–Zumino gauge as well as the real scalar D. The latter is an auxiliary field since the corresponding equations of motion obtained from (1.32) are purely algebraic.

This construction can be generalized to the non Abelian case [19, 20]. For a generic non Abelian compact gauge group G the fields Φ, $\Phi^\dagger$ and V are matrices.

$$(\Phi)_{ij} = T^a_{ij} \Phi_a\,, \qquad \left(\Phi^\dagger\right)_{ij} = T^a_{ij} \Phi^\dagger_a\,, \qquad V_{ij} = T^a_{ij} V_a\,,$$

with generators T^a of G satisfying

$$\mathrm{tr}(T^a T^b) = C(\mathbf{r}) \delta^{ab} \qquad\qquad [T^a, T^b] = i f^{ab}{}_c T^c\,,$$

$C(\mathbf{r}) = d_{\mathbf{r}}$ being the Dynkin index of the representation $\mathbf{r}$.

The gauge transformation takes the form

$$\Phi' = e^{-i\Lambda} \Phi\,, \qquad\qquad \Phi^{\dagger\prime} = \Phi^\dagger e^{i\Lambda^\dagger}$$

and for the vector superfield

$$V \longrightarrow V' : \quad e^{V'} = e^{-i\Lambda^\dagger} e^V e^{i\Lambda}\,. \tag{1.33}$$

It can be proved that in this case as well it is possible to choose the Wess–Zumino gauge in which $V^3 = 0$.

For infinitesimal gauge transformations use of the Becker–Hausdorff's formula allows to write

$$\delta V = V' - V = i\mathcal{L}_{V/2}\left[(\Lambda + \Lambda^\dagger + \coth(\mathcal{L}_{V/2})(\Lambda - \Lambda^\dagger)\right] , \tag{1.34}$$

where $\mathcal{L}_A(B) = [A, B]$ is the Lie derivative and (1.34) is a compact form to be understood in terms of the power expansion of $\coth(\mathcal{L})$.

The non Abelian generalization of the field strength superfield W_α is

$$W_\alpha = -\frac{1}{4}\overline{DD}e^{-V}D_\alpha e^V .$$

In the Wess–Zumino gauge the non Abelian W_α has the same form as in (1.27) with

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + ig[A_\mu, A_\nu] .$$

The transformation law of W_α induced by (1.33) is easily obtained and reads

$$W_\alpha \longrightarrow W'_\alpha = e^{-i\Lambda}W_\alpha e^{i\Lambda} ,$$

so that while in the Abelian case W_α was gauge invariant for a non Abelian gauge group G it turns out to be covariant.

The most general renormalizable action for interacting chiral and vector superfields is therefore

$$\begin{aligned} S \;=\; & \int d^4x\, \frac{1}{d_{\mathbf{r}}}\mathrm{tr}\left\{\frac{1}{16g^2}\left[\int d^2\theta\, W^\alpha W_\alpha + \int d^2\overline{\theta}\,\overline{W}_{\dot\alpha}\overline{W}^{\dot\alpha}\right] + \int d^4\theta\, e^{-V}\Phi^\dagger e^V\Phi + \right. \\ & \left. + \left[\int d^2\theta\, \left(\frac{1}{2}m_{ij}\Phi_i\Phi_j + \frac{1}{3}g_{ijk}\Phi_i\Phi_j\Phi_k\right) + \text{h.c.}\right]\right\} . \end{aligned} \tag{1.35}$$

The corresponding component field expansion in the Wess–Zumino gauge for the case $m_{ij} = 0,\ g_{ijk} = 0$ is

$$\begin{aligned} S = \int d^4x\, & \left\{-\frac{1}{4}F^a_{\mu\nu}F^{a\mu\nu} - i\overline{\lambda}^a\overline{\sigma}^\mu\mathcal{D}_\mu\lambda^a + \frac{1}{2}D^aD^a - \mathcal{D}_\mu\phi^\dagger\mathcal{D}^\mu\phi + \right. \\ & \left. -i\overline{\psi}\overline{\sigma}^\mu\mathcal{D}_\mu\psi + F^\dagger F + i\sqrt{2}g\left(\phi^\dagger T^a\psi\lambda^a - \overline{\lambda}^a T^a\phi\overline{\psi}\right) + gD^a\phi^\dagger T^a\phi\right\} , \end{aligned} \tag{1.36}$$

where the correct factors of the coupling constant g have been recovered substituting in (1.35) $V \longrightarrow 2gV$. In (1.36) $\mathcal{D}_\mu$ denotes the ordinary covariant derivative

$$\begin{aligned} \mathcal{D}_\mu\phi &= \partial_\mu\phi + igA^a_\mu T^a\phi \\ \mathcal{D}_\mu\psi &= \partial_\mu\psi + igA^a_\mu T^a\psi \\ \mathcal{D}_\mu\lambda^a &= \partial_\mu\lambda^a + igf^a{}_{bc}A^b_\mu\lambda^c . \end{aligned}$$

Eliminating the auxiliary fields F and D through the equations of motion produces the scalar potential given by (1.30) plus the so called "D-term"

$$\mathcal{V}_D = \frac{1}{8} g^2 \left([\phi^\dagger, \phi] \right)^2 .$$

The action (1.36) can be proved to be invariant under the transformations

$$\begin{aligned}
\delta_\xi \phi &= \sqrt{2}\xi\psi \\
\delta_\xi \psi &= i\sqrt{2}\sigma^\mu \overline{\xi} \mathcal{D}_\mu \phi + \sqrt{2}\xi F \\
\delta_\xi F &= i\sqrt{2}\sigma^\mu \overline{\xi} \mathcal{D}_\mu \psi + 2igT^a \phi \overline{\xi\lambda}^a \\
\delta_\xi A^a_\mu &= -i\overline{\lambda}^a \overline{\sigma}_\mu \xi + i\overline{\xi}\overline{\sigma}_\mu \lambda^a \\
\delta_\xi \lambda^a &= \sigma^{\mu\nu}\xi F^a_{\mu\nu} + i\xi D^a \\
\delta_\xi D^a &= -\xi\sigma^\mu \mathcal{D}_\mu \overline{\lambda}^a - \mathcal{D}_\mu \lambda^a \sigma^\mu \overline{\xi} .
\end{aligned} \tag{1.37}$$

The Wess–Zumino condition is not supersymmetric so that it not only (partially) breaks gauge invariance, but it also breaks supersymmetry, that is recovered through a non linear realization given by equations (1.37). These transformations are more precisely a mixture of supersymmetry and the residual non Abelian gauge symmetry. The choice of the Wess–Zumino gauge leads to subtleties at the quantum level, as will be discussed in chapter 3.

The super-gauge invariant action (1.35) can be generalized by introducing a complexified coupling constant

$$\tau = \frac{\theta}{2\pi} + \frac{4\pi i}{g^2}$$

obtaining, in the pure super Yang–Mills case, the analytic expression

$$\begin{aligned}
S &= \frac{1}{8\pi} \mathrm{Im} \left[\tau \int d^4x d^2\theta \, \mathrm{tr} W^\alpha W_\alpha \right] = \\
&= \frac{1}{g^2} \int d^4x \, \mathrm{tr} \left[-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - i\lambda\sigma^\mu \mathcal{D}_\mu \overline{\lambda} + \frac{1}{2} D^2 \right] + \\
&\quad -\frac{\theta}{32\pi^2} \int d^4x \, \mathrm{tr} F_{\mu\nu} \tilde{F}^{\mu\nu} ,
\end{aligned}$$

where $\tilde{F}^{\mu\nu} = \frac{1}{2}\varepsilon^{\mu\nu\rho\sigma} F_{\rho\sigma}$.

1.5 Theories of extended supersymmetry

The most straightforward way to construct multiplets for theories with extended ($\mathcal{N} > 1$) supersymmetry is to assemble together $\mathcal{N}$=1 multiplets in such a way as to

obtain a field content corresponding to the representations of section 1.2. It turns out that in general this program can be achieved only on-shell; in other words it is not possible to realize $\mathcal{N}$-extended supersymmetry in a closed form through an off-shell multiplet of fields. With the exception of $\mathcal{N}$=2 supersymmetric Yang–Mills theories closure of the algebra without imposing the equations of motion would require an infinite set of auxiliary fields. This problem is related to the presence of central charges in the supersymmetry algebra for $\mathcal{N} > 1$.

As in the $\mathcal{N}$=1 case the notion of superspace will be introduced to start with and then explicit realizations in specific models will be considered.

1.5.1 Extended superspace: a brief survey

The aim of this section is to give a brief account of ideas beyond the superspace formulation for theories with extended supersymmetry, no claim of completeness is made. In the following extended superspace will not be used in explicit calculations, but it will rather be invoked to justify some statements. A detailed and self-contained review of extended superspace is given in [21].

The general construction of superspace as a coset space introduced in section 1.3 can be generalized to define extended superspace as the coset K=SP*/L, where SP* is the extended super Poincaré group generated by the general algebra (1.1), (1.3) and L is the Lorentz group SO(3,1). The generic element of SP* can be written

$$g = e^{i(a^\mu P_\mu + \eta_i^\alpha Q_\alpha^i + \overline{\eta}_{\dot\alpha}^i \overline{Q}_i^{\dot\alpha} + b^r Z_r + \frac{1}{2} w_{\mu\nu} M^{\mu\nu})} ,$$

where $Z_{ij} = \Omega_{ij}^r Z_r$ are the central charges of the supersymmetry algebra and in this case there are $\mathcal{N}$ anticommuting generators Q_α^i. Elements of the coset are obtained setting $w_{\mu\nu} = 0$ and can be expressed in the form

$$k = e^{i(x^\mu P_\mu + \theta_i^\alpha Q_\alpha^i + \overline{\theta}_{\dot\alpha}^i \overline{Q}_i^{\dot\alpha} + \zeta^r Z_r)} . \tag{1.38}$$

Consequently they are parametrized by the enlarged set of coordinates

$$z_m = (x_\mu, \theta_\alpha^i, \overline{\theta}_{\dot\alpha}^j, \zeta_r) . \tag{1.39}$$

Repeating the steps of section 1.3 allows to obtain the following supersymmetric

coordinate transformations

$$\begin{aligned} x'^{\mu} &= x^{\mu} + i\eta_i^{\alpha}\sigma^{\mu}_{\alpha\dot{\alpha}}\overline{\theta}^{\dot{\alpha}i} - i\theta_i^{\alpha}\sigma^{\mu}_{\alpha\dot{\alpha}}\overline{\eta}^{\dot{\alpha}i} \\ \theta_i'^{\alpha} &= \theta_i^{\alpha} + \eta_i^{\alpha} \\ \overline{\theta}'^{\dot{\alpha}j} &= \overline{\theta}^{\dot{\alpha}j} + \overline{\eta}^{\dot{\alpha}j} \\ \zeta'^{r} &= \zeta^r + i\eta_i^{\alpha}\theta_{\alpha j}(\Omega^r)^{ij} + i\overline{\eta}^i_{\dot{\alpha}}\overline{\theta}^{\dot{\alpha}j}(\Omega^r)_{ij}\,. \end{aligned} \tag{1.40}$$

Under the action of the central charges x^{μ}, θ^i_{α} and $\overline{\theta}_{\dot{\alpha}j}$ do not transform while

$$\zeta'^{r} = \zeta^r + b^r\,. \tag{1.41}$$

Superfields are functions defined on the manifold parametrized by z_m

$$F = F(x_{\mu}, \theta^i_{\alpha}, \overline{\theta}^j_{\dot{\alpha}}, \zeta_r)\,. \tag{1.42}$$

The action of the super Poincaré group on superfields is defined through

$$R(g)\,F(z) = F(z')$$

where z' is given in (1.40) and $R(g)$ acts as the left regular representation.

Differential operators generating the infinitesimal transformations now read

$$\begin{aligned} \ell_{\mu} &= \partial_{\mu} \\ \ell^i_{\alpha} &= \frac{\partial}{\partial\theta^{\alpha}_i} + i\sigma^{\mu}_{\alpha\dot{\alpha}}\overline{\theta}^{\dot{\alpha}i}\partial_{\mu} + i(\Omega^r)^{ij}\theta_{\alpha j}\frac{\partial}{\partial\zeta^r} \\ \overline{\ell}_{\dot{\alpha}i} &= \frac{\partial}{\partial\overline{\theta}^{\dot{\alpha}i}} + i\theta^{\alpha}_i\sigma^{\mu}_{\alpha\dot{\alpha}}\partial_{\mu} + i\overline{\theta}^j_{\alpha}(\Omega^r)_{ij}\frac{\partial}{\partial\zeta^r} \\ \ell^r &= \frac{\partial}{\partial\zeta_r} \\ \ell_{\mu\nu} &= -(x_{\mu}\partial_{\nu} - x_{\nu}\partial_{\mu}) - \frac{1}{2}\theta^{\beta}\sigma^{\mu\nu\,\alpha}_{\beta}\frac{\partial}{\partial\theta^{\alpha}} + \frac{1}{2}\overline{\theta}^{\dot{\beta}}\overline{\sigma}^{\mu\nu\,\dot{\alpha}}_{\dot{\beta}}\frac{\partial}{\partial\overline{\theta}^{\dot{\alpha}}} \\ \ell_a &= -\theta^{\alpha}_i B^i_{aj}\frac{\partial}{\partial\theta^{\alpha}_j} - \overline{\theta}^{\dot{\alpha}i}\left(B^i_{aj}\right)^*\frac{\partial}{\partial\overline{\theta}^{\dot{\alpha}}_j}\,. \end{aligned}$$

Relation to component field multiplets is established by Taylor expanding in the θ, $\overline{\theta}$ as well as in the ζ coordinates. Notice that the ζ's are "bosonic" coordinates, so that the expansion is a truly infinite series

$$F(x,\theta,\overline{\theta},\zeta) = F^{(0)}(x,\theta,\overline{\theta}) + F^{(1)}_{(r)}(x,\theta,\overline{\theta})\,\zeta^{(r)} + \dots\,, \tag{1.43}$$

where

$$\begin{aligned} F^{(0)}(x,\theta,\overline{\theta}) &= F(x,\theta,\overline{\theta},0) \\ F^{(1)}_{(r)}(x,\theta,\overline{\theta}) &= \frac{\partial}{\partial\zeta^r}\, F(x,\theta,\overline{\theta},\zeta)\Big|_{\zeta=0}\,. \end{aligned}$$

"Super-covariant" derivatives generalizing those defined in section 1.3 are given by

$$\begin{aligned} D_\mu &= \partial_\mu \\ D^i_\alpha &= \frac{\partial}{\partial\theta^\alpha_i} + i\sigma^\mu_{\alpha\dot\alpha}\overline{\theta}^{\dot\alpha i}\partial_\mu - i(\Omega^r)^{ij}\theta_{\alpha j}\frac{\partial}{\partial\zeta^r} \\ \overline{D}_{\dot\alpha i} &= -\frac{\partial}{\partial\overline{\theta}^{\dot\alpha i}} - i\theta^\alpha_i\sigma^\mu_{\alpha\dot\alpha}\partial_\mu - i(\Omega^r)_{ij}\overline{\theta}^j_{\dot\alpha}\frac{\partial}{\partial\zeta^r} \\ D_r &= \frac{\partial}{\partial\zeta^r}\,. \end{aligned} \tag{1.44}$$

Just as in the case of $\mathcal{N}$=1 superspace constraints defining superfields corresponding to irreducible representations of supersymmetry are implemented through the covariant derivatives (1.44). In general constrained superfields still depend on the coordinates ζ, so that the component supermultiplets contain an infinite number of fields; the only exception is the $\mathcal{N}$=2 *vector superfield* defined by the condition

$$D_r V(x,\theta,\overline{\theta},\zeta) = \frac{\partial}{\partial\zeta^r}V(x,\theta,\overline{\theta},\zeta) = 0\,. \tag{1.45}$$

As will be discussed in the next subsection the component fields associated to this superfield are those defining the $\mathcal{N}$=2 supersymmetric Yang–Mills model.

$\mathcal{N} > 1$ irreducible on-shell representations constructed in section 1.2 contain a finite number of states ($2^{2\mathcal{N}}$ in the massive case and $2^{\mathcal{N}}$ in the massless case). The reason for the appearance here of an infinite number of component fields is that closure of the supersymmetry algebra off-shell in the presence of central charges requires infinitely many auxiliary fields, the only exception being the $\mathcal{N}$=2 super Yang–Mills multiplet related to the superfield in equation (1.45).

The $\mathcal{N}$=2 supersymmetry algebra contains in principle two central charges, but one of them can always be eliminated through a chiral rotation, so that the only relevant anticommuting relation is

$$\{D^i_\alpha, D^j_\beta\} = 2\varepsilon_{\alpha\beta}\varepsilon^{ij}D_\zeta\,.$$

One can consider a superfield W satisfying

$$\overline{D}^i_{\dot\alpha} W = 0\,, \qquad\qquad D_\zeta W = 0\,. \tag{1.46}$$

A multiplet of fields can be obtained taking the $\theta = 0$ component of the fields

$$\begin{aligned} &W\,,\quad D^i_\alpha W\,,\quad D^{\alpha i} D^j_\alpha W\,,\quad D^i_\alpha D_{\beta i} W\,,\\ &D^i_\alpha D^j_\beta D^k_\gamma W\,,\quad D^i_\alpha D^j_\beta D^k_\gamma D^l_\delta W\,. \end{aligned} \tag{1.47}$$

Exploiting the conditions (1.46) in (1.47) does not produce an irreducible multiplet. A further reduction is necessary which is achieved by imposing the reality condition

$$D^{\alpha i} D^j_\alpha W = \overline{D}^i_{\dot\alpha} \overline{D}^{\dot\alpha j} \overline{W}\,.$$

This gives rise to a multiplet containing the fields

$$D\,,\quad \chi^i_\alpha\,,\quad C^{ij}\,,\quad F_{\mu\nu}\,, \tag{1.48}$$

where D and $C^{ij} = C^{ji}$ are real scalars, χ^i_α Weyl spinors and $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the field strength for a real vector field A_μ. These are the fields of the $\mathcal{N}$=2 super Yang–Mills model: they sum up a chiral and a vector $\mathcal{N}$=1 multiplet.

An example of a $\mathcal{N}$=2 multiplet possessing central charge arises if one considers a superfield satisfying

$$\begin{aligned} &D^i_\alpha \Phi_j = \frac{1}{2} \delta^i_j D^k_\alpha \Phi_k\\ &\overline{D}_{\dot\alpha i} \Phi_j + \overline{D}_{\dot\alpha j} \Phi_i = 0\,. \end{aligned} \tag{1.49}$$

Using these constraints a multiplet of fields can be obtained taking the $\theta = 0$ component of the fields

$$\Phi\,,\quad D^k_\alpha \Phi_k\,,\quad \overline{D}^k_{\dot\alpha} \Phi_k\,,\quad D_\zeta \Phi\,.$$

Some D-algebra manipulations allow to construct a supermultiplet containing two Weyl spinors and four complex scalars

$$\psi_\alpha\,,\quad \chi_\alpha\,,\quad \phi_i\,,\quad F_i\,. \tag{1.50}$$

This multiplet is known as *hypermultiplet* and can be viewed as the combination of two $\mathcal{N}$=1 chiral multiplets. It will be shown to describe $\mathcal{N}$=2 matter in supersymmetric gauge theories.

In general the constraints that need to be imposed on $\mathcal{N} > 1$ superfields cannot be explicitly solved for the whole superfield; in other words considering all the θ-components yields a set of non independent fields. The lack of an explicit solution of the constraints does not allow to exploit the potentiality of the superspace formalism for extended supersymmetry, so that it does not prove so powerful a tool as in the $\mathcal{N}$=1 case.

These problems were partially solved in [22] by the introduction of *harmonic superspace*. It is based on a parametrization of $\mathcal{N}$=2 superspace as a homogeneous space that leads to different bosonic coordinates associated to the central charges. Use of such coordinates allows to construct $\mathcal{N}$=2 hypermultiplets in terms of unconstrained superfields. A generalization of this approach leading to the so called *analytic superspace* has been proposed in order to give a manifestly supersymmetric description of $\mathcal{N}$=4 theories (see the review [21] for the details), however it only works on-shell.

In the following all the explicit calculations using superfield formalism will be carried out within the framework of $\mathcal{N}$=1 superspace.

1.5.2 $\mathcal{N} > 1$ supersymmetric models

In this section only examples of $\mathcal{N}$=2 theories will be discussed, the general properties of $\mathcal{N}$=4 models, that are the main subject of this work, will be studied in detail in next chapter.

$\mathcal{N}$=2 supersymmetric Yang–Mills model.

The superfield formulation of $\mathcal{N}$=2 pure supersymmetric Yang–Mills models is a straightforward generalization of the $\mathcal{N} = 1$ case because it is constructed in terms of a $\mathcal{N}$=2 superfield satisfying the constraint (1.45), which implies no dependence on the central charges. Since no central charges are present a full off-shell formulation is possible. Superspace in this case is parametrized by

$$z_m = (x_\mu, \theta^i_\alpha, \overline{\theta}_{\dot\alpha j}) = (x_\mu, \theta_\alpha, \overline{\theta}_{\dot\alpha}, \tilde\theta_\alpha, \overline{\tilde\theta}_{\dot\alpha})$$

The starting point is a chiral superfield Ψ defined by

$$\overline{D}_{\dot\alpha}\Psi = 0\,, \qquad \overline{\tilde D}_{\dot\alpha}\Psi = 0 \tag{1.51}$$

There is a natural choice of variables to describe Ψ, namely

$$\tilde y_\mu = x_\mu + i\theta\sigma_\mu\overline{\theta} + i\tilde\theta\sigma_\mu\overline{\tilde\theta}$$

Ψ can be expressed in terms of $\tilde{y}$ as

$$\Psi(\tilde{y},\theta) = \Phi(\tilde{y},\theta) + i\sqrt{2}\tilde{\theta}^{\alpha}W_{\alpha}(\tilde{y},\theta) + \tilde{\theta}\tilde{\theta}G(\tilde{y},\theta)\,,$$

where Φ and G are $\mathcal{N}$=1 chiral superfields and W_α is a chiral spinor superfield.

To construct a non Abelian gauge theory Ψ is taken to be a matrix

$$\Psi_{ij} = T^a_{ij}\Psi_a\,.$$

A $\mathcal{N}$=2 supersymmetric action can then be written as

$$S = \frac{1}{4\pi}\mathrm{Im}\left\{\tau\int d^4x\int d^2\theta d^2\tilde{\theta}\,\mathrm{tr}\left(\frac{1}{2}\Psi^2\right)\right\}\,, \tag{1.52}$$

where

$$\Psi^2\big|_{\theta^2\tilde{\theta}^2} = W^{\alpha}W_{\alpha}|_{\theta^2} + 2G\Phi|_{\theta^2}\ .$$

$\mathcal{N}$=2 super Yang–Mills is obtained by further constraining the chiral superfield G. This is achieved by requiring Ψ to satisfy

$$D^{\alpha i}D^j_{\alpha}\Psi = \overline{D}^i_{\dot{\alpha}}\overline{D}^{\dot{\alpha}j}\Psi^{\dagger}\,.$$

Solving the constraint provides for G the following expression

$$G = \int d^2\theta\,\Phi^{\dagger}(\tilde{y} - i\theta\sigma\overline{\theta},\overline{\theta})e^{V(\tilde{y}-i\theta\sigma\overline{\theta},\overline{\theta})}\,.$$

Substituting into the general form (1.52) gives the action of $\mathcal{N}$=2 super Yang–Mills theory. It can be viewed as a particular $\mathcal{N}$=1 supersymmetric gauge theory coupled to chiral matter, namely it corresponds to the case of one single flavor in the adjoint representation of the gauge group. Supersymmetry does not allow a superpotential for the model. Integration over the θ variables gives the component expansion. The superfield Ψ is a singlet under the SU(2) R-symmetry group. The resulting component multiplet obtained eliminating the auxiliary fields in the Wess–Zumino gauge contains a vector A_μ and a complex scalar ϕ that are SU(2) singlets and two Weyl spinors (ψ,λ) that transform in the **2** of SU(2). As already noticed, this field content is what would be obtained putting together a vector and a chiral $\mathcal{N}$=1 superfield both in the adjoint of the gauge group.

The most general $\mathcal{N}$=2 action that can be constructed in terms of the superfield Ψ of equation (1.51) is

$$S = \frac{1}{4\pi}\mathrm{Im}\left\{\int d^4x\int d^2\theta d^2\tilde{\theta}\,\mathcal{F}(\Psi)\right\}\,,$$

where $\mathcal{F}$ is a holomorphic function of Ψ called *prepotential*.

$\mathcal{N}$=2 matter.

The $\mathcal{N}$=2 super Yang–Mills model can be generalized introducing the coupling to matter described by hypermultiplets. Hypermultiplets are associated to superfields satisfying (1.49). A kinetic term for these superfields can be written in the form

$$S_K = \int d^4x\, D^{\alpha i} D^j_\alpha \left[\Phi^{i\,\dagger} D^{\beta k} D^k_\beta \Phi^j\right] .$$

Explicit dependence on the central charge does not allow a manifestly $\mathcal{N}$=2 formulation without resorting to harmonic superspace. Anyway an on-shell action can be constructed in terms of $\mathcal{N}=1$ superfields. Denoting the two chiral superfields in the hypermultiplet by Q and $\tilde{Q}$ the action in $\mathcal{N}$=1 language is given by

$$\begin{aligned} S = &\int d^4x \left\{ \int d^4\theta \, \left[Q^\dagger e^{2V} Q + \tilde{Q}^\dagger e^{2V} \tilde{Q} + \Phi^\dagger e^{2V} \Phi \right] + \right. \\ &\left. + \frac{1}{4\pi} \mathrm{Im} \left[\tau \int d^2\theta \, W^\alpha W_\alpha \right] + \left[\int d^2\theta \, \left(\sqrt{2} \tilde{Q} \Phi Q + m_i \tilde{Q}_i Q_i \right) + \mathrm{h.c.} \right] \right\} . \end{aligned} \tag{1.53}$$

The action (1.53) contains the ordinary kinetic terms, a mass term for chiral fields in the hypermultiplet and a superpotential that couples Q, $\tilde{Q}$ and Φ. Notice that no potential term coming from a self interaction of the fields Q and $\tilde{Q}$ is allowed by $\mathcal{N}$=2 supersymmetry. This follows from the observation that Q and $\tilde{Q}$ are assembled into a superfield Φ^i transforming in the **2** of SU(2) R-symmetry and there exists no trilinear invariant in SU(2).

1.6 Quantization of supersymmetric theories

The functional approach to the quantization of field theories, based on the construction of a generating functional for the time-ordered products of fields, can be applied with no further difficulty to supersymmetric theories in their component field formulation. However, although this formulation allows to extract all the peculiar properties of supersymmetric theories at the quantum level, it makes rather obscure how such properties are a direct consequence of supersymmetry. On the contrary a quantization procedure can be carried out directly in superspace in a step by step manifestly supersymmetric fashion. This approach, that allows to construct directly

Green functions for the superfields, has proved extremely powerful and it is perhaps the most useful application of superspace techniques.

Super Feynman rules for the computation of Green functions of the superfields were first proposed in [14] and then reformulated in a more powerful way in [23]. This latter formulation will be reviewed here. Once the Green functions for the superfields are known contact with the "physical" component field formulation is established by further integrating the fermionic coordinates $(\theta, \overline{\theta})$ to extract correlators for the various components.

The discussion will focus on the quantization of $\mathcal{N}$=1 superfields; it is also possible to generalize the method to $\mathcal{N}$-extended superspace, *e.g.* to harmonic superspace, but the $\mathcal{N}$=1 formulation appears to be the most suitable for application to explicit calculations, even in the extended case.

1.6.1 General formalism

In ordinary quantum field theories the path integral quantization is based on the construction of a generating functional for the Green functions. Denoting by ϕ_i the generic fields of the model and by $S[\phi]$ the classical action the generating functional is defined as

$$Z[J] = N \int [\mathcal{D}\phi]\, e^{iS[\phi]+i\int dx \phi_i(x) J_i(x)}\,,$$

where N is a normalization such that $Z[0] = 1$ and $J_i(x)$ are external sources associated to the fields $\phi_i(x)$. Green functions are obtained taking functional derivatives with respect to the sources J_i.

This approach has a natural extension that leads to a generating functional for the Green functions of the superfields in a supersymmetric theory.

The general form of the action of a supersymmetric theory involves, in $\mathcal{N}$=1 language, chiral and vector superfields and can be written

$$S = \int d^4x d^4\theta\, K(\Phi, \Phi^\dagger, V) + \left[\int d^4x d^2\theta\, \mathrm{W}[\Phi] + \mathrm{h.c.}\right]\,. \tag{1.54}$$

To construct a generating functional classical sources for the different superfields must be introduced, that are in turn superfields. Since functional derivatives are to be taken with respect to the classical sources, such sources must be arbitrary. A problem arises with chiral superfields as sources coupled to them must themselves

be chiral. This difficulty can be solved by using a suitable definition of functional derivatives with respect to chiral superfields.

For a generic superfield $F(x,\theta,\overline{\theta})$ functional derivatives are defined according to the rule

$$\frac{\delta F(x',\theta',\overline{\theta}')}{\delta F(x,\theta,\overline{\theta})} = \delta_4(x-x')\delta_4(\theta-\theta')\,,$$

where the δ function of fermionic variables is to be interpreted as

$$\delta_4(\theta-\theta') = \delta_2(\theta-\theta')\delta_2(\overline{\theta}-\overline{\theta}') = (\theta-\theta')^2(\overline{\theta}-\overline{\theta}')^2\,,$$

so that

$$\int d^4\theta\,\delta_4(\theta-\theta')F(x,\theta,\overline{\theta}) = F(x,\theta',\overline{\theta}')$$

and

$$\frac{\delta}{\delta F(x,\theta,\overline{\theta})}\int d^4x'd^4\theta'\,F(x',\theta',\overline{\theta}')G(x',\theta',\overline{\theta}') = G(x,\theta,\overline{\theta})\,. \tag{1.55}$$

For chiral superfields Φ one defines in turn

$$\frac{\delta\Phi(x',\theta',\overline{\theta}')}{\delta\Phi(x,\theta,\overline{\theta})} = -\frac{1}{4}\overline{D}^2\delta_4(x-x')\delta_4(\theta-\theta')\,.$$

In this way the same result as in equation (1.55) is reproduced

$$\begin{aligned}
&\frac{\delta}{\delta\Phi(x,\theta,\overline{\theta})}\int d^4x'd^2\theta'\,\Phi(x',\theta',\overline{\theta}')\tilde{\Phi}(x',\theta',\overline{\theta}') = \\
&= \int d^4x'd^2\theta'\left(-\frac{1}{4}\overline{D}^2\right)\delta_4(x-x')\delta_4(\theta-\theta')\tilde{\Phi}(x',\theta',\overline{\theta}') = \\
&= \int d^4x'd^4\theta'\delta_4(x-x')\delta_4(\theta-\theta')\tilde{\Phi}(x',\theta',\overline{\theta}') = \tilde{\Phi}(x,\theta,\overline{\theta})\,,
\end{aligned}$$

where in the last line it was used the property that $d^2\overline{\theta}$ and $-\frac{1}{4}\overline{D}^2$ are equivalent under space-time integration, as can be proved by direct calculation by an integration by parts.

Given these definitions of functional derivatives the generating functional can be written as

$$Z[J,J^\dagger,J_V] = N\int[\mathcal{D}\Phi\mathcal{D}\Phi^\dagger\mathcal{D}V]\,e^{iS[\Phi,\Phi^\dagger,V]+i(\Phi,J)+i(\Phi^\dagger,J^\dagger)+i(V,J_V)}\,, \tag{1.56}$$

where S is the action (1.54), J ($J^\dagger$) is a chiral (antichiral) source, J_V is real and

$$\begin{aligned}(\Phi, J) &= \int d^4x d^2\theta\, \Phi(x,\theta,\overline{\theta}) J(x,\theta,\overline{\theta})\,,\\ (\Phi^\dagger, J^\dagger) &= \int d^4x d^2\overline{\theta}\, \Phi^\dagger(x,\theta,\overline{\theta}) J^\dagger(x,\theta,\overline{\theta})\,,\\ (V, J_V) &= \int d^4x d^4\theta\, V(x,\theta,\overline{\theta}) J_V(x,\theta,\overline{\theta})\,.\end{aligned}$$

Green functions can then be calculated by the formula

$$\begin{aligned}&\mathcal{G}_n(z_1,\dots,z_i,z_{i+1},\dots,z_j,z_{j+1},\dots,z_n) =\\ &= \langle 0|T\{\Phi(z_1)\dots\Phi(z_i)\Phi^\dagger(z_{i+1})\dots\Phi^\dagger(z_j)V(z_{j+1})\dots V(z_n)\}|0\rangle = \\ &= (-i)^n \frac{\delta^n Z[J,J^\dagger,J_V]}{\delta J(z_1)\dots\delta J(z_i)\delta J^\dagger(z_{i+1})\dots\delta J^\dagger(z_j)\delta J_V(z_{j+1})\dots\delta J_V(z_n)}\bigg|_{J=J^\dagger=J_V=0}\,.\end{aligned} \tag{1.57}$$

A generating functional for the connected Green functions can be obtained by taking the logarithm of Z

$$W[J,J^\dagger,J_V] = (-i)\,\log\left(Z[J,J^\dagger,J_V]\right)\,.$$

The quantum effective action Γ that acts as a generating functional for the one particle irreducible Green functions is obtained through the Legendre transform. Defining the "classical" fields

$$\tilde{\Phi} = \frac{\delta W}{\delta J}\,, \qquad \tilde{\Phi}^\dagger = \frac{\delta W}{\delta J^\dagger}\,, \qquad \tilde{V} = \frac{\delta W}{\delta J_V}\,, \tag{1.58}$$

Γ is given by

$$\Gamma[\tilde{\Phi},\tilde{\Phi}^\dagger,\tilde{V}] = \left(W[J,J^\dagger,J_V] - \left[(J,\Phi)+(J^\dagger,\Phi^\dagger)+(J_V,V)\right]\right)\Big|_{(\tilde{\Phi},\tilde{\Phi}^\dagger,\tilde{J}_V)}\,,$$

where on the right hand side $J = J[\tilde{\Phi},\tilde{\Phi}^\dagger,\tilde{V}]$, $J^\dagger = J^\dagger[\tilde{\Phi},\tilde{\Phi}^\dagger,\tilde{V}]$ and $J_V = J_V[\tilde{\Phi},\tilde{\Phi}^\dagger,\tilde{V}]$ come from the inverting (1.58).

The generating functional Z can be expressed in terms of the corresponding functional in the free theory, Z_0. The general action (1.54) can be divided up into a free part and an interacting part

$$S[\Phi,\Phi^\dagger,V] = S_0[\Phi,\Phi^\dagger,V] + S_{\rm int}[\Phi,\Phi^\dagger,V]\,,$$

where S_0 is quadratic in the fields and contains the kinetic and mass terms. Then one can write

$$Z[J,J^\dagger,J_V] = e^{iS_{\rm int}\left[\frac{\delta}{\delta J},\frac{\delta}{\delta J^\dagger},\frac{\delta}{\delta J_V}\right]} Z_0[J,J^\dagger,J_V]\,, \tag{1.59}$$

where

$$Z_0[J, J^\dagger J_V] = N \int [\mathcal{D}\Phi \mathcal{D}\Phi^\dagger \mathcal{D}V]\, e^{iS_0[\Phi,\Phi^\dagger,V]+i(\Phi,J)+i(\Phi^\dagger,J^\dagger)+i(V,J_V)}$$

and $e^{iS_{\rm int}[\frac{\delta}{\delta J},\frac{\delta}{\delta J^\dagger},\frac{\delta}{\delta J_V}]}$ is to be understood as a power expansion. Since S_0 is quadratic Z_0 is a Gaussian integral that can be explicitly computed. Writing

$$\begin{aligned} Z_0[J, J^\dagger, J_V] = \exp\Bigg\{ & -\frac{i}{2}\int d^4x d^4\theta\, (\Phi, \Phi^\dagger)\mathcal{M}_\Phi \begin{pmatrix} \Phi \\ \Phi^\dagger \end{pmatrix} + \\ & + \frac{i}{2}\int d^4x d^4\theta\, V\mathcal{M}_V V + i(\Phi, J) + i(\Phi^\dagger, J^\dagger) + i(V, J_V)\Bigg\} \end{aligned} \tag{1.60}$$

one gets

$$\begin{aligned} Z_0[J, J^\dagger, J_V] &= \exp\left\{ \frac{i}{2}\int d^8z\, d^8z'\, (J(z), J^\dagger(z))\Delta_\Phi(z,z') \begin{pmatrix} J(z') \\ J^\dagger(z') \end{pmatrix} + \right. \\ &+ \left. \frac{i}{2}\int d^8z\, d^8z'\, J_V(z)\Delta_V(z,z')J_V(z') \right\}, \end{aligned} \tag{1.61}$$

where superfield propagators, Δ_Φ and Δ_V, have been introduced.

1.6.2 Superfield propagators

To compute the propagators the free action must be put in a canonical quadratic form. Since the computation leads to different kinds of difficulties for chiral and vector fields the two cases will be considered separately.

Chiral superfield propagator.

The general form of the free action is

$$\begin{aligned} S_0 &= \int d^4x \left\{ \int d^4\theta\, \Phi_i^\dagger \Phi_i + \left[\int d^2\theta\, \frac{1}{2} m_{ij}\Phi_i\Phi_j + \int d^2\overline{\theta}\, \frac{1}{2} m_{ij}\Phi_i^\dagger\Phi_j^\dagger \right] \right\} = \\ &= \int d^4x d^4\theta \left[\Phi_i^\dagger\Phi_i - \frac{1}{8} m_{ij} \left(\Phi_i \frac{DD}{\Box}\Phi_j + \Phi_i^\dagger \frac{\overline{DD}}{\Box}\Phi_j^\dagger \right) \right], \end{aligned} \tag{1.62}$$

where the last line follows from the equivalence $d^2\theta \leftrightarrow -\frac{1}{4}D^2$, $d^2\overline{\theta} \leftrightarrow -\frac{1}{4}\overline{D}^2$, valid under a space-time integration, and the action of the projection operators

$$\begin{aligned} P_1 &= \frac{1}{16}\frac{D^2\overline{D}^2}{\Box}, \qquad & P_1\Phi^\dagger = \Phi^\dagger, \qquad & P_1\Phi = 0 \\ P_2 &= \frac{1}{16}\frac{\overline{D}^2 D^2}{\Box}, \qquad & P_2\Phi = \Phi, \qquad & P_2\Phi^\dagger = 0. \end{aligned} \tag{1.63}$$

Hence the action can be rewritten as

$$S_0 = \frac{1}{2}\int d^4x d^4\theta \, \left(\Phi_i, \Phi_i^\dagger\right) \mathcal{M}_{ij} \left(\begin{array}{c} \Phi_j \\ \Phi_j^\dagger \end{array} \right) , \tag{1.64}$$

where

$$\mathcal{M}_{ij} = \left(\begin{array}{cc} -\frac{1}{4}\frac{m_{ij}}{\Box} DD & \delta_{ij} \\ \delta_{ij} & -\frac{1}{4}\frac{m_{ij}}{\Box}\overline{DD} \end{array} \right) .$$

Considering for simplicity the case of one single flavor and taking into account the source terms, the exponent of the Gaussian integral that yields Z_0 turns out to be

$$i\int d^4x d^4\theta \, \left[\frac{1}{2}\left(\Phi, \Phi^\dagger\right)\mathcal{M}\left(\begin{array}{c} \Phi \\ \Phi^\dagger \end{array} \right) + \left(\Phi, \Phi^\dagger\right)\left(\begin{array}{cc} -\frac{1}{4}\frac{D^2}{\Box} & 0 \\ 0 & -\frac{1}{4}\frac{\overline{D}^2}{\Box} \end{array} \right)\left(\begin{array}{c} J \\ J^\dagger \end{array} \right)\right] .$$

A simple but lengthy calculation using D-algebra and properties of the projection operators allows to write Z_0 in the form of equation (1.61), where the chiral superfield propagator, known as *Grisaru–Roček–Siegel propagator* [23], reads

$$\Delta_\Phi(x,\theta,\overline{\theta};x',\theta',\overline{\theta}') \;=\; \frac{1}{\Box - m^2}\left(\begin{array}{cc} \frac{m}{4}\frac{D^2}{\Box} & 1 \\ 1 & \frac{m}{4}\frac{\overline{D}^2}{\Box} \end{array} \right)\delta_8(z-z') \tag{1.65}$$

and $\delta_8(z-z') = \delta_4(x-x')\delta_4(\theta-\theta')$.

Vector superfield propagator.

The free field action for a vector superfield is

$$S_0 = \int d^4x \left\{ \int d^2\theta \, \frac{1}{4} W^\alpha W_\alpha + \int d^2\overline{\theta} \, \frac{1}{4}\overline{W}_{\dot\alpha}\overline{W}^{\dot\alpha} + \int d^4\theta \, m^2 V^2 \right\} . \tag{1.66}$$

No massive vector field will enter the models considered in the following so m will be put to zero from the beginning.

The kinetic operator in (1.66) has zero-modes so that just like in ordinary gauge theories a gauge fixing procedure *à la* Faddeev–Popov is necessary. This requires the introduction of a gauge fixing term in (1.66) and a further integration over the Faddeev–Popov ghost fields will appear. The gauge-fixed action thus becomes

$$\begin{aligned} & S = S_0 + S_{\text{GF}} + S_{\text{FP}} = \\ & = \int d^4x d^4\theta \, \left[\frac{1}{4} W^\alpha W_\alpha \delta_2(\overline{\theta}) + \frac{1}{4}\overline{W}_{\dot\alpha}\overline{W}^{\dot\alpha}\delta_2(\theta)\right] + \\ & + \int d^4x d^4\theta \, \left[-\frac{\xi}{8}(\overline{D}^2 V)(D^2 V)\right] + \\ & + i\int d^4x d^4\theta \, (C' + \overline{C}')\, \mathcal{L}_{\frac{V}{2}} \left[(C+\overline{C}) + \coth \mathcal{L}_{\frac{V}{2}}(C-\overline{C})\right] , \end{aligned} \tag{1.67}$$

where C, C' are chiral superfields for the ghosts and no source term has been introduced for such fields. The gauge fixing term here introduced corresponds to a class of gauges that for the component field A_μ interpolate between the Lorentz gauge ($\xi = \infty$) and the Fermi–Feynman gauge ($\xi = 1$).

The free propagators for the ghosts are exactly the same as for ordinary chiral superfields. Furthermore there is a non polynomial interaction with the vector superfield coming from the expansion of $\coth(\mathcal{L}_{V/2})$.

The vector superfield propagator is calculated from the quadratic part of the action

$$\begin{aligned} S &= S_0 + S_{\rm GF} = \\ &= \int d^4x d^4\theta \, \{V \left[-\Box P_T - \xi(P_1 + P_2)\Box\right] V\} = \\ &= \int d^4x d^4\theta \, V \mathcal{M}_V V \,. \end{aligned} \tag{1.68}$$

Introducing the source term for V the generating functional becomes

$$Z[J_V] = \int [\mathcal{D}V] \, e^{i \int V \mathcal{M}_V V + i(V, J_V)} \,.$$

The Gaussian integral can now be performed yielding a result of the form (1.61), where the vector superfield propagator is

$$\begin{aligned} \Delta_V(x, \theta, \overline{\theta}; x', \theta', \overline{\theta}') &= \left[-\frac{1}{\Box} P_T - \frac{\alpha}{\Box}(P_1 + P_2)\right] \delta_8(z - z') = \\ &= -\frac{1}{\Box} \left[1 + (\alpha - 1)(P_1 + P_2)\right] \delta_8(z - z') \,, \end{aligned} \tag{1.69}$$

with $\alpha = \frac{1}{\xi}$.

1.6.3 Feynman rules

Once the free field propagators are known the perturbative expansion for the Green functions of the superfields can be derived from the generating functional in the form (1.59) that yields

$$\begin{aligned} &\mathcal{G}_n(z_1, \ldots, z_i, z_{i+1}, \ldots, z_j, z_{j+1}, \ldots, z_n) = \\ &= \langle 0 | T\{\Phi(z_1) \ldots \Phi(z_i) \Phi^\dagger(z_{i+1}) \ldots \Phi^\dagger(z_j) V(z_{j+1}) \ldots V(z_n)\} | 0 \rangle = \\ &= \frac{\delta}{\delta J(z_1)} \cdots \frac{\delta}{\delta J(z_i)} \frac{\delta}{\delta J^\dagger(z_{i+1})} \cdots \frac{\delta}{\delta J^\dagger(z_j)} \frac{\delta}{\delta J_V(z_{j+1})} \cdots \frac{\delta}{\delta J_V(z_n)} \cdot \\ &\cdot \sum_{k=0}^{\infty} \frac{(-i)^k}{k!} \left(S_{\rm int} \left[\frac{\delta}{\delta J}, \frac{\delta}{\delta J^\dagger}, \frac{\delta}{\delta J_V} \right] \right)^k Z_0[J, J^\dagger, J_V] \Big|_{J = J^\dagger = J_V = 0} \,. \end{aligned} \tag{1.70}$$

A super Feynman graph technique can be developed which is a straightforward generalization of what is done in ordinary field theories. Super Feynman rules for the evaluation of one particle irreducible Green functions can be derived given the above generating functional. The rules can be summarized as follows.

- To each external line is associated the corresponding superfield.
- At each vertex there is an integration over the whole superspace $d^4xd^4\theta$; factors, as for example powers of the coupling constant, to be associated to the vertices, are read directly from the action.
- A factor of $-\frac{1}{4}\overline{D}^2$ or $-\frac{1}{4}D^2$ acting on the internal chiral or antichiral lines respectively must be included at each vertex. Such factors come from the definition of the functional derivatives. At purely chiral (antichiral) vertices one factor of $-\frac{1}{4}\overline{D}^2$ $(-\frac{1}{4}D^2)$ must be suppressed, because it is used to reconstruct the measure over the whole superspace.
- For internal lines the propagators previously derived are used.
- The specific form of the interaction at each vertex as well as the combinatoric analysis are worked out exactly like in ordinary perturbation theory.
- The expressions for the one particle irreducible Green functions in momentum space are obtained by Fourier transforming in the x variables, but not in the fermionic θ coordinates.

1.7 General properties of supersymmetric theories

Supersymmetry has far reaching consequences in the quantum theory both at the perturbative and at the non-perturbative level. Some general results will be briefly recorded in this section. More can be found in the books and reviews quoted in the references.

Some dramatic effects on the divergences encountered in perturbation theory can be easily derived from the Feynman rules. Explicit calculation shows that in Feynman diagrams four covariant derivatives are associated at each vertex, with the exception of vertices involving external (anti) chiral lines that have two derivatives less. Moreover $\Phi\Phi$ $(\Phi^\dagger\Phi^\dagger)$ propagators with momentum p have an additional factor of $\frac{D^2}{p^2}$ $(\frac{\overline{D}^2}{p^2})$.

Simple dimensional analysis allows to compute the superficial degree of divergence d_s of a diagram which is

$$d_s = 4L - 2P + 2V - C - E - 2L\,,$$

where L is the number of loops, P the number of internal propagators, V the number of vertices, C the number of $\Phi\Phi$ or $\Phi^\dagger\Phi^\dagger$ propagators and E the number of external chiral or antichiral lines. Use of the topological constraint $L - P + V = 1$ yields [24]

$$d_s = 2 - C - E\,.$$

For graphs with only external V lines gauge invariance requires the presence of four D or $\overline{D}$ factors on the external lines so that the result is

$$d_s = -C$$

For diagrams with only lines of a given chirality there are further factors of D^2 or $\overline{D}^2$ associated to the external lines so that one obtains

$$d_s = 1 - C - n\,,$$

where n is the number of external chiral or antichiral lines.

This calculation allows to determine all the potentially divergent diagrams. The only possible divergences are logarithmic and can arise in diagrams with only external V lines with $C = 0$ (*e.g.* corrections to the vector superfield propagator) and in $\Phi^\dagger\Phi$ propagators with $C = 0$. This shows that in supersymmetric models only wave function renormalizations may be required; there is no direct mass or coupling constant renormalization, *i.e.* with $\mathrm{W} = m\phi^2$ one has $Z_\mathrm{W} = 1$, but $Z_\phi \neq 1$ implies $Z_m = Z_\phi^{-1/2} \neq 1$.

This and other results can be obtained from the general non renormalization theorem for $\mathcal{N}$=1 theories [25]: *Any perturbative quantum contribution to the effective action* Γ *can be expressed as an integral over the whole superspace and is local in the* θ *variables*, namely

$$\begin{aligned}\Gamma \;&=\; \sum_n \int d^4x_1 \ldots d^4x_n d^4\theta\, G(x_1, \ldots, x_n)\cdot \\ &\qquad \cdot f_n[\Phi, \Phi^\dagger, V, D_\alpha\Phi, \overline{D}_{\dot\alpha}\Phi^\dagger, D_\alpha V, \overline{D}_{\dot\alpha}V, \ldots]\,. \end{aligned} \tag{1.71}$$

The theorem can be easily proved using Feynman rules and successively integrating by parts the θ variables exploiting the properties of Grassmannian integration. Explicit

calculations in the $\mathcal{N}$=4 supersymmetric Yang–Mills theory showing the mechanism leading to this result will be presented in chapter 3, so the proof of the above statement will not be discussed here.

Notice that the non renormalization theorem does not forbid in principle quantum corrections to the superpotential. It is easy to see that a contribution of the form of (1.71) can produce a correction to the superpotential [26, 27]. As an example consider a term of the form

$$\int d^4x d^2\theta d^2\overline{\theta} \frac{D^2}{16\Box} \Phi^n = \int d^4x d^2\theta \frac{\overline{D}^2 D^2}{16\Box} \Phi^n = \int d^4x d^2\theta \Phi^n \,,$$

where Φ is chiral. The right hand side, which follows since $P_1 = \frac{\overline{D}^2 D^2}{16\Box}$ is the projector on chiral superfields, has the form of a correction to the superpotential. This example shows that in principle an arbitrary correction to the superpotential can be generated perturbatively. This effect is related to an infrared singular behaviour that can be present only in theories containing massless fields. The example shows that the correction, which is non-local when written as an integral over the whole superspace, can be rewritten as a local term integrated over a subspace. The same construction could not be repeated if the theory does not contain massless particles, since in that case one would obtain a factor of $\frac{1}{\Box+m^2}$, which cannot be re-expressed as a subintegral. In [27] it was shown that such a correction as the one considered above actually occurs at the two-loop level in the Wess–Zumino model. In a $\mathcal{N}$=1 super Yang–Mills theory coupled to the Wess–Zumino model the correction appears at one loop [28]. An alternative approach to the study of this kind of effects has been given in [29, 30].

Other non renormalization theorems hold for theories of extended supersymmetry. In particular considerations based on the so called *anomalies argument* allow to prove the finiteness of $\mathcal{N}$=2 theories beyond one loop. More precisely it can be proved that the only perturbative contributions to the β function of extended supersymmetric theories can come from one loop diagrams. This result allows to single out a class of finite supersymmetric models and among them the $\mathcal{N}$=4 super Yang–Mills theory that will be studied in detail in the following chapters.

Supersymmetry puts severe restrictions on the non-perturbative properties as well. A review of the general results on the non-perturbative dynamics of supersymmetric theories is provided by [31, 32]. Many astonishing results in this context can be traced back to the properties of *holomorphy* of supersymmetric theories. In supersymmetric models for example the superpotential is a holomorphic function of the chiral super-

fields. Analogously, since the parameters of the theories (such as masses and coupling constants) can be viewed as expectation values of auxiliary fields, various quantities depend analytically on such parameters. Using holomorphy properties many exact result have been derived. In [33] an expression for the β function of supersymmetric gauge theories was proposed that is exact at the perturbative as well as at the non-perturbative level for a class of models. Also on holomorphy relies the construction of the exact solution for the low energy dynamics of a class of $\mathcal{N}$=2 theories proposed by Seiberg and Witten [34].

Chapter 2

$\mathcal{N} = 4$ supersymmetric Yang–Mills theory

Four dimensional $\mathcal{N}$=4 supersymmetric Yang–Mills theory is a very special quantum field theory. It possesses several peculiar properties that will be reviewed in this chapter. The general properties of the model are discussed here, original results in the study of $\mathcal{N}$=4 Yang–Mills will be presented in the following chapters.

The action of the theory was given for the first time in [35, 36] within the framework of string theory toroidal compactifications. The theory has the maximal amount of supersymmetry for a rigid supersymmetric theory in four dimensions, namely sixteen real supercharges; a larger number of supercharges would require fields of spin larger than one and therefore the inclusion of gravity.

Historically the interest in this model was raised by its property of finiteness; the β function of Gell-Mann and Low has been proved to be vanishing in perturbation theory and the same is supposed to be true at the non-perturbative level. As a consequence the superconformal invariance displayed by the classical theory is believed to be preserved after quantization.

Moreover $\mathcal{N}$=4 super Yang–Mills theory possesses exact electric-magnetic duality. It contains beyond the elementary fields an infinite set of dyonic states which are believed to realize exactly the duality symmetry proposed by Montonen and Olive in [37].

More recently there has been a renewal of interest in $\mathcal{N}$=4 Yang–Mills theory

following the proposal made by Maldacena in [38] of a new duality relating type IIB supergravity in $d+1$ dimensional anti-de Sitter space and d dimensional (super)conformal theories. This subject will be discussed in detail in chapter 5.

The present chapter is organized as follows. In section 2.1 the various formulations of the model, which will be employed at different stages in the following, are reviewed. Section 2.2 presents the classical symmetries and conserved currents of the theory. The properties of finiteness are discussed in section 2.3. The final section is then devoted to electric-magnetic duality in $\mathcal{N}$=4 supersymmetric Yang–Mills theory. The general aspects of electric-magnetic duality are reported separately in appendix B.

2.1 Diverse formulations of $\mathcal{N}$=4 supersymmetric Yang–Mills theory

$\mathcal{N}$=4 supersymmetric Yang–Mills theory in four dimensions was obtained for the first time in [35, 36] by applying the method of dimensional reduction to $\mathcal{N}$=1 super Yang–Mills in ten dimensions. The latter is the low energy effective theory coming from type I superstring theory and describes a $\mathcal{N}=1$ vector multiplet in ten dimensions consisting of one real vector and one Majorana–Weyl spinor. The action is [1]

$$S = \int d^{10}x \,\mathrm{tr}\left\{-\frac{1}{4}F_{\Gamma\Lambda}F^{\Gamma\Lambda} + \frac{i}{2}\overline{\lambda}\Gamma^{\Lambda}D_{\Lambda}\lambda\right\}, \tag{2.1}$$

where $F_{\Gamma\Lambda} = \partial_{\Gamma}A_{\Lambda} - \partial_{\Lambda}A_{\Gamma} + ig[A_{\Gamma}, A_{\Lambda}]$ and λ satisfies the Majorana and Weyl conditions

$$\lambda = C_{(10)}\overline{\lambda}^{T}, \qquad \lambda = \pm\Gamma_5\lambda,$$

where $C_{(10)}$ is the charge conjugation operator. The action (2.1) is invariant under the supersymmetry transformations

$$\begin{aligned} \delta_{\eta}A_{\Lambda} &= i\overline{\eta}\Gamma_{\Lambda}\lambda \\ \delta_{\eta}\lambda &= \Sigma_{\Gamma\Lambda}F^{\Gamma\Lambda}\eta\,. \end{aligned} \tag{2.2}$$

In the previous equations capital Greek letters Γ and Σ denote ten dimensional matrices.

Dimensional reduction *á la* Kaluza–Klein on a six dimensional torus T^6 leads to a multiplet of fields in four dimensions possessing an additional SU(4)$\sim$SO(6) global

[1] Capital Greek indices refer to ten dimensional space-time.

symmetry, which is a direct consequence of the ten dimensional Lorentz invariance. More precisely it is a consequence of the fact that the torus T^6 has a trivial holonomy group. In general in the dimensional reduction on a Riemannian manifold of dimension k the spin connection is a SO(k) gauge field and consequently the spinors transform, upon parallel transport around a closed contractible curve, under a subgroup of SO(k), which is the holonomy group. In the case at hand of T^6 every spinor is covariantly constant, so that the whole SO(6)∼SU(4) becomes a global symmetry of the model. From the point of view of the $\mathcal{N}$=4 theory this SU(4) global symmetry is identified with the R-symmetry group of the $\mathcal{N}$=4 supersymmetry algebra, as will be discussed in the next section.

In the bosonic sector the compactification gives rise to a four dimensional real vector and to six real scalars from the internal components of the ten dimensional vector. More precisely one defines

$$\begin{aligned} A^a_\mu &= A^a_\mu & \mu &= 0,1,2,3 \\ \varphi^a_{B4} &= \tfrac{1}{\sqrt{2}}(A^a_{B+3} + iA^a_{B+6}) & B &= 1,2,3 \\ \varphi^{a\,AB} &= \tfrac{1}{2}\varepsilon^{ABC4}\varphi^a_{C4} = \left(\varphi^{a\,AB}\right)^* & A,B,C &= 1,2,3\,. \end{aligned} \tag{2.3}$$

In the fermionic sector a suitable choice of the ten-dimensional Γ matrices allows to write, in Majorana notation, the 32-component Majorana–Weyl spinor λ as

$$\lambda = \begin{pmatrix} L\chi^1 \\ \vdots \\ L\chi^4 \\ R\widetilde{\chi}_1 \\ \vdots \\ R\widetilde{\chi}_4 \end{pmatrix}, \qquad \tilde{\chi}_A = C(\overline{\chi}^A)^T\,,$$

where L and R denote the left and right chirality projection operators and C is the four-dimensional charge conjugation operator related to the ten-dimensional one by $C_{(10)} = C \otimes \begin{pmatrix} 0 & \mathbb{1}_4 \\ \mathbb{1}_4 & 0 \end{pmatrix}$. In four dimensions one obtains four Majorana (or equivalently Weyl) spinors

$$\lambda^A = \begin{pmatrix} L\chi^A \\ \\ R\widetilde{\chi}_A \end{pmatrix} \qquad A = 1,2,3,4\,. \tag{2.4}$$

At the end of the day the result of the compactification is the $\mathcal{N} = 4$ multiplet which contains six real scalars $\varphi^{a\,AB}$ satisfying (2.3), one real vector A^a_μ and four

Weyl spinors λ_α^{aA}, all in the adjoint representation of the gauge group. The construction here described from the ten dimensional theory allows to derive immediately the transformation of the fields under the SU(4) global symmetry. Because of the constraint (2.3) the scalars $\varphi^{a\,AB}$ are in the second rank complex self dual **6** of SU(4); the spinors λ^{aA} are in the **4** while the $\overline{\lambda}^a_A$ transform in the $\overline{\mathbf{4}}$; the gauge field A^a_μ is a singlet. In the following a different notation for the scalar fields will also be used, namely one can define

$$\varphi^i = \frac{1}{2}\overline{t}^i{}_{AB}\varphi^{AB}\,, \qquad\qquad i = 1, 2, \ldots, 6\,,$$

where $(t_i)^{AB}$ are Clebsch–Gordan coefficients that couple two **4**'s to a **6** (these are six-dimensional generalizations of the four-dimensional $\sigma^\mu{}_{\alpha\dot\alpha}$ matrices).

In Weyl notation the action of the four dimensional $\mathcal{N}$=4 theory turns out to be

$$\begin{aligned} S = \int d^4x\, \mathrm{tr}\Big\{ &(D_\mu\varphi^{AB})(D^\mu\overline{\varphi}_{AB}) - \frac{1}{2}i(\lambda^{\alpha A}\overrightarrow{\not{D}}_{\alpha\dot\alpha}\overline{\lambda}^{\dot\alpha}{}_A) - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \\ &-g\lambda^{\alpha A}[\lambda_\alpha{}^B, \overline{\varphi}_{AB}] - g\overline{\lambda}_{\dot\alpha A}[\overline{\lambda}^{\dot\alpha}{}_B, \varphi^{AB}] + 2g^2[\varphi^{AB}, \varphi^{CD}][\overline{\varphi}_{AB}, \overline{\varphi}_{CD}]\Big\}\,, \end{aligned} \tag{2.5}$$

where tr denotes a trace over the colour indices and $D^{ab}_\mu = \delta^{ab}\partial_\mu + igf^{abc}A_{\mu c}$ is the covariant derivative in the adjoint representation.

The action (2.5) is invariant under the supersymmetry transformations

$$\begin{aligned} \delta\varphi^{AB} &= \frac{1}{2}(\lambda^{\alpha A}\eta_\alpha{}^B - \lambda^{\alpha B}\eta_\alpha{}^A) + \frac{1}{2}\varepsilon^{ABCD}\overline{\eta}_{\dot\alpha C}\overline{\lambda}^{\dot\alpha}_D \\ \delta\lambda_\alpha{}^A &= -\frac{1}{2}F^-_{\mu\nu}\sigma^{\mu\nu}{}_\alpha{}^\beta\eta_\beta{}^A + 4i(\not{D}_{\alpha\dot\alpha}\varphi^{AB})\overline{\eta}^{\dot\alpha}_B - 8g[\overline{\varphi}_{BC}, \varphi^{CA}]\eta_\alpha{}^B \\ \delta A^\mu &= -i\lambda^{\alpha A}\sigma^\mu{}_{\alpha\dot\alpha}\overline{\eta}^{\dot\alpha}{}_A - i\eta^{\alpha A}\sigma^\mu{}_{\alpha\dot\alpha}\overline{\lambda}^{\dot\alpha}{}_A\,, \end{aligned} \tag{2.6}$$

which can be easily obtained from the ten dimensional transformations (2.2) using (2.3) and (2.4).

A manifestly off-shell formulation of $\mathcal{N}$=4 supersymmetric Yang–Mills theory would require the introduction of an infinite set of auxiliary fields as a consequence of the presence of bosonic coordinates associated to the twelve central charges in the $\mathcal{N}$=4 superspace, as noticed in [39, 40]. A superfield formulation can be constructed using analytic superspace, however such an approach is rather involved and cannot be applied to explicit calculations because it is intrinsically on-shell. An on-shell superfield description can be given in terms of a superfield $W^{AB} = W^{AB}(x, \theta^A_\alpha, \overline{\theta}_{\dot\alpha A})$ satisfying [41] the reality condition

$$\overline{W}_{AB} = \frac{1}{2}\varepsilon_{ABCD}W^{CD}\,, \tag{2.7}$$

together with the constraint,

$$\mathcal{D}_\alpha^A W^{BC} = \mathcal{D}_\alpha^{[A} W^{BC]}\,, \tag{2.8}$$

where $\mathcal{D}_\alpha$ is the super-covariant derivative. The fields entering the action (2.5) are combined into the lowest components of W^{AB}.

In the next chapter perturbative calculations in $\mathcal{N}$=4 Yang–Mills will be carried out using a formulation of the model in terms of $\mathcal{N}=1$ superfields. The field content of the theory can be obtained by taking one $\mathcal{N}$=1 vector superfield V and three $\mathcal{N}$=1 chiral superfields Φ^I all in the adjoint representation of the gauge group. In this approach the six real scalars are combined into three complex fields that are the scalar components of the chiral superfields Φ^I, three of the Weyl fermions are the spinors of Φ^I and the fourth fermion is the gaugino that together with the real vector constitute the vector superfield V. As a result only a SU(3)×U(1) subgroup of the original SU(4) symmetry is manifest in this formulation. More precisely the representations of SU(4) decompose according to $\mathbf{6} \to \mathbf{3}+\overline{\mathbf{3}}$, $\mathbf{4} \to \mathbf{3}+\mathbf{1}$, so that the chiral superfields Φ^I transform in the $\mathbf{3}$ of SU(3), the antichiral $\Phi_I^\dagger$ in the $\overline{\mathbf{3}}$ and the vector V is a singlet under SU(3).

The action in the $\mathcal{N}$=1 superfield formulation reads

$$\begin{aligned} S &= \frac{1}{d_{\mathbf{r}}}\mathrm{tr}\left\{\int d^4x \left[d^4\theta\, e^{-gV}\Phi_I^\dagger e^{gV}\Phi^I + \frac{1}{4g^2}\left(\int d^2\theta \frac{1}{4} W^\alpha W_\alpha + \mathrm{h.c.}\right) + \right.\right. \\ &+ \left.\left. ig\frac{\sqrt{2}}{3!}\left(\int d^2\theta\, \varepsilon_{IJK}\Phi^I[\Phi^J,\Phi^K] + \int d^2\overline{\theta}\, \varepsilon^{IJK}\Phi_I^\dagger[\Phi_J^\dagger,\Phi_K^\dagger]\right)\right]\right\}, \end{aligned} \tag{2.9}$$

where $d_{\mathbf{r}}$ denotes the Dynkin index of the representation to which the fields belong. To recover the correct dependance on the coupling constant g one must substitute $v \to 2gV$ in the second term in (2.9).

The action (2.9) is non-polynomial as is always the case in supersymmetric gauge theories in $\mathcal{N}$=1 superspace. In particular the scalar potential of (2.5) comes from the superpotential term

$$\begin{aligned} &\int d^4x \left\{\int d^2\theta\, \mathrm{W}(x,\theta,\overline{\theta}) + \mathrm{h.c.}\right\} = \\ &= \int d^4x \left\{\int d^2\theta \left[-i\frac{\sqrt{2}}{3!}\varepsilon_{IJK} f^{abc}(\Phi_a^I \Phi_b^J \Phi_c^K)(x,\theta,\overline{\theta})\right] + \mathrm{h.c.}\right\} \end{aligned}$$

By power expanding e^V one obtains

$$
\begin{aligned}
S = \int d^4x\, d^4\theta\, \Big\{ & V^a(x,\theta,\overline{\theta})\left[-\Box P_T - \xi(P_1+P_2)\Box\right] V_a(x,\theta,\overline{\theta}) + \\
& + \Phi^{\dagger a}_I(x,\theta,\overline{\theta})\Phi^I_a(x,\theta,\overline{\theta}) + igf_{abc}\Phi^{\dagger a}_I(x,\theta,\overline{\theta})V^b(x,\theta,\overline{\theta})\Phi^{Ic}(x,\theta,\overline{\theta}) + \\
& - \frac{1}{2}g^2 f_{ab}{}^e f_{ecd}\Phi^{\dagger a}_I(x,\theta,\overline{\theta})V^b(x,\theta,\overline{\theta})V^c(x,\theta,\overline{\theta})\Phi^{Id}(x,\theta,\overline{\theta}) + \ldots \\
& - \frac{i}{4}gf_{abc}\left[\overline{D}^2\left(D^\alpha V^a(x,\theta,\overline{\theta})\right)\right] V^b(x,\theta,\overline{\theta})\left(D_\alpha V^c(x,\theta,\overline{\theta})\right) + \\
& - \frac{1}{8}g^2 f_{ab}{}^e f_{ecd} V^a(x,\theta,\overline{\theta})\left(D^\alpha V^b(x,\theta,\overline{\theta})\right)\left[\left(\overline{D}^2 V^c(x,\theta,\overline{\theta})\right)\left(D_\alpha V^d(x,\theta,\overline{\theta})\right)\right] + \\
& + \ldots - \frac{\sqrt{2}}{3!}gf^{abc}\left[\varepsilon_{IJK}\Phi^I_a(x,\theta,\overline{\theta})\Phi^J_b(x,\theta,\overline{\theta})\Phi^K_c(x,\theta,\overline{\theta})\delta(\overline{\theta}) + \right. \\
& \left. + \varepsilon^{IJK}\Phi^\dagger{}_{Ia}(x,\theta,\overline{\theta})\Phi^\dagger{}_{Jb}(x,\theta,\overline{\theta})\Phi^\dagger{}_{Kc}(x,\theta,\overline{\theta})\delta(\theta)\right] + \left(\overline{C}'_a(x,\theta,\overline{\theta})C^a(x,\theta,\overline{\theta}) + \right. \\
& \left. - C'_a(x,\theta,\overline{\theta})\overline{C}^a(x,\theta,\overline{\theta})\right) + \frac{i}{2\sqrt{2}}gf_{abc}\left(C'^a(x,\theta,\overline{\theta}) + \overline{C}'^a(x,\theta,\overline{\theta})\right)\cdot \\
& \cdot V^b(x,\theta,\overline{\theta})\left(C^c(x,\theta,\overline{\theta}) + \overline{C}^c(x,\theta,\overline{\theta})\right) - \frac{1}{8}g^2 f_{ab}{}^e f_{ecd}\left(C'^a(x,\theta,\overline{\theta}) + \overline{C}'^a(x,\theta,\overline{\theta})\right)\cdot \\
& \cdot V^b(x,\theta,\overline{\theta})V^c(x,\theta,\overline{\theta})\left(C^d(x,\theta,\overline{\theta}) + \overline{C}^d(x,\theta,\overline{\theta})\right) + \ldots \Big\}\,,
\end{aligned}
\tag{2.10}
$$

which of course can be made polynomial by choosing the Wess–Zumino gauge which implies $V^3 = 0$. In (2.10) the gauge fixing term as well as the corresponding ghost fields have been included following the prescription of (1.67) of chapter 1 and the expansion has been truncated to the terms that will be relevant in future calculations.

Integrating over the θ variables and eliminating the auxiliary fields through the equations of motion results in a different component field formulation, which again has a manifest SU(3)×U(1) global symmetry. Such a formulation will be employed in the next chapter in the discussion of subtleties related to the problem of gauge fixing in this model. Choosing the Wess–Zumino gauge one obtains

$$
\begin{aligned}
S \;=\; & \int d^4x \left\{ -\frac{1}{4}F^a_{\mu\nu}F^{a\,\mu\nu} - D_\mu\varphi^\dagger_I D^\mu\varphi^I + \frac{i}{2}\overline{\lambda}^a \not{D}\lambda^a + \frac{i}{2}\overline{\psi}^a_I \not{D}\psi^{aI} + \right. \\
& + ig\sqrt{2}f_{abc}\left(\overline{\lambda}^a\overline{\varphi}^b_I\psi^{cI} - \overline{\psi}^a_I\varphi^{bI}\lambda^c\right) - \frac{i\sqrt{2}}{2}f_{abc}\left(\varepsilon^I_{JK}\overline{\psi}^a_I\varphi^{bJ}\psi^{cK} + \right. \\
& \left.\left. - \varepsilon^{IJ}_K\overline{\psi}^a_I\varphi^{\dagger b}_J\psi^{cK}\right) - \frac{1}{2}g^2(f_{abc}\varphi^{\dagger a}_I\varphi^b_I)^2 + \frac{1}{2}g^2 f_{abc}f^a_{de}\varepsilon_{IJK}\varepsilon^{LMK}\varphi^{bI}\varphi^{cJ}\varphi^{\dagger d}_L\varphi^{\dagger e}_M \right\}.
\end{aligned}
\tag{2.11}
$$

This formulation can be easily seen to be perfectly equivalent to (2.5) apart from the the manifest R-symmetry, that is only SU(3)×U(1).

It is also useful to give a third component field formulation of $\mathcal{N}$=4 supersymmetric Yang–Mills theory in terms of $\mathcal{N}=2$ multiplets. This approach will actually be used in instanton calculations in chapter 4. The $\mathcal{N}$=4 super Yang–Mills multiplet decomposes in terms of $\mathcal{N}$=2 multiplets into a vector plus a hypermultiplet. In this description the subgroup of the original SU(4) R-symmetry that is realized explicitly is $\mathrm{SU}(2)_{\mathcal{V}}\times\mathrm{SU}(2)_{\mathcal{H}}\times\mathrm{U}(1)$, where the subscripts $\mathcal{V}$ and $\mathcal{H}$ refer to the vector and hypermultiplet respectively: fields in the two multiplets are charged only with respect to one of the two SU(2) factors. The representations of $\mathrm{SU}(2)_{\mathcal{V}}\times\mathrm{SU}(2)_{\mathcal{H}}\times\mathrm{U}(1)$ will be denoted by $(\mathbf{r}_{\mathcal{V}},\,\mathbf{r}_{\mathcal{H}})_q$ with the subscript q referring to the $U(1)$ charge and the following notation for the component fields will be used

$$\begin{aligned}
\mathcal{V} \quad &\rightarrow \quad \lambda^u \in (\mathbf{2},\mathbf{1})_{+1}, \quad \varphi \in (\mathbf{1},\mathbf{1})_{+2}, \quad A_\mu \in (\mathbf{1},\mathbf{1})_0 \\
\mathcal{H} \quad &\rightarrow \quad \psi^{\dot{u}} \in (\mathbf{1},\mathbf{2})_{-1}, \quad q_{u\dot{u}} \in (\mathbf{2},\mathbf{2})_0 \ ,
\end{aligned} \tag{2.12}$$

where $u,\dot{u}=1,2$. In particular the form of the Yukawa couplings in this formulation will be of relevance in the following calculations. In $\mathcal{N}$=2 notation the Yukawa couplings read

$$\begin{aligned}
\mathcal{L}_Y^{(\mathcal{N}=2)} = \frac{\sqrt{2}}{2} g f_{abc} \ \Big\{ & q_S^a \left(\lambda^{b\alpha\, u} \sigma^S{}_{u\dot{u}} \psi_\alpha{}^{c\dot{u}} + \overline{\psi}^b_{\dot{\alpha}\,\dot{u}} \overline{\sigma}^{S\dot{u}u} \overline{\lambda}^{c\dot{\alpha}}_u \right) + \\
& + \ \varphi^a \left(\varepsilon^{uv} \overline{\lambda}^b_{\dot{\alpha}\,u} \overline{\lambda}^{c\dot{\alpha}}_v + \varepsilon_{\dot{u}\dot{v}} \psi^{b\alpha\,\dot{u}} \psi^{c\,\dot{v}}_\alpha \right) + \\
& + \ \overline{\varphi}^a \left(\varepsilon_{uv} \lambda^{b\alpha\,u} \lambda^{c\,v}_\alpha + \varepsilon^{\dot{u}\dot{v}} \overline{\psi}^b_{\dot{\alpha}\,\dot{u}} \overline{\psi}^{c\dot{\alpha}}_{\dot{v}} \right) \Big\} \ ,
\end{aligned} \tag{2.13}$$

In conclusion of this review of the diverse formulations of $\mathcal{N}=4$ super Yang–Mills it is worth recalling that a truly off-shell analysis of the theory, in components and in the Wess–Zumino gauge, was given in [42] within the framework of algebraic renormalization. The problems related to the necessity of including an infinite set of auxiliary fields are dealt with using the method of Batalin and Vilkoviski. This approach, though technically rather involved, allows to prove that all the classical symmetries of the theory survive quantization.

2.2 Classical symmetries and conserved currents

The field content of $\mathcal{N}$=4 supersymmetric Yang–Mills theory described in the preceding section is unique apart from the choice of the gauge group G. In the Abelian case the theory is free as follows immediately by observing that all the fields belong

to the adjoint representation of the gauge group, so that all the covariant derivatives reduce to ordinary ones if G is Abelian, and all the potential terms are proportional to the structure constants f_{abc}. In the non Abelian case the theory has a moduli space of vacua parameterized by the vacuum expectation values (vev's) of the six real scalars, corresponding to a Coulomb phase. The manifold of vacua is determined by the condition of vanishing scalar potential (F-flatness plus D-flatness) which reads

$$\mathrm{tr}\left([\varphi^{AB},\varphi^{CD}][\overline{\varphi}_{AB},\overline{\varphi}_{CD}]\right)=0 \quad\Longleftrightarrow\quad \mathrm{tr}\left([\varphi^{i},\varphi^{j}][\varphi_{i},\varphi_{j}]\right)=0\,.$$

By writing φ^i explicitly in terms of generators of the adjoint representation one obtains

$$\sum_{i,j}^{1,6}\sum_{a=1}^{\dim \mathrm{G}}\varphi_{ia}^2\varphi_{jb}^2\,\mathrm{tr}\left([T^a,T^b]\right)^2=0\,, \tag{2.14}$$

so that for a rank r gauge group G at most r components φ_{ia} for each field can have non vanishing vacuum expectation value. As a result the space of vacua is parameterized by $6r$ real coordinates. The moduli space is

$$\mathcal{M}=\mathbb{R}^{6r}/\mathcal{S}_r\,, \tag{2.15}$$

where $\mathcal{S}_r$ is the group of permutations of r elements, because configurations in which two or more components of φ^i are exchanged are physically indistinguishable. In conclusion the manifold $\mathcal{M}$ is a $6r$-dimensional real space with *orbifold type singularities*.

In a generic point of the Coulomb phase with nonvanishing vev's for the scalar fields the gauge group G is broken to $\mathrm{U}(1)^r$, so that the theory is again Abelian and thus free. On the contrary the origin of the moduli space, where the vev's of all the scalars vanish, corresponds to a phase that is expected to describe a highly non trivial superconformal field theory. Note that the superconformal phase corresponds to an orbifold singularity of the manifold $\mathcal{M}$. In this phase the classical action (2.5) is actually invariant under a larger group of global transformations than that generated by the super Poincaré algebra (1.3), namely the $\mathcal{N}$=4 superconformal group.

The $\mathcal{N}$-extended superconformal algebra in four dimensions is an extension of (1.3) which includes additional generators D, K_μ, S^i_α and $\overline{S}_{\dot\alpha i}$ corresponding respectively to dilatations, special conformal transformations and the associated special supersymmetry transformations. The complete algebra further includes the follow-

ing (anti)commutation relations, supplementing (1.3)

$$
\begin{aligned}
&[M_{\mu\nu}, K_\lambda] = \eta_{\nu\lambda} K_\mu - \eta_{\mu\lambda} K_\nu \\
&[D, P_\mu] = -P_\mu && [D, K_\mu] = K_\mu \\
&[P_\mu, K_\nu] = -2M_{\mu\nu} + 2\eta_{\mu\nu} D && [K_\mu, K_\nu] = 0 \\
&[S_{\alpha i}, M^{\mu\nu}] = (\sigma^{\mu\nu})_\alpha{}^\beta S_{\beta i} && [\overline{S}^i_{\dot\alpha}, M^{\mu\nu}] = (\overline{\sigma})^{\dot\alpha}{}_{\dot\beta} \overline{S}^{\dot\beta i} \\
&\{S_{\alpha i}, \overline{S}^j_{\dot\alpha}\} = 2\sigma^\mu_{\alpha\dot\alpha} K_\mu \delta^j_i \\
&\{S_{\alpha i}, S_{\beta j}\} = 0 && \{\overline{S}^i_{\dot\alpha}, \overline{S}^j_{\dot\beta}\} = 0 \\
&[Q^i_\alpha, D] = \frac{1}{2} Q^i_\alpha && [\overline{Q}_{\dot\alpha i}, D] = \frac{1}{2} \overline{Q}_{\dot\alpha i} \\
&[S_{\alpha i}, D] = -\frac{1}{2} S_{\alpha i} && [\overline{S}^i_{\dot\alpha}, D] = \frac{1}{2} \overline{S}^i_{\dot\alpha} \\
&[Q^i_\alpha, K^\mu] = i\sigma^\mu_{\alpha\dot\alpha} \overline{S}^{\dot\alpha i} && [\overline{Q}^{\dot\alpha}_i, K^\mu] = -i\overline{\sigma}^{\mu\,\dot\alpha\alpha} S_{\alpha i} \\
&[S_{\alpha i}, K^\mu] = 0 && [\overline{S}^{\dot\alpha i}, K^\mu] = 0 \\
&[S_{\alpha i}, P^\mu] = -i\sigma^\mu_{\alpha\dot\alpha} \overline{Q}^{\dot\alpha}_i && [\overline{S}^{\dot\alpha i}, P^\mu] = i\overline{\sigma}^{\mu\,\dot\alpha\alpha} Q^i_\alpha \\
&[S_{\alpha i}, T^a] = -B^{aj}_i S_{\alpha j} && [T^a, \overline{S}^i_{\dot\alpha}] = -B^{\dagger ai}{}_j \overline{S}^j_{\dot\alpha} \\
&\{Q^i_\alpha, S_{\beta j}\} = 2\varepsilon_{\alpha\beta}\delta^i_j D - i(\sigma^{\mu\nu})_\alpha{}^\gamma \varepsilon_{\gamma\beta} M_{\mu\nu}\delta^i_j - 4i\varepsilon_{\alpha\beta}\delta^i_j A + B^{ai}{}_j T_a \\
&\{\overline{Q}_{\dot\alpha i}, \overline{S}^j_{\dot\beta}\} = 2\varepsilon_{\dot\alpha\dot\beta}\delta^j_i D - i\varepsilon_{\dot\alpha\dot\gamma}(\overline{\sigma}^{\mu\nu})^{\dot\gamma}{}_{\dot\beta} M_{\mu\nu}\delta^j_i + 4i\varepsilon_{\dot\alpha\dot\beta}\delta^j_i A + B^{\dagger\,aj}_i T_a \\
&[Q^i_\alpha, A] = -i\left(\frac{4-\mathcal{N}}{4\mathcal{N}}\right) Q^i_\alpha && [\overline{Q}_{\dot\alpha i}, A] = i\left(\frac{4-\mathcal{N}}{4\mathcal{N}}\right) \overline{Q}_{\dot\alpha i} \\
&[S_{\alpha i}, A] = i\left(\frac{4-\mathcal{N}}{4\mathcal{N}}\right) S_{\alpha i} && [\overline{S}^i_{\dot\alpha}, A] = -i\left(\frac{4-\mathcal{N}}{4\mathcal{N}}\right) \overline{S}^i_{\dot\alpha}\,.
\end{aligned}
\tag{2.16}
$$

The operator A in these relations is the generator of the U(1) factor of the group U($\mathcal{N}$)=SU($\mathcal{N}$)×U(1) of the automorphisms of the algebra. The $\mathcal{N}$=4 case is singular since it implies

$$[Q^i_\alpha, A] = [\overline{Q}_{\dot\alpha i}, A] = 0\,.$$

The chiral rotation operator A must therefore have the same action on all the states in the multiplet. This implies in particular that it must act in the same way on states of opposite helicities $-\frac{1}{2}$ and $\frac{1}{2}$, which is possible only if A is actually zero.

For this reason the R-symmetry group in the $\mathcal{N}=4$ case is SU(4) and not U(4) as would be expected from general arguments.

As already noticed in section 2.1 a SU(4)∼SO(6) global symmetry, that is identified with the R-symmetry, naturally arises in the compactification on T^6 that leads to $\mathcal{N}$=4 super Yang–Mills in $D=4$ from the ten dimensional $\mathcal{N}$=1 theory. In this context the reduction of the global symmetry from U(4) to SU(4) is understood as a consequence of the lack of a real six dimensional representation in U(4). More precisely it follows from the form of the Yukawa coupling $\lambda^{\alpha A}[\lambda_\alpha{}^B, \overline{\varphi}_{AB}]$ in the action (2.5). In fact a transformation, under the U(1) factor contained in U(4), of the type $\lambda \to e^{i\alpha}\lambda$ for the spinor would require $\varphi^{AB} \to e^{-2i\alpha}\varphi^{AB}$ for the scalars which is prohibited by the reality condition (2.3).

The Noether currents associated with the superconformal transformations together with those corresponding to chiral SU(4) R-transformations are components of a unique on-shell multiplet [43]. The complete multiplet of currents was given for the Abelian case in [44]. The conserved currents are

$$\begin{aligned}
\mathcal{T}^{\mu\nu} &= \frac{1}{2}[\delta^{\mu\nu}(F^-_{\rho\sigma})^2 - 4F^{-\mu}{}_\rho F^{-\nu\rho} + \text{ h.c.}] - \frac{1}{2}\lambda^{\alpha A}\sigma^{(\mu}{}_{\alpha\dot\alpha}\overleftrightarrow{\partial^{\nu)}}\overline{\lambda}^{\dot\alpha}_A \\
&+ \delta^{\mu\nu}(\partial_\rho\overline{\varphi}_{AB})(\partial^\rho\varphi^{AB}) - 2(\partial^\mu\overline{\varphi}_{AB})(\partial^\nu\varphi^{AB}) \\
&- \frac{1}{3}(\delta^{\mu\nu}\Box - \partial^\mu\partial^\nu)(\overline{\varphi}_{AB}\varphi^{AB}) \\
\Sigma^\mu{}_{\alpha A} &= -\sigma^{\kappa\nu}F^-_{\kappa\nu}\sigma^\mu{}_{\alpha\dot\alpha}\overline{\lambda}^{\dot\alpha}{}_A + 2i\overline{\varphi}_{AB}\overleftrightarrow{\partial^\mu}\lambda_\alpha{}^B + \frac{4}{3}i\sigma^{\mu\nu}{}_\alpha{}^\beta\partial_\nu(\overline{\varphi}_{AB}\lambda_\beta{}^B) \\
\mathcal{J}^\mu{}_A{}^B &= \overline{\varphi}_{AC}\overleftrightarrow{\partial^\mu}\varphi^{CB} + \overline{\lambda}_{\dot\alpha A}\overline{\sigma}^{\mu\dot\alpha\alpha}\lambda_\alpha{}^B - \frac{1}{4}\delta_A{}^B\lambda^{\alpha C}\sigma^\mu{}_{\alpha\dot\alpha}\overline{\lambda}^{\dot\alpha}{}_C\,.
\end{aligned} \tag{2.17}$$

The remaining components of the supermultiplet are obtained by supersymmetry using the equations of motion and read

$$\begin{aligned}
\mathcal{C} &= (F^-{}_{\mu\nu})^2 \\
\hat\Lambda^A_\alpha &= -\sigma^{\mu\nu}{}_\alpha{}^\beta F^-{}_{\mu\nu}\lambda_\beta{}^A \\
\mathcal{E}^{AB} &= \lambda^{\alpha A}\lambda_\alpha{}^B \\
\mathcal{B}_{\mu\nu}{}^{AB} &= \lambda^{\alpha A}\sigma_{\mu\nu\alpha}{}^\beta\lambda_\beta{}^B + 2i\varphi^{AB}F^-{}_{\mu\nu} \\
\hat\chi^C_{\alpha\,AB} &= \frac{1}{2}\varepsilon_{ABDE}(\varphi^{DE}\lambda_\alpha{}^C + \varphi^{CE}\lambda_\alpha{}^D) \\
\mathcal{Q}^{AB}{}_{CD} &= \varphi^{AB}\overline{\varphi}_{CD} - \frac{1}{12}\delta^A{}_{[C}\delta^B{}_{D]}\varphi^{EF}\overline{\varphi}_{EF}\,.
\end{aligned} \tag{2.18}$$

In (2.17) $\mathcal{T}_{\mu\nu}$ is the (improved) energy momentum tensor, $\Sigma^\mu_{\alpha A}$ are the spin $\frac{1}{2}$ supersymmetry currents and $\mathcal{J}^{\mu B}_A$ are the SU(4) currents. $\mathcal{T}_{\mu\nu}$ is a singlet under SU(4)

whereas $\Sigma^{\mu}_{\alpha A}$ transforms in the $\overline{\mathbf{4}}$ and $\mathcal{J}^{\mu B}_{A}$ in the $\mathbf{4}\times\overline{\mathbf{4}}$. The multiplet further includes three scalars $\mathcal{C}$, $\mathcal{E}^{(AB)}$ and $\mathcal{Q}^{ij}$, two fermionic spin $\frac{1}{2}$ components $\hat{\chi}^{C}_{\alpha\, AB}$ and $\hat{\Lambda}^{A}_{\alpha}$ and one antisymmetric tensor $\mathcal{B}^{[AB]}_{\mu\nu}$. Their transformation under SU(4) is as follows: the scalars $\mathcal{C}$, $\mathcal{E}^{(AB)}$ and $\mathcal{Q}^{ij}$ are respectively in the **1**, **10** and **20**, the two fermions $\hat{\chi}^{C}_{\alpha\, AB}$ and $\hat{\Lambda}^{A}_{\alpha}$ belong to the $\mathbf{4}\times\mathbf{6}$ and **4** respectively and $\mathcal{B}^{[AB]}_{\mu\nu}$ is in the **6**.

All of these fields can be obtained as lowest components of an on-shell superfield $\mathcal{W}^{ij}_{(2)}$ which is a bilinear in $W^{i} \equiv \frac{1}{2} t^{i}_{AB} W^{AB}$

$$\mathcal{W}^{ij}_{(2)} = \mathrm{tr}\left(W^{i} W^{j} - \frac{\delta^{ij}}{6} W_{k} W^{k} \right) .$$

In the non Abelian case in (2.17) and (2.18) there is a trace over the colour indices and additional terms are present. Such terms comprise beyond those required for the covariantization of all the derivatives also 'potential' terms. For example the complete non Abelian expression for $\mathcal{E}^{(AB)}$ is

$$\mathcal{E}^{(AB)} = \mathrm{tr}\left(\lambda^{\alpha A} \lambda_{\alpha}{}^{B} \right) + g\, t^{(AB)+}_{[ijk]} \mathrm{tr}\left(\varphi^{i} \varphi^{j} \varphi^{k} \right) , \tag{2.19}$$

where $t^{(AB)+}_{[ijk]}$ is the completely antisymmetrized product of three t^{AB}_{i} matrices.

2.3 Finiteness of $\mathcal{N}$=4 supersymmetric Yang–Mills theory

The first argument suggesting the possibility that the $\mathcal{N}$=4 supersymmetric Yang–Mills theory could be a finite theory at the quantum level, free of ultraviolet as well as infrared divergences, was given in [19]. In that paper within the study of the general problem of the coupling of scalar (chiral) multiplets to vector multiplets it was observed that a number of three chiral multiplets in a supersymmetric Yang–Mills theory would lead to a Gell-Mann and Low β function vanishing in the one-loop approximation. In fact for a gauge group G=SU(N) the one loop β function is given by

$$\beta = -\frac{g^{2}}{16\pi^{2}} (3 - n) N \, ,$$

where n is the number of chiral multiplets coupled to the vector multiplet in the $\mathcal{N}$=1 language.

Successively this result was improved in [45], where the same was shown to hold at the two loop level by an explicit computation in the component field formulation.

Further improvement of this result beyond two loops proves extremely complicated using ordinary techniques based on Feynman rules for the component theory. There exist a computer calculation [46] of the three-loop β function performed by using ordinary Feynman rules which gives a vanishing result as well.

Superspace techniques allow to derive this result in a much simpler way as was shown in [23, 47, 48]. The basic ingredient is the observation that a vertex coupling three chiral superfields is finite as a direct consequence of dimensional analysis. By applying the rules of section 1.7 the superficial degree of divergence of a Green function with three external (anti) chiral superfields is negative, $D = 2 - 3 = -1$, *i.e.* the diagram is finite. This means that the β function is completely determined by the wave function renormalization constant calculated from the divergent part of the propagator of, say, the chiral superfield. The equation for the β function [48] is

$$\beta(g) = \frac{3}{2} g^2 \frac{\partial Z_g^{(1)}}{\partial g} \,,$$

where $Z_g^{(1)}$ is the coefficient of the simple pole term in the Fourier transform of the chiral superfield propagator. Calculations presented in the papers of references [47, 48] show that the propagator of the chiral superfield is finite, in the supersymmetric generalization of the Fermi–Feynman gauge (α=1 in the notation of section 1.6), up to three loops, implying the vanishing of the β function at the same order in perturbation theory.

The lack of manifest $\mathcal{N}$=2 supersymmetry in the calculations described up to now prevents from using the non renormalization theorem for $\mathcal{N}$=2 in order to extend the result to every order. A formulation based on unconstrained $\mathcal{N} = 2$ superfields (which is basically a version of harmonic superspace) was developed in [49]; such an approach allows a description of $\mathcal{N}$=4 super Yang–Mills with manifest $\mathcal{N}$=2 supersymmetry. The finiteness of the theory at all orders in perturbation theory is then a consequence of the $\mathcal{N}$=2 non renormalization theorem once the absence of divergences is proved at one loop.

Another proof of the finiteness of the theory was given in [50] using a light-cone superspace formulation. The advantage of this formulation relies on the fact that only physical fields are involved so that no auxiliary fields are necessary, however the lack of Lorentz invariance introduces other difficulties and makes the interpretation of the results less transparent.

A different approach based on anomaly arguments has been presented in [51].

The vanishing of the β function is obtained as a consequence the superconformal invariance since (see [52])

$$\frac{\beta(g)}{g} F^a_{\mu\nu} F_a^{\mu\nu} = \mathcal{T}_\mu^{\ \mu} \,,$$

so that invariance under dilatations, implying the vanishing of $\mathcal{T}_\mu^{\ \mu}$, immediately gives $\beta(g) = 0$. The problem is therefore the proof of superconformal invariance, which is obtained as a consequence of the SU(4) symmetry being preserved at the quantum level. As already remarked a different demonstration of superconformal symmetry was given in [42] using somewhat more general arguments.

All of the results here discussed either apply to gauge invariant quantities or are obtained avoiding possible subtleties related to the choice of the gauge. As will be discussed in the next chapter subtleties are actually present in the calculation of non gauge invariant quantities such as Green functions both in the component field formulation and in the description in terms of $\mathcal{N}$=1 superfields.

2.4 S-duality

During the last few years the concept of duality has been the basic ingredient for striking results in the analysis of the non-perturbative behaviour of supersymmetric gauge theories as well as of superstring theories. The fundamental idea of duality is that of a transformation relating two different and somehow complementary descriptions of a physical system. The rôle that duality plays in the study of the non-perturbative properties of a physical model relies on the possibility of constructing a transformation that maps the non-perturbative regime of the system into the perturbative regime of the 'dual' system. Examples of this kind of duality have been known for a long time; in particular the duality shown by Kramers and Wannier [53] in the case of the 2-D Ising model and that established between the Sine-Gordon and the Thirring models [54, 55] appear very representative.

The two dimensional Ising model is defined by a set of spin variables σ_i taking values ± 1 and living on a two dimensional square lattice. The interaction is restricted to nearest neighbours and the partition function reads

$$Z(K) = \sum_{\{\sigma\}} \exp\left(K \sum_{(i,j)} \sigma_i \sigma_j \right) \,, \tag{2.20}$$

where K is related to the temperature T and the ferromagnetic coupling constant J through $K = J/k_BT$ (k_B being Boltzmann's constant). In (2.20) the sum on $\{\sigma\}$ is over all spin configurations and the sum on (i, j) is over all pairs of nearest neighbours. The model was explicitly solved by Onsager and exhibits a single phase transition to a ferromagnetic state at a critical temperature T_c. Kramers and Wannier computed the critical temperature by using duality arguments, before the exact solution of the model was found. The crucial point is that the theory can be reformulated in a different ("dual") way by writing the partition function as a sum over the elementary plaquettes of a dual lattice whose sites are the centers of the plaquettes of the original one, with a new coupling K^*. The two descriptions are then equivalent, namely $Z^*(K^*) = Z(K)$, if the couplings are related by

$$\sinh 2K^* = \frac{1}{\sinh 2K} .$$

Notice that this implies that the high temperature (or weak coupling) regime, $K \ll 1$, of the original description is mapped by the duality transformation to the low temperature (or strong coupling) regime, $K^* \gg 1$, of the dual system. The computation of the critical temperature is based on the hypothesis that the system has a single transition point. The critical point must then be the self-dual point of the transformation, $K = K^*$, which gives $\sinh[2J/k_BT_c] = 1$.

A second interesting example of duality occurs in two dimensional relativistic field theory and relates the Sine-Gordon and the Thirring models. The Sine-Gordon model is defined by the 2-D action

$$S_{\text{SG}} = \int d^2x \, \left[\frac{1}{2}\partial_\mu\phi\partial^\mu\phi + \frac{\alpha}{\beta^2}(\cos\beta\phi - 1)\right] , \tag{2.21}$$

where ϕ is a scalar field and the parameter β plays the rôle of a coupling constant as can be established by power expanding the potential term. The spectrum of the theory contains 'meson' excitations of mass $M_m = \sqrt{\alpha}$ and solitonic states with mass $M_s = \frac{8\sqrt{\alpha}}{\beta^2}$. Therefore the solitonic degrees of freedom have large mass in the weak coupling regime. It can be shown that the Sine-Gordon model is completely equivalent to the Thirring model which describes a system of interacting fermions. The action of the latter is

$$S_{\text{T}} = \int d^2x \, \left[\overline{\psi}i\gamma_\mu\partial^\mu\psi + m\overline{\psi}\psi - \frac{g}{2}\overline{\psi}\gamma_\mu\psi\overline{\psi}\gamma^\mu\psi\right] . \tag{2.22}$$

S_{T} can be mapped into the action (2.21) through bosonization [54, 55]. More precisely it can be proved that the duality transformation maps the soliton of the Sine-Gordon

model into the elementary fermion of the Thirring model and the meson states of (2.21) into fermion–anti-fermion bound states. The relation between the coupling constants is

$$\frac{\beta^2}{4\pi} = \frac{1}{1+\frac{g}{\pi}} ,$$

so that in this case as well duality establishes a strong/weak coupling correspondence.

The crucial rôle that duality plays in the context of supersymmetric gauge theories relies on this fundamental feature. The kind of duality that is relevant in four dimensional quantum field theories is a generalization of the electric-magnetic duality introduced for the first time by Dirac in [56]. The general features of electric-magnetic duality are reviewed in appendix B, where the case of the Georgi–Glashow (or Yang–Mills–Higgs) model is described. The application to the $\mathcal{N}$=4 super Yang–Mills theory is discussed in this section. There are various reviews on electric-magnetic duality, see in particular [58, 59, 60].

2.4.1 Montonen–Olive and SL(2,$\mathbb{Z}$) duality

Most of the analysis of this section will be carried out for a SU(2) gauge group, but the results can be generalized to larger groups. The classical spectrum of non Abelian gauge theories can contain non-perturbative states associated with solitonic solutions of the equations of motion. The corresponding field configurations are characterized by a topological charge that is interpreted as a magnetic charge, so that these states describe monopoles and in general dyons (states with both electric and magnetic charge) [61]. As discussed in appendix B the Georgi–Glashow model with Lagrangian

$$\mathcal{L} = -\frac{1}{4}F^a_{\mu\nu}F^{a\mu\nu} + \frac{1}{2}D^\mu\Phi^a D_\mu\Phi_a - \frac{\lambda}{4}\left(\Phi^a\Phi_a - v^2\right)^2 ,$$

possesses, in the (BPS) limit of vanishing potential, a 'classical' spectrum, containing one scalar massive Higgs field, one massless 'photon', massive $W^\pm$ bosons and monopoles $M^\pm$, that is summarized in the following table

	Mass	(Q_e, Q_m)	Spin
Higgs	0	(0,0)	0
Photon	0	(0,0)	1
$W^{\pm}$	ve	$(\pm e,0)$	1
$M^{\pm}$	vg	$(0,\pm g)$	0

where Q_e and Q_m are the electric and magnetic charge operators defined as

$$\begin{aligned} Q_m &= \frac{1}{v}\int d^3x\, D_i\Phi^a B_a^i \\ Q_e &= \frac{1}{v}\int d^3x\, D_i\Phi^a E_a^i\,, \end{aligned} \tag{2.23}$$

with E_a^i and B_a^i the non Abelian electric and magnetic fields respectively. All of the states saturate the Bogomol'nyi bound [62]

$$m \geq v\sqrt{Q_e^2 + Q_m^2}\,, \tag{2.24}$$

where $Q_m = \frac{4\pi n_m}{e}$ and $Q_e = n_e e$, with $n_e, n_m \in \mathbb{Z}$. Of course quantum corrections might in principle modify this situation, however the above picture is expected to be a good approximation in the weak coupling limit, $e \ll 1$. In this limit

$$m_{_W} = ev \ll v\,, \qquad m_{_M} = gv = \frac{4\pi}{e}v \gg v\,.$$

Montonen and Olive have suggested in [37] that a duality transformation exchanging "electric" states associated to $W^{\pm}$ bosons and "magnetic" states associated to monopoles $M^{\pm}$ and at the same time electric and magnetic charges could be an exact symmetry of the full quantum theory in the BPS limit. The exchange of electric and magnetic charges implies, because of the Dirac quantization condition (see appendix B),

$$e \longrightarrow g = \frac{4\pi}{e}\,,$$

so that the proposed symmetry is intrinsically non-perturbative.

The conjecture of [37] is based on the classical spectrum and on the following arguments. First of all the mass formula (2.24) is invariant under duality as can be easily checked. Moreover a semiclassical analysis [63] based on the so called *moduli*

space approximation (to be discussed in the final subsection) shows that the interaction between two monopoles vanishes, whereas there is an attractive interaction between monopoles and anti-monopoles. The duality symmetry would require that the same should be true for the $W^{\pm}$ bosons. It can be proved at the semi-classical level that this is indeed the case because the repulsive interaction between equal charge W bosons is exactly cancelled by an identical contribution due to the exchange of scalars, that are massless in the BPS limit.

There are some basic problems with the Montonen–Olive proposal, namely

- Quantum corrections are expected to spoil the results of the semi-classical analysis. In particular a non-vanishing potential $V(\Phi)$ could be generated.
- The elementary $W^{\pm}$ bosons have spin one while the monopoles that should be related to them by the duality have spin zero, making an exact matching of the quantum numbers impossible.
- The rôle of the dyons has been neglected in the previous analysis.

It will be shown that all of these problems can be solved in supersymmetric theories. In particular the $\mathcal{N}$=4 supersymmetric Yang–Mills theory is supposed to realize exactly a generalization of the Montonen–Olive duality called *S-duality*. This generalization emerges when the effects of a non-vanishing vacuum angle are taken into account. In the presence of a θ-term (see the end of section B.2) the Lagrangian of the Georgi–Glashow model can be written

$$\mathcal{L} = -\frac{1}{32\pi}\mathrm{Im}\,\left[\tau\left(F^{\mu\nu} + i * F^{\mu\nu}\right)\left(F_{\mu\nu} + i * F_{\mu\nu}\right)\right] - \frac{1}{2}D^{\mu}\Phi D_{\mu}\Phi\,, \tag{2.25}$$

with $\tau = \frac{\theta}{2\pi} + i\frac{4\pi}{e^2}$. The electric and magnetic charges of the dyon states saturating the Bogomol'nyi bound are $Q_e = n_e e + n_m \frac{e\theta}{2\pi}$ and $Q_m = n_m \frac{4\pi}{e}$ respectively.

Periodicity in θ requires invariance under

$$\tau \longrightarrow \tau + b\,, \qquad b \in \mathbb{Z}\,, \tag{2.26}$$

which amounts to a shift in the electric charge $n_e \to n_e - b n_m$, $n_m \to n_m$, or in matrix form

$$\begin{pmatrix} n_e \\ n_m \end{pmatrix} \longrightarrow \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}\begin{pmatrix} n_e \\ n_m \end{pmatrix}\,. \tag{2.27}$$

Furthermore at $\theta = 0$ the duality of Montonen and Olive becomes

$$\tau \longrightarrow -\frac{1}{\tau}\,, \tag{2.28}$$

that is equivalent to the exchange of electric and magnetic charges, $e \to g = \frac{4\pi}{e}$, $g \to -e$. In Montonen–Olive duality this is supplemented by the exchange of the corresponding quantum numbers, $Q_e \leftrightarrow Q_m$, implying $n_e \to n_m$, $n_m \to -n_e$. The latter transformation can be written in matrix form as

$$\begin{pmatrix} n_e \\ n_m \end{pmatrix} \longrightarrow \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} n_e \\ n_m \end{pmatrix}. \tag{2.29}$$

It is thus natural to generalize the duality symmetry to the set of transformations generated by (2.26) and (2.28) with arbitrary θ. The resulting group of transformations is SL(2,$\mathbb{Z}$) acting projectively on the parameter τ

$$\tau \longrightarrow \frac{a\tau + b}{c\tau + d}\,, \quad a, b, c, d \in \mathbb{Z}\,, \qquad ad - bc = 1\,. \tag{2.30}$$

The corresponding action on the quantum numbers n_e and n_m, generated by (2.27) and (2.29), is

$$\begin{pmatrix} n_e \\ n_m \end{pmatrix} \longrightarrow \begin{pmatrix} -a & b \\ c & -d \end{pmatrix} \begin{pmatrix} n_e \\ n_m \end{pmatrix}. \tag{2.31}$$

One can then easily check that the mass formula, written in terms of τ as

$$m^2 \geq 4\pi v^2(n_e, n_m)\, \frac{1}{\operatorname{Im}\tau} \begin{pmatrix} 1 & -\operatorname{Re}\tau \\ -\operatorname{Re}\tau & |\tau|^2 \end{pmatrix} \begin{pmatrix} n_e \\ n_m \end{pmatrix} \tag{2.32}$$

is invariant.

2.4.2 S-duality in $\mathcal{N}$=4 supersymmetric Yang–Mills theory

The general properties of the $\mathcal{N}$=4 supersymmetric Yang–Mills theory previously discussed allow to give an intuitive explanation of why it is considered the original example of a theory possessing exact S-duality. In the Coulomb phase for each scalar with non-vanishing vev the same construction leading to the 't Hooft–Polyakov monopole (see the appendix) can be carried on. As a result the classical spectrum contains monopoles and dyons as well. Unlike the case of the Georgi–Glashow model the finiteness of the theory makes the semiclassical analysis of the spectrum sensible even at the quantum level. Moreover the theory has the correct number of fermions in

the adjoint representation to give spin one dyon/monopole configurations through the mechanism sketched in section B.3. More precisely, in the case of a SU(2) gauge group being discussed here, creation and annihilation operators $a_{\pm}^{i\dagger}$ and $a_{\pm}^{i}$, with $i = 1, 2$, associated with the fermionic zero modes can be defined so that the spectrum of zero-energy states, in the charge-one monopole sector, consists of the following states [64]

State	Spin (S_z)
$\|\Omega\rangle$	0
$a_{\pm}^{i\dagger}\|\Omega\rangle$	$\pm\frac{1}{2}$
$a_{-}^{i\dagger}a_{+}^{j\dagger}\|\Omega\rangle$	0
$a_{+}^{1\dagger}a_{+}^{2\dagger}\|\Omega\rangle$	1
$a_{-}^{1\dagger}a_{-}^{2\dagger}\|\Omega\rangle$	-1
$a_{\mp}^{1\dagger}a_{\mp}^{2\dagger}a_{\pm}^{i\dagger}\|\Omega\rangle$	$\mp\frac{1}{2}$
$a_{+}^{1\dagger}a_{+}^{2\dagger}a_{-}^{1\dagger}a_{-}^{2\dagger}\|\Omega\rangle$	0

where $|\Omega\rangle$ is the magnetic charge-one Clifford vacuum. The multiplet generated in this way contains sixteen states: eight bosons (six spin 0 scalars and two spin ±1 vectors) and eight spin $\pm\frac{1}{2}$ fermions. As a result the spin quantum numbers of the monopole multiplet exactly match those of the multiplet of elementary fields.

Furthermore in extended supersymmetric theories the Bogomol'nyi bound comes as a consequence of the supersymmetry algebra, so that it holds in the full quantum theory. In supersymmetric models a vanishing scalar potential in correspondence of non trivial scalar field configurations is perfectly natural because of the presence of flat directions. In the case of $\mathcal{N}$=4 super Yang–Mills this scalar field configuration corresponds to a generic point of the Coulomb phase.

The relation of the BPS limit with the supersymmetry algebra is better understood in the $\mathcal{N}$=2 case. The $\mathcal{N}$=2 supersymmetry algebra possesses two central charges which are related to the magnetic and electric charge operators of the theory [65]. Using a Majorana notation, the $\mathcal{N}$=2 algebra contains the anticommutator

$$\{Q_{\alpha}^{i}, Q_{\beta}^{j}\} = \delta^{ij}\gamma_{\alpha\beta}^{\mu}P_{\mu} + \delta_{\alpha\beta}U^{ij} + (\gamma_5)_{\alpha\beta}V^{ij}\,, \quad i, j = 1, 2\,,$$

where the central charges have been denoted by U^{ij} and V^{ij}. By expressing U^{ij} and V^{ij} in terms of the fields of the theory Olive and Witten have proved the relations

$$U^{ij} = \varepsilon^{ij} v Q_e \,, \qquad V^{ij} = \varepsilon^{ij} v Q_m \,,$$

where Q_e and Q_m are the electric and magnetic charge operators respectively, that generalize those in equation (2.23). In the previous equation $v^2 = \langle \mathrm{tr}\,(S^2)\rangle + \langle \mathrm{tr}\,(P^2)\rangle$, where S and P denote the complex Higgs fields of the model. As a consequence of this relation the Bogomol'nyi bound follows directly from the supersymmetry algebra because of the positive definiteness of the operator $\{Q, Q\}$ in the rest frame.

In the $\mathcal{N}$=4 case the situation is more involved because there are twelve central charges. Correspondingly there are six electric and six magnetic charge operators associated to the six scalars, φ^i, of the $\mathcal{N}$=4 multiplet

$$\begin{aligned} Q_e^i &= \frac{1}{v}\int_{S^2_\infty} d\vec{\sigma}\cdot \mathrm{tr}\left(\vec{E}\varphi^i\right) \\ Q_m^i &= \frac{1}{v}\int_{S^2_\infty} d\vec{\sigma}\cdot \mathrm{tr}\left(\vec{B}\varphi^i\right) \,, \end{aligned} \tag{2.33}$$

where $v^2 = \sum_{i=1}^{6} \langle \mathrm{tr}\,(\varphi^i)^2\rangle$. The Bogomol'nyi bound reads

$$m^2 \geq v^2 \left[(Q_e^i)^2 + (Q_m^i)^2\right]$$

and can be saturated only by states annihilated by exactly one half of the supercharges. The BPS states which saturate the limit correspond to short multiplets of the $\mathcal{N}$=4 superalgebra and for this reason are protected in the full quantum theory. The mass of the BPS states is completely determined by their electric and magnetic quantum numbers and is exact because supersymmetry is preserved at the quantum level.

All the BPS states in the spectrum can be displayed on six two-dimensional lattices (one for each couple of charges Q_e^i and Q_m^i) and lie at points of coordinates

$$\mathcal{Q} = Q_e + iQ_m = \left(n_e e + n_m \frac{e\theta}{2\pi}\right) + i n_m \frac{4\pi}{e} \,.$$

Supersymmetry implies that the BPS states in the spectrum can only decay into a couple of BPS states, so that a generic state of mass m is unstable if and only if there exist BPS states of masses m_1 and m_2 such that

$$m \geq m_1 + m_2 \,.$$



Figure 2.1: Lattice of BPS states

It immediately follows that stable BPS states are all and only those which have integer quantum numbers n_e and n_m that are coprime. These are the states marked by empty circles in figure 2.1, while the full circles denote unstable states.

Thanks to the fact that the BPS states are protected by supersymmetry in the $\mathcal{N}$=4 theory a semi-classical analysis of the spectrum is sensible at the quantum level as well. Some arguments supporting the conjectured S-duality of the $\mathcal{N}$=4 super Yang–Mills theory based on the moduli space approximation are briefly reviewed in the next subsection.

2.4.3 Semiclassical quantization

In appendix B it is discussed how to construct a classical solution of the equations of motion describing a monopole in the Georgi–Glashow model, the aim of this section is to show how the semiclassical quantization of the monopole solution can be used to study the spectrum of the model [63, 66]. The notion of collective coordinates will be introduced starting with the 't Hooft–Polyakov monopole and then the semiclassical quantization will be applied to the $\mathcal{N}$=4 super Yang–Mills case. A readable review of these topics is provided by [67].

In the BPS limit the monopole solution satisfies the Bogomol'nyi equation $B_i^a = D_i\Phi^a$ ($i = 1, 2, 3$, $a = 1, \ldots, \dim \mathrm{G}$). Given a field configuration solving this equation in general one can find a multi-parameter family of solutions with the same energy.

The parameters labeling the degenerate solutions are called *moduli* or *collective coordinates* of the monopole and the corresponding space *moduli space*.

For the 't Hooft–Polyakov monopole described in appendix B three collective coordinates are associated with the translational invariance of the Bogomol'nyi equation. Given a solution $(A_i^{\mathrm{cl}\,a}(\vec{x}), \Phi^{\mathrm{cl}\,a}(\vec{x}))$ the field configuration

$$A_i^{\mathrm{cl}\,a}(\vec{x}+\vec{X})\,, \qquad \Phi^{\mathrm{cl}\,a}(\vec{x}+\vec{X})\,, \tag{2.34}$$

where $\vec{X}$ is a constant vector, still satisfies the equation and has the same energy (and magnetic charge). Thus the three coordinates $\vec{X}$ are moduli.

A fourth collective coordinates is related to gauge transformations. Gauge invariance implies that the functional space of physical field configurations is

$$\mathcal{C} = \mathcal{A}/\mathcal{G}\,,$$

where $\mathcal{A} = (A_i^a, \Phi^a)$ and $\mathcal{G}$ is the group of "small" gauge transformations, *i.e.* gauge transformations that do not reduce to the identity at spatial infinity. "Large", or "global", gauge transformations which do not approach the identity at infinity are true symmetries of the theory and so field configurations related by large gauge transformations are inequivalent. The BPS solution for the 't Hooft–Polyakov monopole is symmetric under the diagonal SO(3) subgroup of $\mathrm{SO(3)_R}\times\mathrm{SU(2)_G}$, where $\mathrm{SO(3)_R}$ is the group of spatial rotations and $\mathrm{SU(2)_G}$ the group of large gauge transformations, but it is not invariant under $\mathrm{SU(2)_G}$. As a result by acting with the U(1) subgroup of $\mathrm{SU(2)_G}$ that is not broken in the Higgs vacuum one generates a new solution with the same energy, so that the fourth collective coordinate is associated with the U(1) group of global 'electromagnetic' transformations. By letting $\vec{X}$ depend on time one describes a moving monopole, $(A_i^{\mathrm{cl}\,a}(\vec{x}+\vec{X}(t)), \Phi^{\mathrm{cl}\,a}(\vec{x}+\vec{X}(t)))$. Since the Hamiltonian for the system in the temporal gauge $A_0 = 0$ is

$$H = T + V = \frac{1}{2}\int d^3x\,, \left[\dot{A}_i^a \dot{A}_a^i + \dot{\Phi}^a \dot{\Phi}_a\right] + \frac{1}{2}\int d^3x\,, \left[B_i^a B_a^i + D_i\Phi^a D^i \Phi_a\right]$$

the time dependence of $\vec{X}$ generates a non vanishing electric field so that the kinetic energy increases while the potential energy remains constant. This is a general feature of a motion in the moduli space. One can then consider a general deformation of a BPS monopole in the $A_0 = 0$ gauge

$$\delta A_i^a(\vec{x}, t)\,, \qquad \delta\Phi^a(\vec{x}, t)\,. \tag{2.35}$$

To keep the potential energy fixed (2.35) must satisfy the linearized Bogomol'nyi equation

$$\varepsilon_{ijk} D^j \delta A^k - D_i \delta\Phi + e[\delta A_i, \Phi^{\rm cl}] = 0 \tag{2.36}$$

and the Gauss law constraint

$$D_i \delta A^i + e[\Phi^{\rm cl}, \delta\Phi] = 0\,. \tag{2.37}$$

The general solution can be written in terms of a single function of time $\chi(t)$

$$\begin{aligned} \delta A_i &= D_i \left(\chi(t)\Phi^{\rm cl}\right) \\ \delta\Phi &= 0 \\ \delta A_0 &= D_0 \left(\chi(t)\Phi^{\rm cl}\right) - \dot\chi(t)\Phi^{\rm cl}\,. \end{aligned} \tag{2.38}$$

If χ is taken to be constant and $(\delta A_i^a, \delta\Phi^a)$ does not contain a translation $\delta\vec{x} = \vec{x} + \vec{X}$, the deformation reduces to a large gauge transformation, so that χ is identified with the fourth collective coordinate. For $\dot\chi \neq 0$ a non-vanishing kinetic energy is generated. Moreover χ is periodic since it parameterizes a compact U(1) (it is a subgroup of $\mathrm{SU}(2)_{\rm G}$), therefore the moduli space with coordinates $(\vec{X}, \chi)$ is topologically $\mathcal{M}_1 = \mathbb{R}^3 \times S^1$.

The previous construction of the moduli space can be generalized to a charge k monopole solution. A rigorous derivation of multimonopole solutions is quite involved [68], the resulting moduli space $\mathcal{M}_k$ is $4k$-dimensional and roughly speaking the $4k$ collective coordinates correspond to the locations of the k monopoles and their dyon degrees of freedom. On physical grounds the existence of multimonopole BPS configurations is related to the fact that the repulsion between monopoles of the same charge can be compensated by an attractive interaction mediated by the massless Higgs field, thus allowing the construction of static configurations.

For a charge k monopole deformations $(\delta_\alpha A_i^a, \delta_\alpha \Phi^a)$, with $\alpha = 1, \ldots, k$, satisfying the linearized Bogomol'nyi equations and the Gauss law constraint define tangent vectors to $\mathcal{M}_k$. Given the tangent vectors the standard metric on $\mathcal{M}_k$ can be constructed as

$$\mathcal{G}_{\alpha\beta} = -\int d^3x\, \mathrm{tr}\left(\delta_\alpha A_i \delta_\beta A^i + \delta_\alpha\Phi\delta_\beta\Phi\right)\,. \tag{2.39}$$

This metric is actually induced by the form of the action for the gauge theory in $\mathcal{C}$ as can be seen by writing the monopole solution as $A_i^{\rm cl}(\vec{x}, z_\alpha)$, $\Phi^{\rm cl}(\vec{x}, z_\alpha)$, where z_α's

denote the moduli. The tangent vectors $(\delta_\alpha A_i, \delta_\alpha \Phi)$ are calculated by differentiating with respect to the z_α. In general one has to include a gauge transformation to ensure that $(\delta_\alpha A_i, \delta_\alpha \Phi)$ are tangent to the moduli space of physical fields, *i.e.* that the fields $A_i' = A_i^{\rm cl} + \delta_\alpha A_i \delta z^\alpha$, $\Phi' = \Phi^{\rm cl} + \delta_\alpha \Phi \delta z^\alpha$ belong to the space $\mathcal{C}$. The general form of the tangent vectors is

$$\begin{aligned} \delta_\alpha A_i &= \frac{\partial A_i}{\partial z^\alpha} - D_i \epsilon_\alpha \\ \delta_\alpha \Phi &= \frac{\partial \Phi}{\partial z^\alpha} - e[\Phi^{\rm cl}, \epsilon_\alpha]\,, \end{aligned}$$

where $\epsilon_\alpha(\vec{x}, z^\beta)$ is a suitable gauge parameter that is fixed by requiring that the linearized Bogomol'nyi equation (2.36) and the Gauss law constraint (2.37) be satisfied. The action of the theory on the BPS configuration is then obtained after substituting

$$\begin{aligned} A_i(\vec{x},t) &\longrightarrow A_i^{\rm cl}(\vec{x}, z^\alpha(t)) \\ A_0 &\longrightarrow \dot{z}^\alpha(t)\epsilon_\alpha \\ \Phi(\vec{x},t) &\longrightarrow \Phi^{\rm cl}(\vec{x}, z^\alpha(t)) \end{aligned} \tag{2.40}$$

and reads

$$S = -\frac{1}{2}\int d^3x dt\, {\rm tr}\left(F_{0i}^{\rm cl} F_{\rm cl}^{0i}\right) = \int dt\, \mathcal{G}_{\alpha\beta}\dot{z}^\alpha \dot{z}^\beta - \frac{4\pi v}{e}k\,. \tag{2.41}$$

The action (2.41) is the starting point for the semiclassical quantization which is achieved by studying the corresponding quantum mechanics.

In the $k=1$ case the moduli are $z^\alpha = (\vec{X}, \chi)$ and the action becomes

$$S = \frac{1}{2}\int dt\, \left[\frac{4\pi v}{e}\dot{\vec{X}}^2 + \frac{4\pi}{v e^2}\dot{\chi}^2\right] - \frac{4\pi v}{e}\,. \tag{2.42}$$

The resulting wave functions are plane waves

$$\psi = e^{i\vec{P}\cdot\vec{X}} e^{i n_e \chi}\,,$$

with $n_e \in \mathbb{Z}$, and one can compute the electric charge, $Q_e = -ie\partial_\chi$, so that the quantum mechanical system possesses an infinite tower of states with $Q_e = n_e e$. For the mass of these states one gets

$$m = \frac{n_e^2 v e^3}{8\pi} + \frac{4\pi v}{e} \approx v[Q_e^2 + Q_m^2]\,,$$

showing that the Bogomol'nyi bound is saturated.

The preceding arguments can be extended to the case of the $\mathcal{N}$=4 super Yang–Mills theory to find stronger evidence for the exact S-duality of the model. In order to be able to use the previous considerations the simplest case will be considered, namely the gauge group will be taken to be G=SU(2), broken by the vacuum configuration

$$\begin{aligned}\langle\varphi^2\rangle = \langle\varphi^3\rangle = \ldots = \langle\varphi^6\rangle = 0\\ \langle\varphi^1\rangle = \Phi\,, \qquad \langle \mathrm{tr}\Phi^2\rangle = v^2\,.\end{aligned}$$

The perturbative states in the bosonic sector of the theory, labeled by quantum numbers (n_m, n_e) consist of a massless photon multiplet with charge $(0,0)$ and massive $W_\pm$ boson multiplets with charge $(0,\pm 1)$. The tower of stable BPS states predicted by S-duality have charges (l,k) (with l and k coprimes) and can be studied within the framework of semiclassical quantization since they can be put in correspondence with certain geometric structures on $\mathcal{M}_k$. Then the non-renormalization properties of the theory make the semiclassical result reliable in the full quantum theory.

In the Georgi–Glashow model $\mathcal{M}_k$ is parameterized by $4k$ collective coordinates which are bosonic zero modes. In the $\mathcal{N}$=4 supersymmetric Yang–Mills theory there are fermionic zero modes as well in the monopole background and the index theorem of [71] shows that the number of such zero modes in the charge k sector is precisely $4k$. As a consequence the moduli space has fermionic coordinates too and the low energy ansatz to be substituted into the Bogomol'nyi equation is given by (2.40) supplemented by

$$\lambda(\vec{x},t) \sim \psi(t)\lambda^{\mathrm{cl}}(\vec{x}, z_\alpha(t)) \tag{2.43}$$

for the Weyl spinors in the monopole background. In equation (2.43) the ansatz for the spinors $\lambda(\vec{x},t)$ has been schematically written as the product of a c-number $\lambda^{\mathrm{cl}}(\vec{x}, z_\alpha)$ satisfying a zero mode equation in the monopole background and a time dependent fermionic collective coordinate $\psi(t)$. The zero modes can be shown to constitute a short multiplet preserving half of the supersymmetries just like is expected for BPS states. The exact ansatz for the fields is rather complicated and was discussed in detail in [69]. A technically involved computation allows to derive the quantum mechanical action to be employed in the semiclassical analysis. The action reads

$$S = \frac{1}{2}\int dt\, \left[\mathcal{G}_{\alpha\beta}\left(\dot{z}^\alpha\dot{z}^\beta + i\overline{\psi}^\alpha\gamma^0 D_t\psi^\beta\right) + \frac{1}{6}R_{\alpha\beta\gamma\delta}\overline{\psi}^\alpha\psi^\gamma\overline{\psi}^\beta\psi^\delta\right] - \frac{4\pi v}{e^2}k\,, \tag{2.44}$$

where a Majorana notation is used. In (2.44) $D_t\psi^\alpha = \dot{\psi}^\alpha + \Gamma^\alpha_{\beta\gamma}\dot{z}^\beta\psi^\gamma$ is the covariant derivative, with $\Gamma^\alpha_{\beta\gamma}$ the Christoffel symbols of the metric $\mathcal{G}_{\alpha\beta}$, and $R_{\alpha\beta\gamma\delta}$ is the corresponding Riemann tensor [69]. It can be shown that thanks to the fact that the monopole moduli space is a hyper-Kähler manifold, S in equation (2.44) is invariant under the action of eight real supercharges. The quantization of the supersymmetric quantum mechanics (2.44) was performed in [70], where it was shown that the states can be put in one to one correspondence with differential forms on $\mathcal{M}_k$ and the Hamiltonian of the system H can be expressed in terms of the exterior derivative acting on forms. This construction leads to a basis of sixteen independent forms that are eigenstates of H in the $n_m = k = 1$ sector. These states constitute a BPS multiplet as can be verified by a computation of the corresponding quantum numbers. The complete wave functions contain a plane wave factor from the bosonic sector of the form $e^{i\vec{P}\cdot\vec{X}}e^{in_e\chi}$ so that again one finds a tower of states with $Q_e = n_e + \frac{e\theta}{2\pi}$. These states all saturate the Bogomol'nyi bound and therefore are exactly those predicted by S-duality.

For $n_m = k > 1$ the problem is much more complicated. The moduli space is

$$\mathcal{M}_k = \mathbb{R}^3 \times (S^1 \times \tilde{\mathcal{M}}^0_k)/\mathbb{Z}_k\,,$$

where $\tilde{\mathcal{M}}^0_k$ is a $4(k-1)$-dimensional hyper-Kähler manifold. The states are tensor products of forms on $\mathbb{R}^3 \times S^1$ with forms on $\tilde{\mathcal{M}}^0_k$, $|s\rangle = |\omega, n_e\rangle \otimes |\alpha\rangle$ [2]. The forms $|\alpha\rangle$ can be proved to be unique and much in the same way as in the $k = 1$ case there are sixteen forms $|\omega, n_e\rangle$ for all k that give rise to a BPS multiplet. The resulting states carry electric charge $Q_e = n_e e + n_m \frac{e\theta}{2\pi}$ and saturate the Bogomol'nyi bound. The unique form $|\alpha\rangle$ for the case $k = 2$ has been explicitly determined by Sen in [72]. Arguments suggesting a possible generalization to $k > 2$ were given in [73].

[2] There are actually further subtleties related to the $\mathbb{Z}_k$ identification.

Chapter 3

Perturbative analysis of $\mathcal{N} = 4$ supersymmetric Yang–Mills theory

As discussed in the previous chapter, the $\mathcal{N}$=4 supersymmetric Yang–Mills theory has been proved to be finite at the perturbative level up to three loops and strong arguments have been proposed in order to extend this result to all orders and even non perturbatively. For this reason the theory has long been considered to be rather trivial. However recent developments, within the context of the correspondence with type IIB superstring theory on anti de Sitter space, that will be studied in the final chapter, have shown that it is actually an extremely interesting superconformal field theory, possessing very peculiar properties.

Even in perturbation theory a careful analysis allows to point out various problems mostly related to the choice of gauge. Both in the component field formulation and using $\mathcal{N}$=1 superfields the gauge fixing procedure appears very subtle. As will be discussed in detail in both cases one finds divergences in off-shell Green functions.

In the component formulation the propagators of the elementary fields are ultraviolet divergent in the Wess–Zumino (WZ) gauge and these infinities are exactly cancelled when the contributions of the gauge-dependent fields, that are put to zero in the WZ gauge, are taken into account. The choice of the WZ gauge, that is almost unavoidable in explicit computations, introduces divergences that require a wave function renormalization. This is a general result that applies to theories with less supersymmetry as well, but appears particularly dramatic in the $\mathcal{N}$=4 case be-

cause the theory is finite, so that the divergences cannot be reabsorbed through a redefinition of the bare parameters.

Different problems emerge in the formulation of the theory in terms of $\mathcal{N}$=1 superfields. Almost all the calculations showing the vanishing of the quantum corrections to two- and three-points functions that are presented in the literature were performed in the supersymmetric generalization of the Fermi–Feynman gauge (α=1 in the notation of section 1.6). With a different choice of gauge two and three point functions develop infrared singularities of difficult interpretation, leading to the result that the choice α=1 is somehow privileged. This conclusion was proposed for the first time in [74] and then it was discussed in the case of the $\mathcal{N}$=4 theory in [75]; however no possible explanation was proposed. Unlike those of the elementary fields the correlation functions of gauge invariant composite operators, that play a crucial rôle in the correspondence with AdS type IIB supergravity/superstring theory, should not suffer from problems related to the gauge fixing.

Also, it will be discussed, in the superfield formulation, the effect of the introduction of a mass term for the chiral superfields, which breaks supersymmetry from $\mathcal{N}$=4 down to $\mathcal{N}$=1. It will be shown that this deformation of the model does not modify the ultraviolet properties of the original theory. This result was first proposed in [76], where it was proved that the inclusion of mass terms for the (anti) chiral superfields does not generate divergent corrections to the effective action. It will be argued that this statement can be reinforced, showing that, at least at one loop, no new divergences, not even corresponding to wave function renormalizations, appear as a consequence of the addition of the mass terms. This result supports the claim put forward in [77, 78], where the 'mass deformed' $\mathcal{N}$=4 theory was proposed as a supersymmetry-preserving regularization scheme for a class of $\mathcal{N}$=1 theories. It will also be argued that the same approach should work in the case of $\mathcal{N}$=2 theories.

All of this problems are examined in detail in this chapter. Section 3.1 deals with the difficulties introduced by the choice of the Wess–Zumino gauge when the component formalism is used. The subsequent sections report calculations performed in the $\mathcal{N}$=1 superfield formalism of two-, three- and four-point functions. The material presented in this chapter is an original and systematic re-analysis of the perturbative properties of $\mathcal{N}$=4 supersymmetric Yang–Mills theory, the treatment follows closely [79].

3.1 Perturbation theory in components: problems with the Wess–Zumino gauge

Throughout this chapter calculations will be carried on in Euclidean space-time and unless otherwise stated the gauge group will be taken to be SU(N). To discuss perturbation theory in components in this section the formulation of equation (2.11), with a SU(3)×U(1) subgroup of the SU(4) R-symmetry group manifest, will be employed. To correctly deal with the lower components of the vector superfield V, that are put to zero in the Wess–Zumino gauge, it is however useful not to eliminate the auxiliary fields F and D through their equations of motion, but rather keep them in the action: in the computation of Green functions the corresponding x-space propagators are simply δ-functions. The complete expression for the vector superfield V was given in equation (1.24) and contains beyond the physical fields, A_μ and λ, and the auxiliary field, D, the 'gauge-dependent' fields C, χ and S. Note that the scalar field C and the spinor χ have "wrong" physical dimension, resulting in non standard free propagators, as will be shown. The non Abelian field strength superfield W_α is

$$W_\alpha = -\frac{1}{4}\overline{DD}e^{-V}D_\alpha e^V = \sum_{k=1}^{\infty} W_\alpha^{(k)}\,, \tag{3.1}$$

where

$$\begin{aligned} W_\alpha^{(1)} &= -\frac{1}{4}\overline{DD}D_\alpha V \\ W_\alpha^{(2)} &= \frac{1}{8}\overline{DD}[V, D_\alpha V] \end{aligned} \tag{3.2}$$

and the terms $W_\alpha^{(k)}$, with $k \geq 3$, contain k factors of V and vanish in the WZ gauge.

The Euclidean action in $\mathcal{N}$=1 superfields can be written

$$\begin{aligned} S^{(\mathrm{E})} = \frac{1}{d_{\mathbf{r}}} tr \int d^4x\, d^4\theta \, \Big\{ -\Phi_I^\dagger \Phi^I - \frac{1}{4g^2}\Big[\frac{1}{4} W^{(1)\alpha} W_\alpha^{(1)} \delta(\overline{\theta}) + \frac{1}{4}\overline{W}_{\dot\alpha}^{(1)} \overline{W}^{(1)\dot\alpha} \delta(\theta) + \\ - \frac{1}{8\alpha}\overline{D}^2 V D^2 V \Big] - g[\Phi_I^\dagger V]\Phi^I - \Big[\Big(\frac{1}{8g^2} W^{(1)\alpha} W_\alpha^{(2)} + \frac{1}{16g^2} W^{(2)\alpha} W_\alpha^{(2)} \Big) \delta(\overline{\theta}) + \\ + \Big(\frac{1}{8g^2} \overline{W}_{\dot\alpha}^{(1)} \overline{W}^{(2)\dot\alpha} + \frac{1}{16g^2} \overline{W}_{\dot\alpha}^{(2)} \overline{W}^{(2)\dot\alpha} \Big) \delta(\theta) \Big] - \frac{1}{2} g^2 [V, [V, \Phi_I^\dagger]] \Phi^I + \dots \Big\}\,, \end{aligned} \tag{3.3}$$

where the gauge fixing term has been included, whereas no ghost term is displayed since it will not be relevant for the computations to be discussed in this section. In equation (3.3) the dots denote terms of higher order in V, which do not contribute

to the Green functions that will be considered and that will be suppressed from now on. In the following calculations the Fermi–Feynman gauge, α=1, will be used. With this choice one obtains [1]

$$\begin{aligned} S &= \frac{1}{d_{\mathbf{r}}}\mathrm{tr}\int d^4x\, d^4\theta\, \left\{\left[V\Box V - \Phi_I^\dagger \Phi^I\right] - \left[\left(\frac{1}{8g^2}W^{(1)\alpha}W^{(2)}_\alpha + \frac{1}{16g^2}W^{(2)\alpha}W^{(2)}_\alpha\right)\delta(\overline{\theta}) + \right.\right. \\ &+ \left.\left.\left(\frac{1}{8g^2}\overline{W}^{(1)}_{\dot\alpha}\overline{W}^{(2)\dot\alpha} + \frac{1}{16g^2}\overline{W}^{(2)}_{\dot\alpha}\overline{W}^{(2)\dot\alpha}\right)\delta(\theta)\right] - \left[-g[\Phi_I^\dagger V]\Phi^I + \frac{1}{2}g^2[V,[V,\Phi_I^\dagger]]\Phi^I\right]\right\}, \end{aligned}$$

where from the definition (3.2) it follows

$$\begin{aligned} W^{(1)}_\alpha &= -i\lambda_\alpha + \left[\delta^\beta_\alpha D - \frac{1}{2}\delta^\beta_\alpha \Box C - \frac{i}{2}(\sigma^\mu\overline{\sigma}^\nu)_\alpha{}^\beta\,(\partial_\mu A_\nu - \partial_\nu A_\mu)\right]\theta_\beta + \\ &\quad + \theta\theta\sigma^\mu_{\alpha\dot\alpha}\partial_\mu\overline{\lambda}^{\dot\alpha}\,. \end{aligned} \tag{3.4}$$

The explicit form of $W^{(2)}_\alpha$ will not be necessary for the moment (see however equation (3.18)). Expansion of the action in components using the complete expression of V gives

$$S = S_0 + S_{\mathrm{int}}\,.$$

The free action S_0 comes from the terms $-V\Box V$ and $\Phi_I^\dagger\Phi^I$ and reads

$$\begin{aligned} S_0 &= \int d^4x\, \left\{\left[(\partial_\mu\varphi_I^{a\dagger})(\partial^\mu\varphi^I_a) + \overline{\psi}^a_I\overline{\sigma}^\mu(\partial_\mu\psi^I_a) + F_I^{a\dagger}F^I_a\right] + \left[S_a^\dagger\Box S^a + \right.\right. \\ &-\frac{1}{2}C^a\Box D_a - \frac{1}{2}D^a\Box C_a - \frac{1}{2}C^a\Box^2 C_a + \frac{1}{2}\chi^a\Box\lambda_a + \frac{1}{2}\overline{\chi}^a\Box\overline{\lambda}_a + \frac{1}{2}\lambda^a\Box\chi_a + \\ &\left.\left.+\frac{1}{2}\overline{\lambda}^a\Box\overline{\chi}_a + \frac{1}{2}\chi^a\Box\sigma^\mu(\partial_\mu\overline{\chi}_a) + \frac{1}{2}\overline{\chi}^a\Box\overline{\sigma}^\mu(\partial_\mu\chi_a) + \frac{1}{2}(\partial_\mu A^a_\nu)(\partial_\nu A_{a\mu})\right]\right\}. \end{aligned} \tag{3.5}$$

The interaction part S_{int} contains an infinite number of terms. The propagators of the fermion ψ and of the scalar φ belonging to the $\mathcal{N}$=1 chiral multiplet will now be computed at one loop. The terms that are relevant for these calculations come from the expansion of $\Phi^\dagger V\Phi$ and $\Phi^\dagger V^2\Phi$ in the superfield action. The latter generates tadpole type diagrams in the propagator $\langle\varphi^\dagger\varphi\rangle$, but not in $\langle\overline{\psi}\psi\rangle$. The interaction

[1] From now on the superscript E will be suppressed. Euclidean signature is understood unless otherwise stated.

part of the action to be considered is

$$
\begin{aligned}
S_{\rm int} = \int d^4x \Bigg\{ & ig f_{abc} \left[\frac{1}{2}\phi_I^{a\dagger} D^b \phi^{cI} + \frac{1}{2}\phi_I^{a\dagger}(\Box C^b)\phi^{cI} - \frac{1}{2}(\partial_\mu \phi_I^{a\dagger}) C^b (\partial^\mu \phi^{cI}) + \right. \\
& + \frac{i}{2}\left(\phi_I^{a\dagger} A_\mu^b (\partial^\mu \phi^{cI}) - (\partial^\mu \phi_I^{a\dagger}) A_\mu^b \phi^{cI} \right) + \frac{i}{\sqrt{2}}\left(\phi_I^{a\dagger} S^{b\dagger} F^{cI} - F_I^{a\dagger} S^b \phi^{cI} \right) + \\
& - F_I^{a\dagger} C^b F^{cI} + \frac{i}{\sqrt{2}}\left(\overline{\psi}_I^a \overline{\lambda}^b \phi^{cI} - \phi_I^{a\dagger} \lambda^b \psi^{cI} \right) + \frac{i}{\sqrt{2}}\left(F_I^{a\dagger} \chi^b \psi^{cI} - \overline{\psi}_I^a \overline{\chi}^b F^{cI} \right) + \\
& - \frac{i}{\sqrt{2}}\left(\phi^{a\dagger} (\partial_\mu \overline{\chi}^b) \sigma^\mu \psi^{cI} + \overline{\psi}_I^a \overline{\sigma}^\mu (\partial_\mu \chi^b) \phi^{cI} \right) + \frac{1}{2}\left(C^b \overline{\psi}_I^a \overline{\sigma}^\mu (\partial_\mu \psi^{cI}) - C^b (\partial_\mu \overline{\psi}_I^a) \overline{\sigma}^\mu \psi^{cI} \right) + \\
& \left. - \frac{i}{2}\overline{\psi}_I^a \overline{\sigma}^\mu \psi^{cI} A_\mu^b \right] - \frac{\sqrt{2}}{3!} g f_{abc} \left[\varepsilon^{IJK}\left(\phi_I^{a\dagger} \phi_J^{b\dagger} F_K^{c\dagger} - \phi_I^{a\dagger} \overline{\psi}_J^b \overline{\psi}_K^c \right) + \varepsilon_{IJK}\left(\phi^{aI} \phi^{bJ} F^{cK} + \right. \right. \\
& \left. \left. - \phi^{aI} \psi^{bJ} \psi^{cK} \right) \right] - \frac{g^2}{2} f_{abe} f^e{}_{cd} \Bigg(\Bigg[- C^b \psi^{dI} \sigma^\mu \overline{\psi}_I^a A_\mu^c + \\
& + (\overline{\chi}^c \overline{\psi}_I^d)(\chi^b \psi^{aI}) + \frac{i}{2}\left(\partial_\mu \psi^{dI} \sigma^\mu \overline{\psi}_I^a - \psi^{dI} \sigma^\mu \partial_\mu \overline{\psi}_I^a \right) C^b C^c \Bigg] + \varphi_I^{a\dagger} \Bigg[C^b \left(D^c + \right. \\
& \left. + \frac{1}{2}\Box C^c \right) - \chi^b \left(\lambda^c + \frac{i}{2}\sigma^\mu \partial_\mu \overline{\chi}^c \right) - \overline{\chi}^b \left(\overline{\lambda}^c + \frac{i}{2}\sigma^\mu \partial_\mu \chi^c \right) + S^{b\dagger} S^c - \\
& \left. - \frac{1}{2} A_\mu^{bc} A^{c\mu} \Bigg] \varphi^{dI} + \frac{i}{2} C^b A^{c\mu} \left(\varphi_I^{a\dagger} (\partial_\mu \varphi^{dI}) - (\partial_\mu \varphi_I^{a\dagger}) \varphi^{dI} \right) + \frac{1}{4} C^b C^c \left(\varphi_I^{a\dagger} (\Box \varphi^{dI}) + \right. \\
& \left. + (\Box \varphi_I^{a\dagger}) \varphi^{dI} \right) + \frac{i}{2} \chi^b \sigma^\mu \overline{\chi}^c \left(\varphi_I^{a\dagger} (\partial_\mu \varphi^{dI}) - (\partial_\mu \varphi_I^{a\dagger}) \varphi^{dI} \right) \Bigg) \Bigg\} . \qquad (3.6)
\end{aligned}
$$

The quadratic part of the action, S_0, can be written in a more compact form introducing the notation

$$
\mathcal{B}^a(x) = \begin{pmatrix} C^a(x) \\ D^a(x) \end{pmatrix} , \qquad \mathcal{F}^a(x) = \begin{pmatrix} \chi^a(x) \\ \overline{\chi}^a(x) \\ \lambda^a(x) \\ \overline{\lambda}^a(x) \end{pmatrix} , \qquad (3.7)
$$

so that

$$
\begin{aligned}
S_0 \;=\; & \int d^4x \left\{ \left[(\partial_\mu \varphi_I^{a\dagger})(\partial^\mu \varphi_a^I) + \overline{\psi}_I^a \overline{\sigma}^\mu (\partial_\mu \psi_a^I) + F_I^{a\dagger} F_a^I \right] + \right. \\
& \left. + \left[S_a^\dagger \Box S^a - \frac{1}{2} A_\mu^a A_a^\mu + \mathcal{B}_a^T M \mathcal{B}^a + \mathcal{F}_a^T N \mathcal{F}^a \right] \right\} , \qquad (3.8)
\end{aligned}
$$

where

$$
M = \frac{1}{2}\begin{pmatrix} -\Box^2 & -\Box \\ -\Box & 0 \end{pmatrix} , \qquad N = \frac{1}{2}\begin{pmatrix} 0 & \Box\sigma^\mu\partial_\mu & -\Box & 0 \\ \Box\overline{\sigma}^\mu\partial_\mu & 0 & 0 & -\Box \\ -\Box & 0 & 0 & 0 \\ 0 & -\Box & 0 & 0 \end{pmatrix} . \qquad (3.9)
$$

Inverting the kinetic matrices M and N one gets the free propagators. From (3.9) it follows

$$M^{-1} = \begin{pmatrix} 0 & \frac{1}{\Box} \\ \frac{1}{\Box} & -1 \end{pmatrix}, \quad N^{-1} = \begin{pmatrix} 0 & 0 & -\frac{1}{\Box} & 0 \\ 0 & 0 & 0 & -\frac{1}{\Box} \\ -\frac{1}{\Box} & 0 & 0 & \frac{\sigma^\mu \partial_\mu}{\Box} \\ 0 & -\frac{1}{\Box} & \frac{\overline{\sigma}^\mu \partial_\mu}{\Box} & 0 \end{pmatrix}, \tag{3.10}$$

so that the free propagators are



$$\longrightarrow \langle \varphi_J^{b\dagger}(x) \varphi_a^I(y) \rangle_{\text{free}} = -\frac{\delta_a^b \delta_J^I}{\Box} \delta(x-y) = \Delta_{aJ}^{bI}(x-y)$$



$$\longrightarrow \langle S^{b\dagger}(x) S_a(y) \rangle_{\text{free}} = \frac{\delta_a^b}{\Box} \delta(x-y) = 2\Delta_a^b(x-y)$$



$$\longrightarrow \langle C^b(x) D_a(y) \rangle_{\text{free}} = \frac{\delta_a^b}{\Box} \delta(x-y) = \Delta_a^b(x-y)$$

$$\longrightarrow \langle D^b(x) D_a(y) \rangle_{\text{free}} = -\delta_a^b \delta(x-y)$$



$$\longrightarrow \langle F_J^{b\dagger}(x) F_a^I(y) \rangle_{\text{free}} = \delta_J^I \delta_a^b \delta(x-y)$$

$$\longrightarrow \langle A_\mu^b(x) A_{a\nu}(y) \rangle_{\text{free}} = -\frac{\delta_{\mu\nu} \delta_a^b}{\Box} \delta(x-y) = \Delta_{a\mu\nu}^b(x-y)$$



$$\longrightarrow \langle \chi^{b\alpha}(x) \lambda_a^\beta(y) \rangle_{\text{free}} = -\frac{\varepsilon^{\alpha\beta} \delta_a^b}{\Box} \delta(x-y) = R_a^{b\alpha\beta}(x-y)$$



$$\longrightarrow \langle \overline{\chi}^{b\dot{\alpha}}(x) \overline{\lambda}_a^{\dot{\beta}}(y) \rangle_{\text{free}} = -\frac{\varepsilon^{\dot{\alpha}\dot{\beta}} \delta_a^b}{\Box} \delta(x-y) = \overline{R}_a^{b\dot{\alpha}\dot{\beta}}(x-y)$$



$$\longrightarrow \langle \overline{\lambda}^{b\dot{\alpha}}(x) \lambda_a^\alpha(y) \rangle_{\text{free}} = \frac{\delta_a^b \overline{\sigma}_{\dot{\alpha}\alpha}^\mu \partial_\mu}{\Box} \delta(x-y) = \overline{S}_{a\dot{\alpha}\alpha}^b(x-y)$$

$$\dot{\alpha}, I, b \longrightarrow \alpha, J, a \qquad \longrightarrow \quad \langle \overline{\psi}_I^{b\dot{\alpha}}(x)\psi_a^{\alpha J}(y)\rangle_{\text{free}} = \frac{\delta_a^b \delta_I^J \overline{\sigma}^\mu_{\dot{\alpha}\alpha}\partial_\mu}{\Box}\delta(x-y) = \\ = \overline{S}^{bJ}_{aI\dot{\alpha}\alpha}(x-y)$$

In conclusion beyond the ordinary propagators for the physical fields, φ, ψ, λ and A_μ, and those for the auxiliary fields, F and D, one further obtains the propagators $\langle S^\dagger S\rangle$, $\langle CD\rangle$, $\langle \chi\lambda\rangle$ and $\langle \overline{\chi}\overline{\lambda}\rangle$. The latter are not present in the Wess–Zumino gauge.

3.1.1 One loop corrections to the propagator of the fermions belonging to the chiral multiplet

The one-loop correction to the propagator of the fermions ψ^I in the chiral multiplet is the simplest to compute. In the WZ gauge there are three contributions at the one loop level, that will be shown to lead to a logarithmically divergent result.

From the action (3.5), (3.6), with $C{=}\chi{=}S{=}0$ one obtains the following three diagrams



The three contributions depicted above can be easily evaluated and give the results

$$A^{\dot{\alpha}\alpha aI}_{bJ}(x;y) = -\frac{1}{2}g^2 f^d{}_{ef} f^l{}_{mn}\int d^4x_1 d^4x_2 \left\{\Delta^{me}_{\nu\mu}(x_2-x_1)\left[\overline{S}^{\dot{\alpha}\beta aI}_{lL}(x-x_2)\sigma^\nu_{\beta\dot{\gamma}}\cdot \right.\right. \\ \left.\left.\cdot\overline{S}^{\dot{\gamma}\gamma nL}_{dK}(x_2-x_1)\sigma^\mu_{\gamma\dot{\beta}}\overline{S}^{\dot{\beta}\alpha fK}_{bJ}(x_1-y)\right]\right\}, \qquad (3.11)$$

$$B^{\dot\alpha\alpha aI}_{bJ}(x;y) = -\frac{1}{2}g^2\varepsilon^{LMN}\varepsilon^{PQ}{}_R f_{de}{}^f f_{lmn}\int d^4x_1 d^4x_2 \left\{\Delta^{ld}_{PL}(x_2-x_1)\cdot \right.$$
$$\left.\cdot\left[\overline{S}^{\dot\alpha\beta aI}_{fN}(x-x_2)S^{em}_{\beta\dot\beta MQ}(x_2-x_1)\overline{S}^{\dot\beta\alpha nR}_{bJ}(x_1-y)\right]\right\}, \qquad (3.12)$$

$$C^{\dot\alpha\alpha aI}_{bJ}(x;y) = -g^2 f_{de}{}^f f_{lm}{}^n\int d^4x_1 d^4x_2 \left\{\Delta^{lK}_{fL}(x_2-x_1)\cdot \right.$$
$$\left.\cdot\left[\overline{S}^{\dot\alpha\beta aI}_{nK}(x-x_2)S^{em}_{\beta\dot\beta}(x_2-x_1)\overline{S}^{\dot\beta\alpha dL}_{bJ}(x_1-y)\right]\right\}. \qquad (3.13)$$

Taking the Fourier transform one obtains

$$\tilde A^{\dot\alpha\alpha aI}_{bJ}(p) = \frac{i}{2}g^2 N(N^2-1)\delta^a_b\delta^I_J\tilde{\overline{S}}^{\dot\alpha\beta}(p)\sigma^\lambda_{\beta\dot\beta}p_\lambda\tilde{\overline{S}}^{\dot\beta\alpha}(p)\left[\int\frac{d^4k}{(2\pi)^4}\frac{1}{\left(k+\frac{p}{2}\right)^2\left(k-\frac{p}{2}\right)^2}\right],$$

$$\tilde B^{\dot\alpha\alpha aI}_{bJ}(p) = -\tilde C^{\dot\alpha\alpha aI}_{bJ}(p) = \tilde A^{\dot\alpha\alpha aI}_{bJ}(p),$$

where

$$\tilde{\overline{S}}^{\dot\alpha\alpha}(p) = -i\frac{\overline{\sigma}^{\dot\alpha\alpha}_\mu p^\mu}{p^2}.$$

In conclusion the one-loop correction to the fermion propagator in the Wess–Zumino gauge turns out to be logarithmically divergent

$$\langle\left(\overline{\psi}\psi\right)^{\dot\alpha\alpha aI}_{bJ}\rangle^{1-\text{loop,WZ}}_{\text{FT}} = \tilde A^{\dot\alpha\alpha aI}_{bJ}(p) + \tilde B^{\dot\alpha\alpha aI}_{bJ}(p) + \tilde C^{\dot\alpha\alpha aI}_{bJ}(p) =$$
$$= \frac{i}{2}g^2 N(N^2-1)\delta^a_b\delta^I_J\tilde{\overline{S}}^{\dot\alpha\beta}(p)\sigma^\lambda_{\beta\dot\beta}p_\lambda\tilde{\overline{S}}^{\dot\beta\alpha}(p)\left[\int\frac{d^4k}{(2\pi)^4}\frac{1}{\left(k+\frac{p}{2}\right)^2\left(k-\frac{p}{2}\right)^2}\right]. \qquad (3.14)$$

Equation (3.14) shows that a logarithmically divergent wave function renormalization is required in the WZ gauge.

This divergence will now be proved to be a gauge artifact due to the choice of the WZ gauge. In fact if the fields C, χ and S are included two further contributions must be taken into account, which correspond to the diagrams



The contribution of these two diagrams exactly cancels the divergence in equation (3.14) giving a net vanishing one-loop result for the propagator. In fact the computation of D and E gives

$$D_{bJ}^{\dot{\alpha}\alpha aI}(x;y) = -\frac{1}{2}g^2 f_{de}{}^f f_{lm}{}^n \int d^4x_1 d^4x_2 \left\{ \Delta_{fL}^{lK}(x_2-x_1)\cdot \right.$$
$$\left. \cdot \left[\overline{S}_{mK}^{\dot{\alpha}\gamma aI}(x-x_2)\sigma_{\gamma\dot{\gamma}}^{\mu} \left(\partial_{\mu}^{(2)} R_{\dot{\beta}}^{\dot{\gamma}ne}(x_2-x_1) \right) \overline{S}_{bJ}^{\dot{\beta}\alpha dL}(x_1-y) \right] \right\} , \qquad (3.15)$$

$$E_{bJ}^{\dot{\alpha}\alpha aI}(x;y) = D_{bJ}^{\dot{\alpha}\alpha aI}(x;y)\,.$$

In momentum space one has

$$\tilde{D}_{bJ}^{\dot{\alpha}\alpha aI}(p) = -\frac{i}{4}g^2 N(N^2-1)\delta_b^a \delta_J^I \tilde{\overline{S}}^{\dot{\alpha}\beta}(p)\sigma_{\beta\dot{\beta}}^{\lambda} p_{\lambda} \tilde{\overline{S}}^{\dot{\beta}\alpha}(p) \left[\int \frac{d^4k}{(2\pi)^4} \frac{1}{\left(k+\frac{p}{2}\right)^2 \left(k-\frac{p}{2}\right)^2} \right] ,$$

$$\tilde{E}_{bJ}^{\dot{\alpha}\alpha aI}(p) = \tilde{D}_{bJ}^{\dot{\alpha}\alpha aI}(p)\,.$$

In conclusion summing up all the terms gives

$$\langle \left(\overline{\psi}\psi\right)_{bJ}^{\dot{\alpha}\alpha aI} \rangle_{\mathrm{FT}}^{1-\mathrm{loop}} =$$
$$= \tilde{A}_{bJ}^{\dot{\alpha}\alpha aI}(p) + \tilde{B}_{bJ}^{\dot{\alpha}\alpha aI}(p) + \tilde{C}_{bJ}^{\dot{\alpha}\alpha aI}(p) + \tilde{D}_{bJ}^{\dot{\alpha}\alpha aI}(p) + \tilde{E}_{bJ}^{\dot{\alpha}\alpha aI}(p) = 0\,, \qquad (3.16)$$

so that the one-loop correction to the $\langle\overline{\psi}\psi\rangle$ propagator, if the Wess–Zumino gauge is not exploited to perform the calculation, is zero, as expected in $\mathcal{N}$=4 super Yang–Mills theory.

Notice that the insertion of tadpoles such as



in a diagram gives a vanishing result because all the propagators are diagonal in colour space, so that the tadpole contains a factor $\delta_{ab}f^{abc} \equiv 0$. The same is true also for diagrams in $\mathcal{N}$=1 superspace that will be discussed in subsequent sections.

3.1.2 One loop corrections to the propagator of the scalars belonging to the chiral multiplet

The calculation of the propagator $\langle\varphi^{\dagger}\varphi\rangle$ is more complicated because more diagrams are involved. In particular tadpole type graphs, coming from the term $\Phi^{\dagger}V^2\Phi$ in the

action, are present. However the result is analogous to what was found for the $\langle\overline{\psi}\psi\rangle$ propagator: in the WZ gauge the one-loop correction is logarithmically divergent, requiring a wave function renormalization, but when the contributions neglected in the WZ gauge are considered the total one-loop result is vanishing.

The diagrams contributing to $\langle\varphi^\dagger\varphi\rangle$ at the one-loop level in the Wess–Zumino gauge are the following









The last three diagrams are all tadpole corrections since the free propagators for the auxiliary fields F and D are δ-functions, so that the corresponding lines shrink to a point. Each of these diagrams is quadratically divergent. There is also a quadratically divergent contribution coming from the two diagrams $B(x,y)$ and $C(x,y)$ and the sum of all these terms exactly vanishes. The sum of the previous diagrams gives a final result that is logarithmically divergent and in momentum space is schematically of the form

$$\langle(\varphi^\dagger\varphi)_{bJ}^{aI}\rangle_{\rm FT}^{1-\text{loop,WZ}} \sim g^2\delta_J^I\delta_b^a p^2 \int \frac{d^4k}{(2\pi)^4}\frac{1}{\left(k+\frac{p}{2}\right)^2\left(k-\frac{p}{2}\right)^2}\,. \tag{3.17}$$

Just like in the case of $\langle\overline{\psi}\psi\rangle$ this divergence corresponds to a wave function renormalization for the field $\varphi(x)$.

Taking into account the contribution of the fields C, χ and S there are eight more diagrams to be calculated



$$\longrightarrow \quad G_{bJ}^{aI}(x;y)$$



$$\longrightarrow \quad H_{bJ}^{aI}(x;y)$$



$$\longrightarrow \quad L_{bJ}^{aI}(x;y)$$



$$\longrightarrow \quad M_{bJ}^{aI}(x;y)$$



$$\longrightarrow \quad N_{bJ}^{aI}(x;y)$$



$$\longrightarrow \quad P_{bJ}^{aI}(x;y)$$



$$\longrightarrow \quad Q_{bJ}^{aI}(x;y)$$



Single diagrams contain quadratically divergent terms that cancel in the sum leaving a total contribution that is again logarithmically divergent. This non-vanishing correction is exactly what is needed to cancel the divergence in (3.17). As a result the complete one-loop correction to the $\langle\varphi^\dagger\varphi\rangle$ propagator is zero, whereas in the Wess–Zumino gauge a wave function renormalization is required.

Notice that the situation induced by the choice of the WZ gauge is a feature common to every supersymmetric gauge theory, since the contribution of the gauge-dependent fields C, χ and S is in general logarithmically divergent. However in theories with less supersymmetry a logarithmic wave function renormalization is usually necessary in any case, so that this effect is completely irrelevant. On the contrary it becomes important in the $\mathcal{N}$=4 Yang–Mills theory and in general in finite theories. Of course the divergences encountered here are gauge artifacts and disappear in gauge invariant correlation functions. Examples of computations of correlators of gauge invariant operators, in which the choice of the WZ gauge does not lead to this kind of problems, will be considered within the discussion of the correspondence with AdS type IIB superstring theory in the final chapter. Note also that the divergences encountered in the propagators are zero on-shell.

The calculations described here seem to suggest the possibility of constructing improved Feynman rules in which the effect of the gauge dependent fields, neglected in the WZ gauge, is dealt with by a suitable redefinition of the free propagators. However this program proves extremely complicated when the propagators of fields in the vector multiplet or three- and more-point functions are considered. The calculation of these correlation functions requires, already at the one-loop level, many other terms to be included in the action $S_{\rm int}$. In particular terms coming from $W^{(1)}W^{(2)}$ and $W^{(2)}W^{(2)}$ must be considered. The calculation of $W^{(2)}$ without the simplifications

introduced by the Wess–Zumino gauge is rather lengthy and gives

$$
\begin{aligned}
W_\alpha^{a(2)} \;=\; & -\frac{i}{2} f^a{}_{bc} \Big\{ -iC^b\lambda_\alpha^c - \frac{1}{2}\sigma^\mu_{\alpha\dot\alpha}\overline{\chi}^{\dot\alpha b}\partial_\mu C^c - \frac{i}{2}\sigma^\mu_{\alpha\dot\alpha}\overline{\chi}^{\dot\alpha b}A_\mu^c + \frac{1}{\sqrt{2}}S^{b\dagger}\chi_\alpha^c + \\
& + \Big[S^{b\dagger}S^c\delta_\alpha{}^\beta + \delta_\alpha{}^\beta C^b D^c - \frac{i}{2}(\sigma^\mu\overline{\sigma}^\nu)_\alpha{}^\beta C^b(\partial_\mu A_\nu^c - \partial_\nu A_\mu^c) + \\
& - \frac{i}{2}(\sigma^\mu\overline{\sigma}^\nu)_\alpha{}^\beta(A_\nu^b\partial_\mu C^c + A_\mu^b\partial_\nu C^c) - \frac{1}{2}\delta_\alpha{}^\beta C^b\Box C^c - \delta_\alpha{}^\beta\overline{\chi}^b\overline{\lambda}^c + \\
& + \frac{1}{2}(\sigma^\mu\sigma^\nu)_\alpha{}^\beta\partial_\mu C^b\partial_\nu C^c - \chi^{b\beta}\lambda_\alpha^c - \lambda^{b\beta}\chi_\alpha^c - i\sigma^\mu_{\alpha\dot\alpha}\overline{\chi}^{b\dot\alpha}\partial_\mu\chi^{c\beta} + \\
& + i\varepsilon^{\gamma\beta}\sigma^\mu_{\gamma\dot\beta}\partial_\mu\overline{\chi}^b\chi_\alpha^c + \frac{1}{2}(\sigma^\mu\sigma^\nu)_\alpha{}^\beta A_\nu^b A_\mu^c \Big]\theta_\beta + \Big[C^b\sigma^\mu_{\alpha\dot\alpha}\partial_\mu\overline{\lambda}^{c\dot\alpha} + \\
& - i\chi_\alpha^b D^c - \frac{1}{4}(\sigma^\mu\overline{\sigma}^\nu)_\alpha{}^\beta\chi_\beta^b(\partial_\mu A_\nu^c - \partial_\nu A_\mu^c) - \frac{1}{2}A_\nu^b(\sigma^\mu\overline{\sigma}^\nu)_\alpha{}^\beta\partial_\mu\chi_\beta^c + \\
& + \frac{1}{2}\partial_\mu A^{b\mu}\chi_\alpha^c - \frac{i}{2}\partial_\nu C^b(\sigma^\mu\overline{\sigma}^\nu)_\alpha{}^\beta\partial_\mu\chi_\beta^c - \frac{i}{\sqrt{2}}\sigma^\mu_{\alpha\dot\alpha}\partial_\mu\overline{\chi}^{\dot\alpha b}S^c + \sqrt{2}S^b\lambda_\alpha^c + \\
& - iA_\mu^b\sigma^\mu_{\alpha\dot\alpha}\overline{\lambda}^{c\dot\alpha}\Big]\theta\theta \Big\} \qquad (3.18)
\end{aligned}
$$

Substituting (3.18) into $S_{\rm int}$ results in a number of relevant interaction terms of the order of 100, making this formulation totally impractical in explicit calculations. In conclusion perturbation theory in components requires almost unavoidably the WZ gauge, but the latter introduces divergences in gauge dependent quantities. In the following sections the superfield formalism, that allows to avoid these problems, will be employed. However as will be shown new difficulties related to the gauge fixing emerge.

3.2 Perturbation theory in $\mathcal{N}$=1 superspace: propagators

The difficulties encountered in the previous section in the calculation of gauge dependent quantities, because of the divergences introduced by the Wess–Zumino gauge, can be overcome using the $\mathcal{N}$=1 superfield formulation of the model.

In chapter 2 it has already been pointed out how the $\mathcal{N}$=1 superfield formalism has proved to be a powerful tool in the proof of the finiteness of the theory up to three loops. In this approach there is no particular difficulty in working without fixing the WZ gauge. If one does not choose to work in the WZ gauge the action is non polynomial, however a finite number of terms is relevant at each order in perturbation

theory, so that only at very high order the Wess–Zumino gauge introduces significant simplifications. The aim of this section is to show that there are other subtleties related to the gauge fixing, even if one does not work in the WZ gauge.

The formulation of the theory in $\mathcal{N}$=1 superspace that will be employed is a 'mass deformed' version of that given by equation (2.10). In other words the action that will be considered is that of equation (2.10) plus mass terms for the (anti) chiral superfields

$$\begin{aligned} S_{\rm m} &= -\int d^4x d^4\theta \left[\frac{1}{2} m \delta_{IJ} \Phi^I_a(x,\theta,\overline{\theta}) \Phi^{Ja}(x,\theta,\overline{\theta}) \delta(\overline{\theta}) + \right. \\ &\left. + \frac{1}{2} m^* \delta^{IJ} \Phi^{\dagger a}_{\ I}(x,\theta,\overline{\theta}) \Phi^{\dagger}_{\ Ja}(x,\theta,\overline{\theta}) \delta(\theta) \right] . \end{aligned} \tag{3.19}$$

The addition of this term to the action breaks supersymmetry down to $\mathcal{N}$=1. In [76] it was argued, by means of dimensional arguments, that this term should not affect the ultraviolet properties of the $\mathcal{N}$=4 theory. More precisely the statement of [76] is that no divergences appear in gauge invariant quantities, so that no divergent contribution to the quantum effective action is generated perturbatively. As a result the model obtained in this way would be an example of a finite $\mathcal{N}$=1 theory. The inverse construction, in which one deforms a $\mathcal{N}$=1 model to $\mathcal{N}$=4 super Yang–Mills plus a mass term, has been proposed in [77, 78] as a regularization procedure preserving supersymmetry. The calculations presented in this section will show that in the presence of the term (3.19) no divergence is generated, at the one-loop level, in the two-, three- and four-point Green functions that are computed. The results presented suggest the possibility of reinforcing the conclusions of [76]; namely the $\mathcal{N}$=4 theory augmented with (3.19) appears to be finite in the sense that no divergences appear in the total n-point irreducible Green functions, at least at one loop. In particular no wave function renormalization is required. Notice that this result is what is actually necessary for the consistency of the approach advocated in [77, 78].

The complete action is [2]

$$\begin{aligned} S = \int d^4x\, d^4\theta \, \Big\{ & V^a(x,\theta,\overline{\theta}) \left[-\Box P_T - \xi (P_1 + P_2) \Box \right] V_a(x,\theta,\overline{\theta}) + \\ & + \Phi^{\dagger a}_{\ I}(x,\theta,\overline{\theta}) \Phi^I_a(x,\theta,\overline{\theta}) - \frac{1}{2} m \delta_{IJ} \Phi^I_a(x,\theta,\overline{\theta}) \Phi^{Ja}(x,\theta,\overline{\theta}) \delta(\overline{\theta}) + \\ & - \frac{1}{2} m \delta^{IJ} \Phi^{\dagger a}_{\ I}(x,\theta,\overline{\theta}) \Phi^{\dagger}_{\ Ja}(x,\theta,\overline{\theta}) \delta(\theta) + i g f_{abc} \Phi^{\dagger a}_{\ I}(x,\theta,\overline{\theta}) V^b(x,\theta,\overline{\theta}) \cdot \end{aligned} \tag{3.20}$$

[2] In the following the mass parameter m will be taken to be real for simplicity of notation.

$$
\begin{aligned}
&\cdot\Phi^{Ic}(x,\theta,\overline{\theta}) - \frac{1}{2}g^2 f_{ab}{}^e f_{ecd}\Phi^{\dagger a}_I(x,\theta,\overline{\theta})V^b(x,\theta,\overline{\theta})V^c(x,\theta,\overline{\theta})\Phi^{Id}(x,\theta,\overline{\theta}) + \\
&-\frac{i}{4}g f_{abc}\left[\overline{D}^2\left(D^\alpha V^a(x,\theta,\overline{\theta})\right)\right]V^b(x,\theta,\overline{\theta})\left(D_\alpha V^c(x,\theta,\overline{\theta})\right) + \\
&-\frac{1}{8}g^2 f_{ab}{}^e f_{ecd}V^a(x,\theta,\overline{\theta})\left(D^\alpha V^b(x,\theta,\overline{\theta})\right)\left[\left(\overline{D}^2 V^c(x,\theta,\overline{\theta})\right)\left(D_\alpha V^d(x,\theta,\overline{\theta})\right)\right] + \\
&+\ldots - \frac{\sqrt{2}}{3!}g f^{abc}\left[\varepsilon_{IJK}\Phi^I_a(x,\theta,\overline{\theta})\Phi^J_b(x,\theta,\overline{\theta})\Phi^K_c(x,\theta,\overline{\theta})\delta(\overline{\theta}) + \right. \\
&\left. + \varepsilon^{IJK}\Phi^\dagger{}_{Ia}(x,\theta,\overline{\theta})\Phi^\dagger{}_{Jb}(x,\theta,\overline{\theta})\Phi^\dagger{}_{Kc}(x,\theta,\overline{\theta})\delta(\theta)\right] + \left(\overline{C}'{}_a(x,\theta,\overline{\theta})C^a(x,\theta,\overline{\theta}) + \right. \\
&\left. - C'{}_a(x,\theta,\overline{\theta})\overline{C}^a(x,\theta,\overline{\theta})\right) + \frac{i}{2\sqrt{2}}g f_{abc}\left(C'^a(x,\theta,\overline{\theta}) + \overline{C}'^a(x,\theta,\overline{\theta})\right)\cdot \\
&\cdot V^b(x,\theta,\overline{\theta})\left(C^c(x,\theta,\overline{\theta}) + \overline{C}^c(x,\theta,\overline{\theta})\right) - \frac{1}{8}g^2 f_{ab}{}^e f_{ecd}\left(C'^a(x,\theta,\overline{\theta}) + \overline{C}'^a(x,\theta,\overline{\theta})\right)\cdot \\
&\cdot V^b(x,\theta,\overline{\theta})V^c(x,\theta,\overline{\theta})\left(C^d(x,\theta,\overline{\theta}) + \overline{C}^d(x,\theta,\overline{\theta})\right) + \ldots\Big\}\,, \qquad (3.21)
\end{aligned}
$$

where the dots stand for terms that are not relevant for the considerations of this chapter.

Notice that in the action (3.21) a gauge fixing term corresponding to a family of gauges parameterized by $\alpha = \frac{1}{\xi}$ has been introduced. It will now be shown, by explicitly computing the propagators of both the chiral and the vector superfields, that the supersymmetric generalization of the Fermi–Feynman gauge, corresponding to $\alpha = 1$, is somehow privileged (see also [75, 80]), since any other choice of the parameter α leads to infrared divergences in Green functions.

3.2.1 Propagator of the chiral superfield

The propagator of the chiral superfield is the simplest Green function to compute. The calculation will be reported in detail in order to illustrate the superfield technique.

It follows from the action (2.10) that there are three diagrams contributing to the propagator $\langle\Phi\Phi^\dagger\rangle$ at the one loop level. The one-particle irreducible parts of these diagrams will be directly evaluated in momentum space using the super Feynman rules of section 1.6. The convention employed is that all momenta are taken to be incoming. The diagrams are the following



where $z=(x,\theta,\overline{\theta})$ and the notation for the internal propagators is



$$\longrightarrow \quad \langle \Phi^{aI}(z)\Phi^{b\dagger}_J(z')\rangle_{\rm free} = \delta^I_J\delta^a_b\frac{1}{\Box+m^2}\delta_8(z-z')$$



$$\longrightarrow \quad \langle V^a(z)V_b(z')\rangle_{\rm free} = -\frac{\delta^a_b}{\Box}[1+(\alpha-1)(P_1+P_2)]\delta_8(z-z')$$

P_1 and P_2 in the $\langle VV\rangle$ propagator are the projectors (1.63). Moreover in the presence of mass terms for Φ and $\Phi^\dagger$ there are free propagators $\langle\Phi\Phi\rangle$ and $\langle\Phi^\dagger\Phi^\dagger\rangle$, that will enter the calculation of the vector superfield propagator at one loop



$$\longrightarrow \quad \langle \Phi^{aI}(z)\Phi^J_b(z')\rangle_{\rm free} = -\delta^{IJ}\delta^a_b\,\frac{m}{4}\frac{D^2}{\Box(\Box+m^2)}\delta_8(z-z')$$



$$\longrightarrow \quad \langle \Phi^{a\dagger}_I(z)\Phi^\dagger_{bJ}(z')\rangle_{\rm free} = -\delta_{IJ}\delta^a_b\,\frac{m}{4}\frac{\overline{D}^2}{\Box(\Box+m^2)}\delta_8(z-z')$$

Using these free propagators and the rules of section 1.6, with the vertices read

from the action (3.21), the three contributions can be evaluated as follows

$$\tilde{A}(p) = \int \frac{d^4k}{(2\pi)^4} d^4\theta_1 d^4\theta_2 \left\{ igf_{acd}\Phi_I^{a\dagger}(p,\theta_1,\overline{\theta}_1)\left[\left(-\frac{1}{4}\overline{D}_1^2\right)\frac{\delta_J^I\delta^{cf}\delta(1,2)}{(p-k)^2+m^2}\cdot\right.\right.$$
$$\left.\left.\cdot\left(-\frac{1}{4}\overleftarrow{D}_1^2\right)\right]\left[\left(1+\gamma(D_1^2\overline{D}_1^2+\overline{D}_1^2D_1^2)\right)\left(-\frac{\delta^{de}\delta(1,2)}{k^2}\right)\right] igf_{efb}\Phi^{bJ}(-p,\theta_2,\overline{\theta}_2)\right\},$$

where $\gamma=(\alpha-1)$ and the compact notation $\delta(1,2)=\delta_2(\theta_1-\theta_2)\delta_2(\overline{\theta}_1-\overline{\theta}_2)$ has been introduced. The computation uses the properties of the δ-function, which imply for example [23, 4]

$$D_{1\alpha}\delta(1,2) = -\delta(1,2)\overleftarrow{D}_{2\alpha}\,, \qquad \overline{D}_{1\dot{\alpha}}\delta(1,2) = -\delta(1,2)\overleftarrow{\overline{D}}_{2\dot{\alpha}}\,,$$
$$\overline{D}_{1\dot{\alpha}}D_{1\alpha}\delta(1,2) = \delta(1,2)\overleftarrow{\overline{D}}_{2\dot{\alpha}}\overleftarrow{D}_{2\alpha}\,, \qquad \overline{D}_1^2D_1^2\delta(1,2) = \delta(1,2)\overleftarrow{\overline{D}}_2^2\overleftarrow{D}_2^2\,,$$

and integrations by parts on the Grassmannian variables in order to remove the D and $\overline{D}$ derivatives from one δ, so that one θ integration can be performed immediately. In this way one obtains an expression that is local in θ as is expected from the $\mathcal{N}$=1 non-renormalization theorem. For $\tilde{A}(p)$ one obtains

$$\tilde{A}(p) = -g^2N(N^2-1)\delta_b^a\delta_J^I\left(\frac{1}{4}\right)^2\int\frac{d^4k}{(2\pi)^4}d^4\theta_1d^4\theta_2\,\frac{1}{k^2[(p-k)^2+m^2]}\left\{\Phi_{aI}^\dagger(p,1)\cdot\right.$$
$$\left.\cdot\Phi^{bJ}(-p,2)\left[\overline{D}_1^2D_1^2\delta(1,2)\right]\left[1+\gamma(D_1^2\overline{D}_1^2+\overline{D}_1^2D_1^2)\delta(1,2)\right]\right\} = \tilde{A}_1(p)+\tilde{A}_2(p)\,.$$

The first term is trivially calculated using [23, 4]

$$\int d^4\theta\,\left[\overline{D}^2D^2\delta(\theta-\theta')\right]\delta(\theta-\theta') = 16$$
$$\int d^4\theta\,\left[\overline{D}^mD^n\delta(\theta-\theta')\right]\delta(\theta-\theta') = 0 \quad \text{if} \quad (m,n)\neq(2,2) \tag{3.22}$$

and gives

$$\tilde{A}_1(p) = -g^2N(N^2-1)\delta_b^a\delta_J^I\int\frac{d^4k}{(2\pi)^4}d^4\theta\,\frac{1}{k^2[(p-k)^2+m^2]}\left\{\Phi_{aI}^\dagger(p,\theta,\overline{\theta})\Phi^{bJ}(-p,\theta,\overline{\theta})\right\}. \tag{3.23}$$

In the computation of the second term one must use the (anti) commutators of covariant derivatives, see appendix A, which in particular imply [23, 4]

$$\overline{D}^2D^2\overline{D}^2 = 16\Box\overline{D}^2 \qquad D^2\overline{D}^2D^2 = 16\Box D^2\,. \tag{3.24}$$

Then integration by parts analogously gives

$$\tilde{A}_2(p) = \gamma\, g^2 N(N^2-1)\delta_b^a\delta_J^I \int \frac{d^4k}{(2\pi)^4} d^4\theta\, \frac{[(p-k)^2+p^2]}{k^4[(p-k)^2+m^2]} \left\{\Phi^\dagger_{aI}(p,\theta,\overline{\theta})\Phi^{bJ}(-p,\theta,\overline{\theta})\right\}. \tag{3.25}$$

In conclusion from the first diagram one obtains two contributions, the first proportional to γ and the second independent of it. The second diagram gives one single γ-independent contribution. The Feynman rules give

$$\begin{aligned}\tilde{B}(p) = \int \frac{d^4k}{(2\pi)^4} d^4\theta_1 d^4\theta_2 \Bigg\{ & \left(-\frac{\sqrt{2}}{3!}\right) g\varepsilon^I{}_{KL} f_{acd} \Phi^{\dagger a}_I(p,1)\cdot \\ & \cdot \left[\left(-\frac{1}{4}D_1{}^2\right)\frac{\delta^{ce}\delta^K_M\delta(1,2)}{(p-k)^2+m^2}\left(-\frac{1}{4}\overleftarrow{\overline{D}}_2{}^2\right)\right]\left[\frac{\delta^d_f\delta^L_N\delta(1,2)}{k^2+m^2}\right]\cdot \\ & \cdot \left(-\frac{\sqrt{2}}{3!}\right)\varepsilon_J{}^{MN} f_{be}{}^f \Phi^{bJ}(-p,2)\Bigg\}\,.\end{aligned}$$

Proceeding exactly as for $\tilde{A}_1$ one obtains

$$\tilde{B}(p) = g^2 N(N^2-1)\delta_b^a\delta_J^I \int \frac{d^4k}{(2\pi)^4}\, d^4\theta\, \left\{\frac{\Phi^\dagger_{aI}(p,\theta,\overline{\theta})\Phi^{bJ}(-p,\theta,\overline{\theta})}{(k^2+m^2)[(p-k)^2+m^2]}\right\}. \tag{3.26}$$

From the last diagram one gets

$$\begin{aligned}\tilde{C}(p) = \int \frac{d^4k}{(2\pi)^4} d^4\theta_1\, \Bigg\{ & \left(-\frac{g^2}{2}\right) f_{ac}{}^d f_{deb} \Phi^{a\dagger}_I(p,1)\cdot \\ & \cdot \left[-\left(1+\gamma(\overline{D}_1^2 D_1^2 + D_1^2\overline{D}_1^2)\right)\frac{\delta^{ce}\delta(1,1)}{k^2}\right]\Phi^{bJ}(-p,1)\Bigg\}\,.\end{aligned}$$

The only non-vanishing contribution comes from the term in which the projection operators act on the δ-function, since $\delta_4(\theta-\theta)=0$, so that

$$\tilde{C}(p) = -\gamma\, g^2 N(N^2-1)\delta^b_a\delta^I_J \int \frac{d^4k}{(2\pi)^4} d^4\theta \frac{1}{k^4}\left\{\Phi^{a\dagger}_I(p,\theta,\overline{\theta})\Phi^J_b(-p,\theta,\overline{\theta})\right\}, \tag{3.27}$$

from which one sees that the C contribution is absent in the gauge $\alpha=1$, *i.e.* $\gamma=0$.

Putting the various corrections together gives the following result. The sum of the terms $\tilde{A}_1(p)$ and $\tilde{B}(p)$, which does not depend on γ, is

$$\begin{aligned}\tilde{A}_1(p)+\tilde{B}(p) \;=\; & g^2 N(N^2-1)\delta^b_a\delta^I_J \int \frac{d^4k}{(2\pi)^4} d^4\theta \left[\Phi^{a\dagger}_I(p,\theta,\overline{\theta})\Phi^J_b(-p,\theta,\overline{\theta})\right]\cdot \\ & \cdot\left\{\frac{1}{k^2\left[(p-k)^2+m^2\right]} - \frac{1}{(k^2+m^2)\left[(p-k)^2+m^2\right]}\right\}.\end{aligned}$$

This is finite and exactly vanishes for $m = 0$, *i.e.* in the limit in which the $\mathcal{N}$=4 theory is recovered.

The sum of the terms $\tilde{A}_2(p)$ and $\tilde{C}(p)$ is proportional to γ and reads

$$\begin{aligned}\tilde{A}_2(p) + \tilde{C}(p) &= \gamma\, g^2 N(N^2-1)\delta_a^b\delta_J^I \int \frac{d^4k}{(2\pi)^4} d^4\theta \left[\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Phi_b^J(-p,\theta,\overline{\theta})\right] \cdot \\ &\cdot \left\{ \left(\frac{(p-k)^2}{k^4\left[(p-k)^2+m^2\right]} - \frac{1}{k^4} \right) + \left(\frac{p^2}{k^4\left[(p-k)^2+m^2\right]} \right) \right\} .\end{aligned}$$

Both of the terms in the last integral are infrared divergent. The first one is zero in the limit $m \to 0$, while the second one gives an infrared divergence that is still present in the $\mathcal{N}$=4 theory, *i.e.* with $m = 0$. More precisely putting

$$\tilde{A}_2(p) + \tilde{C}(p) = \gamma\, g^2 N(N^2-1)\delta_a^b\delta_J^I \left[\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Phi_b^J(-p,\theta,\overline{\theta})\right] \left[I_1(p) + I_2(p)\right] ,$$

one has

$$\begin{aligned} I_1(p) &= \int \frac{d^4k}{(2\pi)^4} \left\{ \frac{(p-k)^2}{k^4\left[(p-k)^2+m^2\right]} - \frac{1}{k^4} \right\} = \\ &= \int \frac{d^4k}{(2\pi)^4} \left\{ \frac{m^2}{k^4\left[(p-k)^2+m^2\right]} \right\} = \\ &= 2m^2 \int_0^1 d\zeta\, \zeta \int \frac{d^4k}{(2\pi)^4} \frac{1}{\left[k^2 + (p^2\zeta + m^2)(1-\zeta)\right]^3} = \\ &= \frac{2m^2}{2(4\pi)^2} \int_0^1 d\zeta\, \zeta \frac{1}{(p^2\zeta + m^2)(1-\zeta)} = \\ &= -\frac{m^2}{(4\pi)^2} \left[\frac{1}{(p^2+m^2)} \log\epsilon \;+\; \frac{m^2}{p^2(p^2+m^2)} \log\left(\frac{p^2+m^2}{m^2} \right) \right] \end{aligned}$$

where a standard Feynman parameterization has been used. Analogously

$$\begin{aligned} I_2(p) &= \int \frac{d^4k}{(2\pi)^4} \left\{ \frac{p^2}{k^4\left[(p-k)^2+m^2\right]} \right\} = \\ &= 2p^2 \int_0^1 d\zeta\, \zeta \int \frac{d^4k}{(2\pi)^4} \frac{1}{\left[k^2 + (p^2\zeta + m^2)(1-\zeta)\right]^3} = \\ &= \frac{p^2}{(4\pi)^2} \int_0^1 d\zeta\, \zeta \frac{1}{(p^2\zeta + m^2)(1-\zeta)} = \\ &= -\frac{1}{(4\pi)^2} \left[\frac{p^2}{(p^2+m^2)} \log\epsilon \;+\; \frac{m^2}{(p^2+m^2)} \log\left(\frac{p^2+m^2}{m^2} \right) \right] . \end{aligned}$$

In the above expressions an infrared regulator ϵ has been introduced.

Notice that the total correction is exactly zero on-shell, *i.e.* for p^2=m^2.

To summarize the results, the propagator of the chiral superfields of the $\mathcal{N}$=4 super Yang–Mills theory in $\mathcal{N}$=1 superspace is logarithmically infrared-divergent for any choice of the gauge parameter $\alpha \neq 1$. This divergence corresponds to a wave function renormalization for the superfields Φ^I and vanishes on-shell. In the Fermi–Feynman gauge α=1 the one-loop correction exactly vanishes.

3.2.2 Propagator of the vector superfield

The one-loop calculation of the propagator of the vector superfield is much more complicated, because many more diagrams are involved producing a large number of contributions. The final result is however completely analogous: in the Fermi–Feynman gauge the one-loop correction is zero, whereas infrared divergences arise for $\alpha \neq 1$. In the presence of a mass term for the (anti) chiral superfields no new divergences are generated.

Form a calculational viewpoint the new feature with respect to the $\langle \Phi \Phi^\dagger \rangle$ case is that there are also diagrams involving the ghosts. There are two multiplets of ghosts, described by the chiral superfields C and C'. The free propagators for these superfields will be denoted by



$$\longrightarrow \quad \langle \overline{C}^{a\prime}(z) C^b(z') \rangle_{\text{free}} = \delta^a_b \frac{1}{\Box} \delta_8(z - z')$$

$$\longrightarrow \quad \langle \overline{C}^{a}(z) C^{b\prime}(z') \rangle_{\text{free}} = \delta^a_b \frac{1}{\Box} \delta_8(z - z')$$

The ghosts are treated exactly like ordinary chiral superfields with the only difference that there is a minus sign associated with the loops, because C and C' are anticommuting fields [23].

The corrections to the $\langle VV \rangle$ propagator at the one-loop level correspond to the diagrams



$\longrightarrow \quad A(z;z')$



$\longrightarrow \quad B(z;z')$



$\longrightarrow \quad C(z;z')$



$\longrightarrow \quad D_1(z;z')$



$\longrightarrow \quad D_2(z;z')$



$\longrightarrow \quad D_3(z;z')$



The contributions A to E are rather straightforward to evaluate much in the same way as the diagrams entering the $\langle \Phi \Phi^\dagger \rangle$ propagator. The last two graphs are more involved because the free propagator for the V superfield is more complicated for generic values of the parameter α. Moreover the cubic and quartic vertices

$$
\begin{aligned}
&-\frac{i}{16\sqrt{2}} g f_{abc} \left[\overline{D}^2 \left(D^\alpha V^a\right)\right] V^b \left(D_\alpha V^c\right) \\
&-\frac{1}{128} g^2 f_{ab}{}^e f_{ecd} V^a \left(D^\alpha V^b\right) \left[\left(\overline{D}^2 V^c\right) \left(D_\alpha V^d\right)\right] ,
\end{aligned}
$$

lead to various terms corresponding to the ways the covariant derivatives can act on

the V lines. The V^3 vertex in particular



gives rise to six different terms.

Schematically the calculation goes as follows. $\tilde{A}(p)$ is a contribution that appears because of the addition of the mass terms (3.19); it is not present in the $\mathcal{N}$=4 theory, *i.e.* when m=0. It is useful to discuss separately the corrections coming from diagrams $\tilde{A}$, $\tilde{B}$ and $\tilde{C}$ and those obtained from $\tilde{D}_i$, $\tilde{E}_i$, $\tilde{F}$ and $\tilde{G}$, since the latter correspond to the one-loop contribution to the vector superfield propagator in the $\mathcal{N}$=1 supersymmetric Yang–Mills theory.

The first diagram, $\tilde{A}$, is logarithmically divergent and reads

$$\tilde{A}(p) = 3g^2N(N^2-1)\delta_{ab}\int \frac{d^4k}{(2\pi)^4}d^4\theta \left\{ \frac{m^2\left[V^a(p,\theta,\overline{\theta})V^b(-p,\theta,\overline{\theta})\right]}{(k^2+m^2)[(p-k)^2+m^2]} \right\}. \qquad (3.28)$$

For the diagram $\tilde{B}$ application of the Feynman rules gives rise to three different contributions, one quadratically divergent and two logarithmically divergent

$$\begin{aligned}\tilde{B}(p) &= 3g^2N(N^2-1)\delta_{ab}\int \frac{d^4k}{(2\pi)^4}d^4\theta\, \frac{1}{(k^2+m^2)[(p-k)^2+m^2]}\cdot \\ &\cdot\left\{ k^2V^a(p,\theta,\overline{\theta})V^b(-p,\theta,\overline{\theta}) - \frac{i}{4}p_\mu\sigma^\mu_{\alpha\dot{\alpha}}V^a(p,\theta,\overline{\theta})\left[(\overline{D}^{\dot{\alpha}}D^\alpha)V^b(-p,\theta,\overline{\theta})\right] + \right. \\ &\left. +\frac{1}{4}V^a(p,\theta,\overline{\theta})\left[(\overline{D}^2D^2)V^b(-p,\theta,\overline{\theta})\right]\right\}. \qquad (3.29)\end{aligned}$$

The tadpole diagram $\tilde{C}$ gives a quadratically divergent result of the form

$$\tilde{C} = -3g^2N(N^2-1)\delta^{ab}\int \frac{d^4k}{(2\pi)^4}d^4\theta\, \frac{1}{(k^2+m^2)}\left\{V^a(p,\theta,\overline{\theta})V^b(-p,\theta,\overline{\theta})\right\}. \qquad (3.30)$$

Summing and subtracting m^2 in the numerator of the first term in (3.29) and putting the three corrections $\tilde{A}$, $\tilde{B}$ and $\tilde{C}$ together gives a net result that is only logarithmi-

cally divergent

$$\begin{aligned}\tilde{A}(p)+\tilde{B}(p)+\tilde{C}(p) &= -3g^2N(N^2-1)\delta_{ab}\int\frac{d^4k}{(2\pi)^4}d^4\theta\,\frac{1}{(k^2+m^2)[(p-k)^2+m^2]}\cdot\\ &\cdot\left\{\frac{i}{4}p_\mu\sigma^\mu_{\alpha\dot\alpha}V^a(p,\theta,\overline{\theta})\left[(\overline{D}^{\dot\alpha}D^\alpha)V^b(-p,\theta,\overline{\theta})\right]+\right.\\ &\left.+\frac{1}{16}V^a(p,\theta,\overline{\theta})\left[(\overline{D}^2D^2)V^b(-p,\theta,\overline{\theta})\right]\right\}. \qquad (3.31)\end{aligned}$$

Notice that in particular the logarithmically divergent contribution proportional to m^2 exactly cancels out. This is crucial because this correction would correspond to a mass renormalization for the vector superfield that is known to be excluded in any gauge theory as well as in supersymmetric theories in general.

The diagrams $\tilde{D}_i$ and $\tilde{E}_i$ are completely analogous to the previous ones, with the only difference that the mass does not appear in the denominators and there is a minus sign associated with the loops. Their sum is logarithmically divergent and takes the form

$$\begin{aligned}&\tilde{D}_1(p)+\tilde{D}_2(p)+\tilde{D}_3(p)+\tilde{E}_1(p)+\tilde{E}_2(p)=\frac{1}{8}g^2N(N^2-1)\delta_{ab}\cdot\\ &\cdot\int\frac{d^4k}{(2\pi)^4}d^4\theta\,\frac{1}{k^2(p-k)^2}\left\{ip_\mu\sigma^\mu_{\alpha\dot\alpha}V^a(p,\theta,\overline{\theta})\left[(\overline{D}^{\dot\alpha}D^\alpha)V^b(-p,\theta,\overline{\theta})\right]+\right.\\ &\left.+\frac{1}{8}V^a(p,\theta,\overline{\theta})\left[(\overline{D}^2D^2)V^b(-p,\theta,\overline{\theta})\right]\right\}. \qquad (3.32)\end{aligned}$$

All of these corrections do not depend on the gauge parameter $\gamma{=}\alpha-1$ and must be summed to those coming from the last two diagrams. $\tilde{F}$ exactly vanishes for any α, so that only $\tilde{E}$ needs to be considered. This diagram produces in principle 72 corrections because the Feynman rules give rise to 18 terms (distributing the covariant derivatives associated with the two vertices), each of which splits into 4, since the free propagator itself contains two terms. Many of these contributions can be easily shown to vanish using the properties of the covariant derivatives. In particular one uses

$$\overline{D}^2D_\alpha\overline{D}^2D^2=0\,,$$

which follows from $\overline{D}^3=0$ and use of the (anti) commutation relations for the D's. It is useful to separate in the non-vanishing part terms proportional to γ and γ^2, from the γ-independent terms. The latter combine to give a logarithmically divergent correction that together with that of equation (3.32) cancel the correction

(3.31) coming from the sum $\tilde{A}+\tilde{B}+\tilde{C}$. Actually if $m\neq 0$ this sum is finite and exactly vanishes at m=0. As a result the only non-vanishing corrections to the vector superfield propagator at the one loop level come from terms proportional to γ and to γ^2 in $\tilde{E}(p)$. The former contain an infrared divergent part of the form

$$J^{(1)}(p)=c_1\gamma g^2\delta^a_b\int\frac{d^4k}{(2\pi)^4}d^4\theta\,\frac{\sigma^\mu_{\alpha\dot\alpha}\sigma^\nu_{\beta\dot\beta}p_\mu p_\nu}{k^4(p-k)^2}\left\{V_a(p,\theta,\overline\theta)\left[D^\alpha\overline{D}^{\dot\alpha}\overline{D}^{\dot\beta}D^\beta V^b(-p,\theta,\overline\theta)\right]\right\}\,,$$

but is ultraviolet finite. Furthermore there is a correction, finite both in the ultraviolet and in the infrared regions, proportional to γ^2, that reads

$$\begin{aligned}J^{(2)}(p) &= c_2\gamma^2 g^2\delta^a_b\int\frac{d^4k}{(2\pi)^4}d^4\theta\,\frac{(\sigma^\nu\overline\sigma^\mu\sigma^\lambda)_{\alpha\dot\alpha}(p-k)_\mu k_\nu p_\lambda}{k^4(p-k)^4}\,\cdot\\ &\quad\cdot\left\{\left[(\overline{D}^2D^\alpha)V_a(p,\theta,\overline\theta)\right]\left[(D^2\overline{D}^{\dot\alpha})V^b(-p,\theta,\overline\theta)\right]\right\}\,.\end{aligned}$$

In the previous expressions for $J^{(1)}$ and $J^{(2)}$ c_1 and c_2 are non-vanishing numerical constants proportional to $N(N^2-1)$.

In conclusion in the $\mathcal{N}$=4 theory, *i.e.* when m=0, the one-loop correction to the vector superfield propagator, just like that of the chiral superfield, is infrared singular unless the Fermi–Feynman gauge, α=1, is chosen, in which case it vanishes. Like in the case of the chiral superfield propagator the non-vanishing γ-dependent corrections correspond to a wave function renormalization for the superfield V and are zero on-shell, *i.e.* for p^2=0.

The proof of the finiteness of the theory in the presence of the mass terms (3.19) given in [76] is based on naive power counting, which gives the superficial degree of divergence, d_s, of a diagram in $\mathcal{N}$=1 superspace [24]

$$d_s=2-E-C\,,$$

where E is the number of external (anti) chiral lines and C the number of $\Phi\Phi$ or $\Phi^\dagger\Phi^\dagger$ propagators. In [76] it is also argued that for corrections to the effective action involving only V superfields the requirement of gauge invariance reduces the degree of divergence to

$$d_s=-C\,.$$

However for the purposes of [77, 78] it appears crucial that no divergences, not even corresponding to a wave function renormalization, are present in total n-point functions for any n. The computation of the two-point functions in this section has shown

that this actually the case at one loop. An argument for the generalization of this result to Green functions with an arbitrary number of external V lines will now be briefly discussed. First note that diagrams involving only vector and ghost superfields are not modified by the inclusion of (3.19), so that one must only consider graphs containing internal chiral lines. The ultraviolet properties of diagrams that are only logarithmically divergent in the original $\mathcal{N}$=4 theory are not modified by the presence of the mass in the propagators. Thus the contributions to be analyzed are the quadratically divergent ones, that can acquire subleading logarithmic singularities, plus eventually new diagrams involving $\Phi\Phi$ and $\Phi^\dagger\Phi^\dagger$ propagators. The relevant quadratic divergences come from tadpole diagrams



The only new diagram containing $\Phi\Phi$ and $\Phi^\dagger\Phi^\dagger$ propagators that must be considered is



Since all the vertices entering these graphs come from the expansion of the term $\mathrm{tr}\left[e^{-V}\Phi^\dagger e^{V}\Phi\right]$ in the action, the only covariant derivatives that are present come from the functional derivatives and must act on the internal (anti) chiral lines to give a non-vanishing result. As a consequence the loop integral in the above diagrams is exactly the same as for the $\tilde{C}$ and $\tilde{A}$ corrections to the VV propagator. Hence

summing to the previous diagrams the contribution of



leads to a net logarithmically divergent correction that is exactly the same as the one obtained in the original $\mathcal{N}$=4 theory. Like in the latter case this divergence will be cancelled by the contributions coming from the other one-loop diagrams involving V and ghost internal lines. The argument given here reinforces the results of [76], at least at the one-loop level, and supports the proposal, put forward in [77, 78], according to which the mass deformed $\mathcal{N}$=4 model can be considered a consistent supersymmetry-preserving regularization for a class of $\mathcal{N}$=1 theories.

Notice that for the previous discussion it is not necessary to consider equal masses for all the (anti) chiral superfields. The same results can be proved giving different masses to the three superfields. This can be easily understood since in each diagram considered in this section only one chiral/anti-chiral pair is involved, because the propagators are diagonal in 'flavour' space and the vertices containing vector and (anti) chiral superfields couple $\Phi_I^\dagger$ and Φ^I with the same index I. Basically this means that the above discussed cancellations apply separately to the contributions of each (anti) chiral superfield. From the viewpoint of the dimensional analysis of [76] having different masses m_I is irrelevant.

As a consequence one can in particular give mass to only two of the (anti) chiral multiplets. This suggests the possibility of generalizing the approach of [77, 78] to the case of $\mathcal{N}$=2 super Yang–Mills theories. A discussion of the effect of a $\mathcal{N}$=2 mass term in $\mathcal{N}$=4 supersymmetric Yang–Mills theory can be found in [81].

3.2.3 Discussion

The infrared divergences found in the calculation of the propagators of the chiral and the vector superfields are due to the fact that the vector superfield is dimensionless, so that it contains in particular, as its lowest component, the scalar C that is itself

dimensionless and hence has a propagator which behaves, in momentum space, like

$$\langle (CC)(k) \rangle \sim \frac{1}{k^4} \,.$$

The contribution of the scalar C to the $\langle VV \rangle$ propagator leads to an infrared divergence whenever a diagram contains a loop involving a VV line. In the Fermi–Feynman gauge the problem is not present because the choice $\alpha = 1$ gives a kinetic matrix of the form of equation (3.9) in the component expansion. The corresponding inverse matrix M^{-1} in (3.10) shows that no $\langle CC \rangle$ free propagator is present. On the contrary any choice $\alpha \neq 1$ produces such a propagator, *i.e.* gives a matrix M^{-1} with a non-vanishing $(M^{-1})_{11}$ entry, leading to the previously discussed problems. An explicit calculation in components with $\alpha \neq 1$ and without fixing the Wess–Zumino gauge should show problems with infrared divergences analogous to those encountered here. These problems with infrared divergences are not peculiar of the $\mathcal{N}$=4 super Yang–Mills theory, but appear in any supersymmetric gauge theory. Analogous infrared divergences in Green functions have been observed in [80, 82]. The conclusion proposed in these papers is that there exist no way to remove the infrared divergences, so that the choice α=1 is somehow necessary, at least for the computation of gauge-dependent quantities.

There are two general theorems concerning infrared divergences in quantum field theory. The first is the Kinoshita–Lee–Nauenberg theorem [83], which deals with infrared divergences in cross sections. It states that no infrared problem is present in physical cross sections of a renormalizable quantum field theory, if the appropriate sums over degenerate initial and final states, associated with soft and collinear particles, are considered. The second theorem was proved by Kinoshita, Poggio and Quinn [84] and concerns Green functions. The statement of the theorem is that the proper (one-particle irreducible) Euclidean Green functions with non-exceptional external momenta are free of infrared divergences in a renormalizable quantum field theory. A set of momenta p_i, $i = 1, 2, \ldots, n$ is said to be exceptional if any of the partial sums

$$\sum_{i \in I} p_i \,, \qquad \text{with} \;\; I \;\; \text{a subset of} \;\; \{1, 2, \ldots, n\} \,,$$

vanishes. The reason for the absence of divergences is that the external momenta, if non-exceptional, play the rôle of an infrared regulator in the Green functions.

The proof of the theorem is based on dimensional analysis which does not work in the presence of a $\frac{1}{k^4}$ propagator. In particular it fails in the case of supersymmetric

gauge theories in general gauges, because the adimensionality of the V superfield implies that factors of $\frac{1}{k^4}$ can enter the loop integrals. The apparent violation of the Kinoshita–Poggio–Quinn theorem can be understood from the viewpoint of the component formulation: supersymmetric gauge theories, being non-polynomial and thus containing an infinite number of interaction terms, are not *formally* renormalizable. The choice of the Wess–Zumino gauge makes the theory polynomial. In fact in this case no infrared divergence is found in Green functions and the general theorems are satisfied. However in the case of the $\mathcal{N}$=4 theory the choice of the WZ gauge results in a change of the ultraviolet properties of the model, at least for what concerns gauge dependent quantities, *e.g.* the propagators.

In theories with less supersymmetry, in which ultraviolet divergences are present the problem might seem less severe, since the subtraction of the ultraviolet infinities could cure also the infrared singularity.

The infrared problems discussed here are however gauge artifacts and cannot affect gauge invariant quantities. The mechanism leading to the cancellation of infrared divergences in gauge independent Green functions has not been verified in detail yet, but can be understood starting from the super Feynman rules. When the propagator $\langle VV \rangle$ is inserted into a Feynman graph it is connected to vertices which carry covariant derivatives. These derivatives come directly from the form of the action for vertices involving only V fields and from the definition of the functional derivatives for vertices involving (anti) chiral superfields. The situation in diagrams for gauge invariant quantities is such that at least one covariant derivative can always be brought to act on the V propagator by integrations by parts. In this way an additional factor of the momentum k is generated in the numerator. More precisely for the infrared problematic term $-\frac{1}{k^4}(\overline{D}^2 D^2 + D^2\overline{D}^2)\delta_4(\theta-\theta')$ one gets

$$D_\alpha \left[-\frac{1}{k^4}(\overline{D}^2 D^2 + D^2\overline{D}^2)\delta_4(\theta-\theta') \right] = -4i\frac{k_\mu \sigma^\mu_{\alpha\dot{\alpha}}}{k^4}\overline{D}^{\dot{\alpha}} D^2$$

and

$$\overline{D}^{\dot{\alpha}} \left[-\frac{1}{k^4}(\overline{D}^2 D^2 + D^2\overline{D}^2)\delta_4(\theta-\theta') \right] = 4i\frac{k_\mu \overline{\sigma}^{\mu\dot{\alpha}\alpha}}{k^4} D_\alpha \overline{D}^2 \,.$$

A rigorous and general check of this mechanism in gauge invariant Green functions has not been achieved yet.

3.3 Three- and four-point functions of $\mathcal{N}$=1 superfields

The computation of Green functions with three and four external legs will now be considered. From now on the Fermi–Feynman gauge will be fixed. The three-point functions should in principle suffer from infrared problems of the kind encountered in the preceding section. This claim will not be studied here since the calculation with $\alpha \neq 1$ is awkward, even at the one-loop level, because of the huge number of contributions. Four-point functions on the contrary should be infrared finite, because they are directly related to physical scattering amplitudes. Notice, however, that infrared divergences in the Green function with four external V lines were found in [82]. In that paper beyond the infrared singularity, it was shown that the adimensionality of the superfield V implies that it requires a non-linear renormalization, *i.e.* the renormalized field V_R is a non-linear function of the bare field V

$$V_R = f(V)\,.$$

An example of three-point function will be briefly discussed here and then the more interesting case of a four-point function will be studied in greater detail.

3.3.1 Three-point functions

The simplest three point function to consider corresponds to the correction to the vertex $\varepsilon_{IJK}\mathrm{tr}\left(\Phi^I\Phi^J\Phi^K\right)$ (or $\varepsilon^{IJK}\mathrm{tr}\left(\Phi^\dagger_I\Phi^\dagger_J\Phi^\dagger_K\right)$), which is determined by one single diagram at the one loop level. However as a less trivial example of computation of three-point functions the one-loop correction to the vertex



will be considered here. The correction at the one-loop level comes from the following diagrams (the notation for the propagators is the same as in the previous section)


$\tilde{\Phi}_b^J(q)$
$\tilde{V}_c(-p-q)$
$\longrightarrow$
$\tilde{A}(p;q)$
$\tilde{\Phi}_I^{a\dagger}(p)$
$\tilde{\Phi}_b^J(q)$
$\tilde{V}_c(-p-q)$
$\longrightarrow$
$\tilde{B}(p;q)$
$\tilde{\Phi}_I^{a\dagger}(p)$
$\tilde{\Phi}_b^J(q)$
$\tilde{V}_c(-p-q)$
$\longrightarrow$
$\tilde{C}(p;q)$
$\tilde{\Phi}_I^{a\dagger}(p)$
$\tilde{\Phi}_b^J(q)$
$\tilde{V}_c(-p-q)$
$\longrightarrow$
$\tilde{D}(p;q)$
$\tilde{\Phi}_I^{a\dagger}(p)$
$\tilde{\Phi}_b^J(q)$
$\tilde{V}_c(-p-q)$
$\longrightarrow$
$\tilde{E}(p;q)$
$\tilde{\Phi}_I^{a\dagger}(p)$



In the presence of a mass term for the (anti) chiral superfields the expressions given above are modified by the presence of the mass in the free propagators and furthermore there are two additional sets of contributions corresponding to the diagrams



The calculation of these diagrams is completely analogous to those that were presented in the previous section. All of these proper contributions can be derived using the Feynman rules. The last two diagrams are finite and vanish in the $\mathcal{N}$=4 theory, *i.e.* at m=0, as they are proportional to m^2; they will not be considered here. The other contributions will be briefly studied in the limit $m \to 0$.

The diagram $\tilde{D}(p,q)$ is zero as a consequence of the contraction among the colour indices, so that there are five contributions to be calculated. The dimensional analysis of section 1.7 gives a vanishing superficial degree of divergence, corresponding to a logarithmic divergence, for the Green function under consideration. Each diagram actually contains a logarithmically divergent term plus finite terms. A straightforward but rather lengthy computation, based on elementary D-algebra, allows to prove that

the divergent part of all the diagrams is of the form

$$I_{\log} = c\,\delta^I_J f^{abc} \int \frac{d^4k}{(2\pi)^4} d^4\theta\, \frac{1}{k^2(p-k)^2} \left\{ \Phi^\dagger_{aI}(p,\theta,\overline{\theta}) V_b(-p-q,\theta,\overline{\theta}) \Phi^J_c(q,\theta,\overline{\theta}) \right\} ,$$

where c is a constant. Moreover one finds, from each diagram a finite contribution of the form

$$\begin{aligned} I_{\text{finite}} \;=\; & \delta^I_J f^{abc} \int \frac{d^4k}{(2\pi)^4} d^4\theta\, \frac{1}{k^2(q+k)^2(p-k)^2} \\ & \Phi^\dagger_{aI}(p,\theta,\overline{\theta}) \left\{ c_1 \left[D^\alpha \overline{D}^2 D_\alpha V_b(-p-q,\theta,\overline{\theta}) \right] \Phi^J_c(q,\theta,\overline{\theta}) + \right. \\ & \left. + c_2 \sigma^\mu_{\alpha\dot{\alpha}} (q+k)_\mu \left[D^\alpha \overline{D}^{\dot{\alpha}} V_b(-p-q,\theta,\overline{\theta}) \right] \Phi^J_c(q,\theta,\overline{\theta}) \right\} , \end{aligned}$$

where c_1 and c_2 are numerical constants. The sum of all the logarithmically divergent terms contained in the diagrams $\tilde{A}$, $\tilde{B}$, $\tilde{C}$, $\tilde{E}$ and $\tilde{F}$ vanishes. The residual finite part can be shown to be zero as well. In conclusion the one-loop correction to the three-point function $\langle \Phi^\dagger V \Phi \rangle$ exactly vanishes in the Fermi–Feynman gauge. The same result can be shown to hold for the other three-point functions.

The superfield formalism does not lead to significant simplifications in the calculation of three-point functions with respect to the same computation in components (in the Wess–Zumino gauge!); actually for the Green function considered here the number of diagrams to be evaluated is approximately the same in the component formulation. However the power of superspace techniques becomes clear in the computation of four-point functions that will be considered in next subsection.

3.3.2 Four-point functions

The computation of four-point functions in components is awkward even at the one-loop level and in the Wess–Zumino gauge. In this case the choice of the WZ gauge should not lead to divergences because the complete four-point function must finally give the physical scattering amplitude. In the $\mathcal{N}$=1 superfield formulation the calculation of four-point functions, though rather lengthy, is much more simple.

In this section the computation of the one-particle irreducible one-loop correction to the Green function $\langle \Phi^\dagger \Phi \Phi^\dagger \Phi \rangle$ in the Fermi–Feynman gauge will be presented. There are several diagrams to be considered: each of them is free of ultraviolet divergences, as immediately follows from dimensional analysis. Moreover in the Fermi–Feynman gauge the single corrections are infrared safe. In conclusion one finds a finite and non-vanishing result.

The first subset of diagrams corresponds to



$$\longrightarrow \quad \tilde{A}(p;q;r)$$

together with those obtained from this one by crossing. There are a total of four diagrams of this kind. Application of the Feynman rules and steps completely analogous to those entering the evaluation of the propagators allow to find the following results.

$$\tilde{A}_1(p,q,r) = \left(\frac{1}{4}\right)^2 g^4\,\kappa N^2(N^2-1)(\delta_a^b\delta_c^d+\delta_{ac}\delta^{bd})\delta_J^I\delta_L^K\, I_{1\,bdIK}^{(A)acJL}(p,q,r)$$

where

$$\begin{aligned}
I_{1\,bdIK}^{(A)acJL}(p,q,r) \;&=\; \int \frac{d^4k}{(2\pi)^4}d^4\theta\frac{1}{k^2(p-k)^2(q+k)^2(q+k-r)^2}\cdot\\
&\cdot\Big\{\Big[(D^2\overline{D}^2)\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Big]\,\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\Phi_d^L(r,\theta,\overline{\theta})\Phi_b^J(q,\theta,\overline{\theta})+\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\cdot\\
&\cdot\Big[(D^2\overline{D}^2)\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\Big]\,\Phi_d^L(r,\theta,\overline{\theta})\Phi_b^J(q,\theta,\overline{\theta})+2\Big[(D^2\overline{D}_{\dot\alpha})\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Big]\cdot\\
&\cdot\Big[\overline{D}^{\dot\alpha}\Phi^{c\dagger}(p+q-r,\theta,\overline{\theta})\Big]\,\Phi_d^L(r,\theta,\overline{\theta})\Phi_b^J(q,\theta,\overline{\theta})+2\Big[\overline{D}_{\dot\alpha}\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Big]\cdot\\
&\cdot\Big[(D^2\overline{D}^{\dot\alpha})\Phi^{c\dagger}(p+q-r,\theta,\overline{\theta})\Big]\,\Phi_d^L(r,\theta,\overline{\theta})\Phi_b^J(q,\theta,\overline{\theta})+4\Big[(D^\alpha\overline{D}_{\dot\alpha})\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Big]\cdot\\
&\cdot\Big[D_\alpha\overline{D}^{\dot\alpha}\Phi^{c\dagger}(p+q-r,\theta,\overline{\theta})\Big]\,\Phi_d^L(r,\theta,\overline{\theta})\Phi_b^J(q,\theta,\overline{\theta})\Big\}\equiv\\
&\equiv\int\frac{d^4k}{(2\pi)^4}d^4\theta\frac{1}{k^2(p-k)^2(q+k)^2(q+k-r)^2}K_{bdIK}^{acJL}(p,q,r;\theta;\overline{\theta})
\end{aligned}\tag{3.33}$$

and the constant κ is a group theory factor defined by

$$f_{aef}f^{bfg}f_{cgh}f^{dhe}=\kappa N^2(N^2-1)(\delta_a^b\delta_c^d+\delta_a^d\delta_c^b)\,;$$

$$\tilde{A}_2(p,q,r)=\left(\frac{1}{4}\right)^2 g^4\,\kappa N^2(N^2-1)(\delta_a^d\delta_c^b+\delta_{ac}\delta^{bd})\delta_L^I\delta_J^K\, I_{2\,bdIK}^{(A)acJL}(p,q,r)\,,$$

where

$$\begin{aligned}
I_{2\,bdIK}^{(A)acJL}(p,q,r) \;&=\; \int\frac{d^4k}{(2\pi)^4}d^4\theta\frac{1}{k^2(p-k)^2(p-k-r)^2(q+k)^2(p+q-k-r)^2}\cdot\\
&\cdot K_{bdIK}^{acJL}(p,q,r;\theta;\overline{\theta})\,;
\end{aligned}\tag{3.34}$$

$$\tilde{A}_3(p,q,r) = \left(\frac{1}{4}\right)^2 g^4\,\kappa N^2(N^2-1)(\delta^b_a\delta^d_c+\delta^d_a\delta^b_c)\delta^I_J\delta^K_L\, I^{(A)acJL}_{3\,bdIK}(p,q,r)\,,$$

where

$$\begin{aligned} I^{(A)acJL}_{3\,bdIK}(p,q,r) \;=\; & \int \frac{d^4k}{(2\pi)^4}d^4\theta\frac{1}{k^2(p-k)^2(q+k)^2(q+k-r)^2}\cdot \\ & \cdot\Big\{\Big[\overline{D}^2\Phi^{a\dagger}_I(p,\theta,\overline{\theta})\Big]\left[D^2\Phi^J_b(q,\theta,\overline{\theta})\right]\Phi^{c\dagger}_K(p+q-r,\theta,\overline{\theta})\Phi^L_d(r,\theta,\overline{\theta})+ \\ & +8i\sigma^\mu_{\alpha\dot\alpha}(p-k-r)_\mu\Big[\overline{D}^{\dot\alpha}\Phi^{a\dagger}_I(p,\theta,\overline{\theta})\Big]\left[D^\alpha\Phi^J_b(q,\theta,\overline{\theta})\right]\Phi^{c\dagger}_K(p+q-r,\theta,\overline{\theta})\cdot \\ & \cdot\Phi^L_d(r,\theta,\overline{\theta})-16(p-k-r)^2\Phi^{a\dagger}_I(p,\theta,\overline{\theta})\Phi^J_b(q,\theta,\overline{\theta})\Phi^{c\dagger}_K(p+q-r,\theta,\overline{\theta})\cdot \\ & \cdot\Phi^L_d(r,\theta,\overline{\theta})\Big\}\;; \end{aligned} \tag{3.35}$$

$$\tilde{A}_4(p,q,r) = \left(\frac{1}{4}\right)^2 g^4\,\kappa N^2(N^2-1)(\delta^b_a\delta^d_c+\delta^d_a\delta^b_c)\delta^I_J\delta^K_L\, I^{(A)acJL}_{4\,bdIK}(p,q,r)\,,$$

where

$$\begin{aligned} I^{(A)acJL}_{4\,bdIK}(p,q,r) \;=\; & \int \frac{d^4k}{(2\pi)^4}d^4\theta\frac{1}{k^2(p-k)^2(p-k-r)^2(q+k)^2}\cdot \\ & \cdot\Big\{\Big[\overline{D}^2\Phi^{a\dagger}_I(p,\theta,\overline{\theta})\Big]\Phi^J_b(q,\theta,\overline{\theta})\Phi^{c\dagger}_K(p+q-r,\theta,\overline{\theta})\left[D^2\Phi^L_d(r,\theta,\overline{\theta})\right]+ \\ & +8i\sigma^\mu_{\alpha\dot\alpha}(q+k)_\mu\Big[\overline{D}^{\dot\alpha}\Phi^{a\dagger}_I(p,\theta,\overline{\theta})\Big]\Phi^J_b(q,\theta,\overline{\theta})\Phi^{c\dagger}_K(p+q-r,\theta,\overline{\theta})\cdot \\ & \cdot\left[D^\alpha\Phi^L_d(r,\theta,\overline{\theta})\right]-16(q+k)^2\Phi^{a\dagger}_I(p,\theta,\overline{\theta})\Phi^J_b(q,\theta,\overline{\theta})\Phi^{c\dagger}_K(p+q-r,\theta,\overline{\theta})\cdot \\ & \cdot\Phi^L_d(r,\theta,\overline{\theta})\Big\}\;. \end{aligned} \tag{3.36}$$

The second kind of contribution corresponds to the diagram



In this case the diagrams obtained by crossing are identical to the one depicted, so that they are accounted for by giving the correct weight to $\tilde{B}(p,q,r)$. For this contribution one finds

$$\tilde{B}(p,q,r) = \frac{g^4}{6}\,\kappa N^2(N^2-1)(\delta^b_a\delta^d_c+\delta^d_a\delta^b_c)(\delta^I_J\delta^K_L+\delta^I_L\delta^K_J)I^{(B)acJL}_{bdIK}(p,q,r)\,,$$

where

$$
\begin{aligned}
I^{(B)acJL}_{bdIK}(p,q,r) \;=\; & \int \frac{d^4k}{(2\pi)^4} d^4\theta \frac{1}{k^2(p-k)^2(p+q-r)^2(q+k)^2}\cdot \\
& \cdot \Big\{ \Big[\overline{D}^2 \Phi^{a\dagger}_I(p,\theta,\overline{\theta})\Big]\, \Phi^J_b(q,\theta,\overline{\theta}) \Phi^{c\dagger}_K(p+q-r,\theta,\overline{\theta})\, \big[D^2 \Phi^L_d(r,\theta,\overline{\theta})\big] + \\
& + 8i\sigma^\mu_{\alpha\dot\alpha}(p-k)_\mu \Big[\overline{D}^{\dot\alpha} \Phi^{a\dagger}_I(p,\theta,\overline{\theta})\Big]\, \Phi^J_b(q,\theta,\overline{\theta}) \Phi^{c\dagger}_K(p+q-r,\theta,\overline{\theta}) \cdot \\
& \cdot \big[D^\alpha \Phi^L_d(r,\theta,\overline{\theta})\big] - 16(p-k)^2 \Phi^{a\dagger}_I(p,\theta,\overline{\theta}) \Phi^J_b(q,\theta,\overline{\theta}) \Phi^{c\dagger}_K(p+q-r,\theta,\overline{\theta}) \cdot \\
& \cdot \Phi^L_d(r,\theta,\overline{\theta}) \Big\}\ . \qquad (3.37)
\end{aligned}
$$

The next one-loop correction is



$$\longrightarrow \quad \tilde C(p;q;r)$$

This contribution is trivially zero because it contains the product

$$\delta_4(\theta_1-\theta_2)\delta_4(\theta_1-\theta_2) \equiv 0\,.$$

This would not be true with a choice of gauge different from the Fermi–Feynman one, *i.e.* with $\alpha \neq 1$, because in that case there would be projectors acting on the δ's. The vanishing of $\tilde C(p,q,r)$, that is completely trivial in the superfield formulation, corresponds, in the component formulation, to a complicated cancellation among various terms coming from graphs with the same topology.

Another subset of diagrams contains



$$\longrightarrow \quad \tilde D(p;q;r)$$

plus the crossed versions of this one. There are three inequivalent crossed diagrams. The result of the calculation is the following.

$$\tilde{D}_1(p,q,r) = \left(\frac{1}{4}\right)^2 g^4\,\kappa N^2(N^2-1)(\delta_{ac}\delta^{bd}+\delta_a^d\delta_c^b)(\delta_J^I\delta_L^K-\delta_L^I\delta_J^K)I_{1\,bdIK}^{(D)acJL}(p,q,r)\,,$$

where

$$\begin{aligned} I_{1\,bdIK}^{(D)acJL}(p,q,r) &= \int \frac{d^4k}{(2\pi)^4}d^4\theta\frac{1}{k^2(p-k)^2(k+r-p-q)^2(p-k-r)^2}\cdot \\ &\cdot\Big\{\Big[\overline{D}^2\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Big]\left[D^2\Phi_b^J(q,\theta,\overline{\theta})\right]\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\Phi_d^L(r,\theta,\overline{\theta})+ \\ &+8i\sigma^\mu_{\alpha\dot\alpha}(k+r-p)_\mu\Big[\overline{D}^{\dot\alpha}\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Big]\left[D^\alpha\Phi_b^J(q,\theta,\overline{\theta})\right]\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\cdot \\ &\cdot\Phi_d^L(r,\theta,\overline{\theta})-16(k+r-p)^2\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Phi_b^J(q,\theta,\overline{\theta})\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\cdot \\ &\cdot\Phi_d^L(r,\theta,\overline{\theta})\Big\}\,; \end{aligned} \tag{3.38}$$

$$\tilde{D}_2(p,q,r) = \left(\frac{1}{4}\right)^2 g^4\,\kappa N^2(N^2-1)(\delta_{ac}\delta^{bd}+\delta_a^b\delta_c^d)(\delta_L^I\delta_J^K-\delta_J^I\delta_L^K)I_{2\,bdIK}^{(D)acJL}(p,q,r)\,,$$

where

$$\begin{aligned} I_{2\,bdIK}^{(D)acJL}(p,q,r) &= \int \frac{d^4k}{(2\pi)^4}d^4\theta\frac{1}{k^2(p-k)^2(k+r-p-q)^2(p+q-k)^2}\cdot \\ &\cdot\Big\{\Big[\overline{D}^2\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Big]\Phi_b^J(q,\theta,\overline{\theta})\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\left[D^2\Phi_d^L(r,\theta,\overline{\theta})\right]+ \\ &+8i\sigma^\mu_{\alpha\dot\alpha}(k-p-q)_\mu\Big[\overline{D}^{\dot\alpha}\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Big]\Phi_b^J(q,\theta,\overline{\theta})\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\cdot \\ &\cdot\left[D^\alpha\Phi_d^L(r,\theta,\overline{\theta})\right]-16(k-p-q)^2\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Phi_b^J(q,\theta,\overline{\theta})\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\cdot \\ &\cdot\Phi_d^L(r,\theta,\overline{\theta})\Big\}\,; \end{aligned} \tag{3.39}$$

$$\tilde{D}_3(p,q,r) = \left(\frac{1}{4}\right)^2 g^4\,\kappa N^2(N^2-1)(\delta_a^d\delta_c^b+\delta_{ac}\delta^{bd})(\delta_J^I\delta_L^K-\delta_L^I\delta_J^K)I_{3\,bdIK}^{(D)acJL}(p,q,r)\,,$$

where

$$\begin{aligned} I_{3\,bdIK}^{(D)acJL}(p,q,r) &= \int \frac{d^4k}{(2\pi)^4}d^4\theta\frac{1}{k^2(p-k)^2(k+r-p-q)^2(p+q-k)^2}\cdot \\ &\cdot\Big\{\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Phi_b^J(q,\theta,\overline{\theta})\Big[\overline{D}^2\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\Big]\left[D^2\Phi_d^L(r,\theta,\overline{\theta})\right]+ \\ &-8i\sigma^\mu_{\alpha\dot\alpha}(p-k-r)_\mu\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Phi_b^J(q,\theta,\overline{\theta})\Big[\overline{D}^{\dot\alpha}\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\Big]\cdot \\ &\cdot\left[D^\alpha\Phi_d^L(r,\theta,\overline{\theta})\right]-16(p-k-r)^2\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Phi_b^J(q,\theta,\overline{\theta})\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\cdot \\ &\cdot\Phi_d^L(r,\theta,\overline{\theta})\Big\}\,; \end{aligned} \tag{3.40}$$

$$\tilde{D}_4(p,q,r) = \left(\frac{1}{4}\right)^2 g^4\,\kappa N^2(N^2-1)(\delta_a^b\delta_c^d+\delta_{ac}\delta^{bd})(\delta_L^I\delta_J^K-\delta_J^I\delta_L^K)I_{4\,bdIK}^{(D)acJL}(p,q,r)\,,$$

where

$$\begin{aligned} I_{4\,bdIK}^{(D)acJL}(p,q,r) &= \int\frac{d^4k}{(2\pi)^4}d^4\theta\frac{1}{k^2(p-k)^2(k+r-p-q)^2(p+q-k)^2}\cdot \\ &\cdot\Big\{\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\left[D^2\Phi_b^J(q,\theta,\overline{\theta})\right]\left[\overline{D}^2\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\right]\Phi_d^L(r,\theta,\overline{\theta})+ \\ &-8i\sigma^\mu_{\alpha\dot\alpha}(p+q-k)_\mu\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\left[D^\alpha\Phi_b^J(q,\theta,\overline{\theta})\right]\left[\overline{D}^{\dot\alpha}\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\right]\cdot \\ &\cdot\Phi_d^L(r,\theta,\overline{\theta})-16(p+q-k)^2\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Phi_b^J(q,\theta,\overline{\theta})\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\cdot \\ &\cdot\Phi_d^L(r,\theta,\overline{\theta})\Big\}\,. \end{aligned} \tag{3.41}$$

The last one-loop family of diagrams is formed by the one below plus those obtained by crossing.



The resulting contributions to the Green function are

$$\tilde{E}_1(p,q,r) = \left(\frac{1}{4}\right)^2\frac{g^4}{2}\,\kappa N^2(N^2-1)\delta_a^b\delta_c^d+\delta_{ac}\delta^{bd})\delta_J^I\delta_L^K I_{1\,bdIK}^{(E)acJL}(p,q,r)\,,$$

where

$$\begin{aligned} I_{1\,bdIK}^{(E)acJL}(p,q,r) &= \int\frac{d^4k}{(2\pi)^4}d^4\theta\frac{1}{k^2(p-k)^2(p+q-k)^2}\cdot \\ &\cdot\Big\{\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Phi_b^J(q,\theta,\overline{\theta})\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\Phi_d^L(r,\theta,\overline{\theta})\Big\}\,; \end{aligned} \tag{3.42}$$

$$\tilde{E}_2(p,q,r) = \left(\frac{1}{4}\right)^2\frac{g^4}{2}\,\kappa N^2(N^2-1)(\delta_a^d\delta_c^b+\delta_{ac}\delta^{bd})\delta_L^I\delta_J^K I_{2\,bdIK}^{(E)acJL}(p,q,r)\,,$$

where

$$\begin{aligned} I_{2\,bdIK}^{(E)acJL}(p,q,r) &= \int\frac{d^4k}{(2\pi)^4}d^4\theta\frac{1}{k^2(p-k)^2(r-k)^2}\cdot \\ &\cdot\Big\{\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Phi_b^J(q,\theta,\overline{\theta})\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\Phi_d^L(r,\theta,\overline{\theta})\Big\}\,; \end{aligned} \tag{3.43}$$

$$\tilde{E}_3(p,q,r) = \left(\frac{1}{4}\right)^2 \frac{g^4}{2}\,\kappa N^2(N^2-1)(\delta_a^b\delta_c^d + \delta_{ac}\delta^{bd})\delta_L^I\delta_J^K I_{3\,bdIK}^{(E)acJL}(p,q,r)\,,$$

where

$$\begin{aligned} I_{3\,bdIK}^{(E)acJL}(p,q,r) &= \int \frac{d^4k}{(2\pi)^4} d^4\theta \frac{1}{k^2(p+q-k)^2(k-r)^2}\cdot \\ &\cdot \left\{\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Phi_b^J(q,\theta,\overline{\theta})\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\Phi_d^L(r,\theta,\overline{\theta})\right\}\,; \end{aligned} \tag{3.44}$$

$$\tilde{E}_4(p,q,r) = \left(\frac{1}{4}\right)^2 \frac{g^4}{2}\,\kappa N^2(N^2-1)(\delta_a^d\delta_c^b + \delta_{ac}\delta^{bd})\delta_L^I\delta_J^K I_{4\,bdIK}^{(E)acJL}(p,q,r)\,,$$

where

$$\begin{aligned} I_{4\,bdIK}^{(E)acJL}(p,q,r) &= \int \frac{d^4k}{(2\pi)^4} d^4\theta \frac{1}{k^2(q+k)^2(k+r-p)^2}\cdot \\ &\cdot \left\{\Phi_I^{a\dagger}(p,\theta,\overline{\theta})\Phi_b^J(q,\theta,\overline{\theta})\Phi_K^{c\dagger}(p+q-r,\theta,\overline{\theta})\Phi_d^L(r,\theta,\overline{\theta})\right\}\,. \end{aligned} \tag{3.45}$$

The sum of all the preceding terms results in a finite and non vanishing total one-loop correction to the four point function. The final expression contains terms with six different tensorial structures

$$\langle\Phi^\dagger\Phi\Phi^\dagger\Phi\rangle = \kappa\left(\frac{1}{4}\right)^2 g^4N^2(N^2-1)\sum_{i=1}^{6}G^{(i)}\,,$$

where

$$\begin{aligned} G^{(1)} &= \delta_a^b\delta_c^d\delta_J^I\delta_L^K\left(I_{1\,bdIK}^{(A)acJL} + I_{3\,bdIK}^{(A)acJL} + \frac{2}{3}I_{bdIK}^{(B)acJL} - I_{2\,bdIK}^{(D)acJL} + \right. \\ &- \left. I_{4\,bdIK}^{(D)acJL} + I_{1\,bdIK}^{(E)acJL} + I_{3\,bdIK}^{(E)acJL}\right) \\ G^{(2)} &= \delta_{ac}\delta^{bd}\delta_J^I\delta_L^K\left(I_{1\,bdIK}^{(A)acJL} + I_{1\,bdIK}^{(D)acJL} - I_{2\,bdIK}^{(D)acJL} + I_{3\,bdIK}^{(D)acJL} + \right. \\ &- \left. I_{4\,bdIK}^{(D)acJL} + I_{1\,bdIK}^{(E)acJL} + I_{3\,bdIK}^{(E)acJL}\right) \\ G^{(3)} &= \delta_a^b\delta_c^d\delta_L^I\delta_J^K\left(I_{2\,bdIK}^{(A)acJL} + \frac{2}{3}I_{bdIK}^{(B)acJL} + I_{2\,bdIK}^{(D)acJL} + I_{4\,bdIK}^{(D)acJL}\right) \\ G^{(4)} &= \delta_a^d\delta_c^b\delta_L^I\delta_J^K\left(I_{2\,bdIK}^{(A)acJL} + I_{4\,bdIK}^{(A)acJL} + \frac{2}{3}I_{bdIK}^{(B)acJL} - I_{1\,bdIK}^{(D)acJL} - \right. \\ &- \left. I_{3\,bdIK}^{(D)acJL} + I_{2\,bdIK}^{(E)acJL} + I_{4\,bdIK}^{(E)acJL}\right) \\ G^{(5)} &= \delta_a^d\delta_c^b\delta_J^I\delta_L^K\left(I_{3\,bdIK}^{(A)acJL} + \frac{2}{3}I_{bdIK}^{(B)acJL} + I_{1\,bdIK}^{(D)acJL} + I_{3\,bdIK}^{(E)acJL}\right) \\ G^{(6)} &= \delta_{ac}\delta^{bd}\delta_L^I\delta_J^K\left(I_{4\,bdIK}^{(A)acJL} - I_{1\,bdIK}^{(D)acJL} + I_{2\,bdIK}^{(D)acJL} - I_{3\,bdIK}^{(D)acJL} + \right. \\ &+ \left. I_{4\,bdIK}^{(D)acJL} + I_{2\,bdIK}^{(E)acJL} + I_{4\,bdIK}^{(E)acJL}\right) \end{aligned}$$

In the presence of a mass term for the (anti) chiral superfields the expressions given above are slightly modified by the presence of the mass in the free propagators and furthermore there are two additional sets of contributions corresponding to the diagrams



$$\longrightarrow \quad \tilde{F}(p;q;r)$$



$$\longrightarrow \quad \tilde{G}(p;q;r)$$

Both of these graphs give corrections proportional to m^2, that can be calculated much in the same way as the previous ones.

With a different choice of gauge the computation of four-point Green functions is more complicated. Single diagrams involving vector superfield propagators contain new contributions, some of which are infrared divergent. The correction $\tilde{C}$ is not zero anymore, because there are projection operators acting on the δ-functions. Moreover further diagrams must be included in the calculation at the same order as a consequence of the non-vanishing of the one-loop correction to the vertices. For example one must consider diagrams such as



The techniques illustrated in this chapter for the calculation of Green functions in the $\mathcal{N}$=1 formalism can be applied to correlation functions of composite operators as well. Green functions of gauge invariant composite operators such as those that form the multiplet of currents (equations (2.17) and (2.18)) play a crucial rôle in the correspondence with type IIB superstring theory on AdS space to be discussed in chapter 5. The application of $\mathcal{N}$=1 superspace to this problem will be sketched in the final chapter. It can be shown that the extension of the formalism to the case of composite operators is rather straightforward, the fundamental difference being that the complete Green functions and not the proper parts must be considered.

Chapter 4

Nonperturbative effects in $\mathcal{N} = 4$ super Yang–Mills theory

In the last chapter the quantum properties of $\mathcal{N}$=4 supersymmetric Yang–Mills theory have been analyzed at the perturbative level, this chapter is devoted to the study of some non-perturbative effects. Instanton calculus will be employed to compute various Green functions of gauge invariant composite operators in the semiclassical approximation.

It is well known that in gauge field theories non-perturbative effects can modify in a dramatic way the perturbative structure of the vacuum. Instanton calculus has proved to be an extremely powerful tool in the analysis of non-perturbative phenomena in quantum field theories, particularly in the case of supersymmetric gauge theories. The rôle of instantons in Yang–Mills theories was pointed out in [85] at the classical level and then at the quantum level in a fundamental work by 't Hooft [86]. Instantons are non trivial finite action solutions of the classical Euclidean equations of motion of the model. The computation of Green functions in the instanton background allows to study non-perturbative properties of the quantum theory. In non-supersymmetric non Abelian gauge theories the contribution of instantons to Green functions turns out to be divergent in the semiclassical approximation [86]. However in the supersymmetric case explicit instanton calculations can be performed leading to finite and well defined results, thanks to the cancellation between fermionic and bosonic quantum fluctuations. Compensations of quantum effects due

bosons and fermions take place in the instanton background and allow to explicitly compute vacuum expectation values (vev's) of composite operators (condensates). These cancellations are reminiscent of boson and fermion compensations in internal loops at the perturbative level. The condensates play the rôle of order parameters in non-perturbative phenomena such as for instance the spontaneous breaking of chiral symmetries or even of supersymmetry in certain models [87]. Furthermore instanton calculus together with holomorphy properties, has allowed to derive some exact results in supersymmetric theories [33] and, more recently, to obtain non trivial checks of the exact result proposed in [34].

As was discussed in previous chapters supersymmetric gauge theories typically possess a moduli space of vacua parameterized by the vev's of scalar fields. The literature on instanton effects in asymptotic free theories includes examples in which the computations are performed with zero vev's for all the scalars [87] as well as examples in which non vanishing vev's for the scalar fields parameterizing the moduli space are considered [88]. The $\mathcal{N}$=4 super Yang–Mills theory has a (exactly) vanishing β-function and a moduli space of vacua, $\mathcal{M}=\mathbb{R}^{6k}\times\mathcal{S}_k$, where k is the rank of the gauge group and $\mathcal{S}_k$ the group of permutations of k elements, parameterized by the vev's of six real scalar fields, $\langle\varphi^i\rangle$. The superconformal phase, corresponding to the origin of $\mathcal{M}$, *i.e.* $\langle\varphi^i\rangle$=0, $\forall i$, cannot be obtained naively as the limit of the theory in the Coulomb phase. In this chapter the approach of [87] will be applied to the computation of various condensates corresponding to operators in the supermultiplet of currents (2.17), (2.18). Four-, eight- and sixteen-point functions of bilinear operators belonging to the current supermultiplet will be computed in the semiclassical approximation in the case of a SU(2) gauge group. In particular the exact complete spatial dependence will be determined for a four-point function at lowest order.

Due to the vanishing of the β-function and of the associated chiral anomalies, the correlation functions that will be studied, unlike the cases encountered in theories with less supersymmetry, receive a non-vanishing perturbative contribution as well. These Green functions will reconsidered in the final chapter in the context of the correspondence with type IIB superstring theory compactified on $\mathrm{AdS}_5\times S^5$. It will be shown that the difficulties encountered in perturbative calculations in chapter 3 do not affect the computations discussed here.

The chapter is organized as follows. The general features of instanton calculus in supersymmetric gauge theories are briefly discussed in sections 1, 2 and 3. Sections 4.4, 4.5 and 4.6 report on original results in the case of $\mathcal{N}$=4 supersymmetric Yang–

Mills theory, following [89]. The final section describes the generalizations of the results of [89] to the case of a SU(N) gauge group and to the K-instanton sector in the large N limit, that were given respectively in [90] and [91, 92].

4.1 Instanton calculus in non Abelian gauge theories

In this section the semiclassical quantization in the background of an instanton configuration will be reviewed for the case of a pure Yang–Mills theory. The extension to fermions will be discussed in the next section.

In quantum field theory one is interested in computing vacuum expectation values of operators $\mathcal{O}(\phi)$, that in general are polynomials in the elementary fields ϕ. In Euclidean space one must calculate

$$\langle \mathcal{O}(x_1, \dots, x_n) \rangle = \int [\mathcal{D}\phi] e^{-\frac{S[\phi]}{\hbar}} \mathcal{O}(\phi(x_1), \dots, \phi(x_n)) \,, \tag{4.1}$$

where the dependence on $\hbar$ has been explicitly indicated in order to discuss the semiclassical approximation.

The classical vacuum of the theory corresponds to a constant expectation value $\langle \phi \rangle$ for the elementary fields and a perturbative series is obtained expanding the interaction part of the action $S[\phi]$ for small fluctuations, $\delta\phi$, around the classical vacuum. In instanton calculations one considers a situation in which there exists a non-trivial solution of the classical equations of motion with finite Euclidean action. Denoting by $\overline{\phi}$ such a field configuration one can then study quantum fluctuation around $\phi = \overline{\phi}$ in the limit $\hbar \to 0$. Putting

$$\phi(x) = \overline{\phi}(x) + \delta\phi(x) = \overline{\phi}(x) + \hbar^{1/2} \eta(x)$$

and expanding $S[\phi]$ in powers of $\hbar$ in the form

$$S[\phi] = S[\overline{\phi}] + \frac{\hbar}{2} \int d^4x d^4y \left(\frac{\delta^2 S}{\delta\phi(x)\delta\phi(y)} \right) \Big|_{\phi=\overline{\phi}} \eta(x)\eta(y) + O(\hbar^{3/2}) \,,$$

the vacuum expectation value of an operator $\mathcal{O}[\phi]$ can be evaluated in the semiclas-

sical limit by a saddle point approximation [1]

$$\begin{aligned}\langle \mathcal{O}(x_1,\dots,x_n)\rangle &= \int [\mathcal{D}\eta] e^{-\frac{S[\overline{\phi}+\eta]}{\hbar}} \mathcal{O}[(\overline{\phi}+\eta)(x_i)] = \\ &= \mathcal{O}(\overline{\phi})\, e^{-\frac{\overline{S}}{\hbar}} \left[\det\left(\frac{\delta^2 S}{\delta\phi(x)\delta\phi(y)}\right)\Big|_{\phi=\overline{\phi}}\right]^{-\frac{1}{2}} (1+O(\hbar))\ . \end{aligned} \tag{4.2}$$

Notice that in principle one could consider a classical solution $\overline{\phi}$ with infinite Euclidean action, but this would give a vanishing contribution in the semiclassical approximation because of the factor $e^{-S[\overline{\phi}]/\hbar}$ in (4.2).

4.1.1 Yang–Mills instantons

The classification of non-trivial solutions of the classical Euclidean equations of motion for the Yang–Mills system has been given in [85], where an explicit solution has also been constructed. Another (multi-center) solution has been proposed by 't Hooft, who also studied the semiclassical quantization [86]. The case of a SU(2) gauge group will be discussed first and then the extension to SU(N) will be sketched.

The Euclidean Yang–Mills action is

$$S_{\rm YM}[A] = -\frac{1}{4 d_{\mathbf{r}} g^2}\int d^4x\, \mathrm{tr}\,(F_{\mu\nu}F^{\mu\nu})\ , \tag{4.3}$$

where the gauge field A_μ and the field strength $F_{\mu\nu}$ are respectively

$$\begin{aligned} A_\mu &= A^a_\mu T_a \\ F^{\mu\nu} &= \partial^\mu A^\nu - \partial^\nu A^\mu + [A^\mu, A^\nu] = \left(\partial_\mu A^a_\mu - \partial_\nu A^a_\mu + i f^a{}_{bc} A^b_\mu A^c_\nu\right) T_a \end{aligned}$$

and T^a are generators of the gauge group in the representation $\mathbf{r}$, satisfying $[T^a, T^b] = i f^{ab}{}_c T^c$ and $\mathrm{tr}\left(T^a T^b\right) = \delta^{ab} d_{\mathbf{r}}$, $d_{\mathbf{r}}$ being the Dynkin index of the representation. Gauge transformations take the form

$$A_\mu(x) \longrightarrow A^\Omega_\mu(x) = \Omega(x)\left[A_\mu(x) + i\partial_\mu\right]\Omega^\dagger(x)\,,$$

with $\Omega(x) = e^{i\alpha_a(x)T^a}$.

The study of finite Euclidean action configurations is performed much in the same way as for the monopoles of the Georgi–Glashow model in appendix B. The requirement of finite action implies that A_μ must approach asymptotically, *i.e.* for

[1] This relation holds for bosonic fields ϕ, see the next section for the case of fermions.

$|x| \to \infty$, a pure gauge configuration. This leads to a topological classification of the field configurations in homotopy classes. The quantity which characterizes the finite energy solutions is the Pontryagin index

$$K[A] = \frac{1}{32\pi^2} \int d^4x F^a_{\mu\nu} \tilde{F}^{\mu\nu}_a \,, \tag{4.4}$$

where $\tilde{F} = \frac{1}{2}\varepsilon_{\mu\nu\rho\sigma}F^{\rho\sigma}$. The Yang–Mills action satisfies a "Bogomol'nyi bound"

$$S_{\rm YM}[A] = \pm\frac{8\pi^2}{g^2}K[A] + \frac{1}{8\pi^2}\int d^4x \left(F^a_{\mu\nu} \mp \tilde{F}^a_{\mu\nu}\right)\left(F^{\mu\nu}_a \mp \tilde{F}^{\mu\nu}_a\right) \geq \frac{8\pi^2}{g^2}\left|K[A]\right| \,. \tag{4.5}$$

Solutions of the classical equations of motion correspond to local minima of the action functional, so that in each homotopy class they saturate the bound in (4.5). As a consequence they are determined by the condition

$$F^a_{\mu\nu} = \pm\tilde{F}^a_{\mu\nu} \,.$$

In the K=0 sector the minimum is reached on the trivial vacuum, A_μ=0, while the configurations minimizing $S[A]$ for $|K| \neq 0$ are (anti) instantons. Self-dual and anti self-dual configurations are referred to as respectively instantons and anti-instantons.

In the K=1 sector and for a gauge group SU(2) the explicit solution was found in [85, 86] through the ansatz

$$\overline{A}_\mu = i\frac{\overline{\sigma}_{\mu\nu}}{2}\partial^\nu \log\Phi \,, \tag{4.6}$$

where Φ is real scalar field. Substituting into the equations of motion one obtains the equation for Φ

$$\frac{\Box\Phi}{\Phi} = 0 \,.$$

A solution can be written in the form

$$\overline{\Phi} = 1 + \frac{\rho^2}{(x-x_0)^2} \,. \tag{4.7}$$

$\overline{\Phi}$ depends on 5 free parameters, ρ associated with the "size" of the instanton and x_0 with its position. The corresponding instanton solution is

$$\overline{A}_\mu = -2i\rho^2\overline{\eta}^a_{\mu\nu}\frac{y^\nu}{y^2(y^2+\rho^2)} \,, \tag{4.8}$$

where $y = x - x_0$ and $\overline{\eta}^a_{\mu\nu}$ are the anti self-dual 't Hooft symbols

$$\overline{\eta}^a_{\mu\nu} = -\overline{\eta}^a_{\nu\mu} = \begin{cases} \varepsilon^{a\mu\nu} & \text{for} \quad \mu,\nu = 1,2,3 \\ -\delta^{a\mu} & \text{for} \quad \nu = 4\,. \end{cases}$$

The general solution depends on 3 additional parameters corresponding to the freedom of performing global gauge rotations. Actually the Yang–Mills system is classically invariant under the four-dimensional conformal group, which partially overlaps with global gauge transformations. In [93] it has been proved that the most general one-instanton solution depends on 8 parameters, referred to as collective coordinates or moduli, that in the following will be denoted by β_i. The moduli space, *i.e.* the space parameterized by the coordinates β_i, will be indicated by $\mathcal{M}$. The field (4.8) has a singularity at y=0, which can be removed by a correspondingly singular gauge transformation, so that one can construct a solution with no singularities at any x

$$\overline{A}_\mu = -2ie^{i\theta_a T^a}\eta^a_{\mu\nu}\frac{y^\nu}{(y^2+\rho^2)}e^{-i\theta_a T^a}\,, \tag{4.9}$$

where $\eta^a_{\mu\nu}$ are the self-dual 't Hooft symbols

$$\eta^a_{\mu\nu} = -\eta^a_{\nu\mu} = \begin{cases} \varepsilon^{a\mu\nu} & \text{for} \quad \mu,\nu = 1,2,3 \\ \delta^{a\mu} & \text{for} \quad \nu = 4\,, \end{cases}$$

which satisfy

$$\sigma_{\mu\nu} = \eta^a_{\mu\nu}\sigma_a\,,$$

with σ^a, $a = 1,2,3$ the standard Pauli matrices. The associated field strength is

$$\overline{F}_{\mu\nu} = F_{\mu\nu}[\overline{A}] = -2ie^{i\theta_a T^a}\sigma_{\mu\nu}\frac{\rho^2}{(y^2+\rho^2)^2}e^{-i\theta_a T^a}\,. \tag{4.10}$$

The ansatz of (4.7) can be generalized to a n-instanton solution considering the scalar field

$$\Phi(x) = 1 + \sum_{i=1}^{n}\frac{\rho_i^2}{(x - x_0^{(i)})^2}\,.$$

Substituting into (4.6) yields a solution, $\overline{A}^{(n)}_\mu$, with singularities at $x_\mu = x^{(i)}_{0\,\mu}$, which can be made regular by a suitable singular gauge transformation. For such a configuration one can prove that

$$S[\overline{A}^{(n)}] = nS[\overline{A}^{(1)}]\,,$$

so that $\overline{A}_\mu^{(n)}(x)$ is a solution in the homotopy class $K[A]{=}n$. The number of collective coordinates for the n-instanton solution is $8n-3$ (with $n>1$). There are 8 moduli for each single instanton, but one must subtract 3 global rotations that are already accounted for by the relative rotations. This result can be proved to hold in general, even if the n instantons are not widely separated, [94].

4.1.2 Semiclassical quantization

To perform a semiclassical quantization around an instanton configuration according to (4.2) one needs the expansion of $S[A]$ around $A_\mu=\overline{A}_\mu$, obtained putting $A_\mu=\overline{A}_\mu+Q_\mu$, where Q_μ denotes the quantum fluctuation. A straightforward calculation gives the gauge kinetic operator in the instanton background

$$M^{ab}_{\mu\nu}=\left(\frac{\delta^2 S[A]}{\delta A^a_\mu \delta A^b_\nu}\right)\Big|_{\overline{A}}=\left[-D^2(\overline{A})\delta_{\mu\nu}+D_\mu(\overline{A})D_\nu(\overline{A})-2\overline{F}_{\mu\nu}\right]\delta^{ab}\,,$$

where $D_\mu(\overline{A})=\partial_\mu+i[\overline{A}_\mu,\,.\,]$ is the covariant derivative.

In the instanton background the Faddeev–Popov quantization must be suitably modified, because the kinetic operator $M_{\mu\nu}$ possesses 8 new zero modes (in the case of a SU(2) gauge group under consideration) beyond those associated with gauge symmetry. These new zero-modes correspond to the symmetries of the action $S[A]$ that are broken by the instanton configuration and are related to the 8 collective coordinates β_i in the general solution (4.9). In fact since $\overline{A}_\mu$ is a solution for any β_i

$$\int d^4y\left(\frac{\delta^2 S[A]}{\delta A_\mu \delta A_\nu}\right)\Big|_{\overline{A}_\beta}\frac{\partial\overline{A}_\nu}{\partial\beta_i}(y,\beta)=0\,,$$

so that (4.2) is formally divergent. This divergence is completely analogous to those introduced in the partition function of gauge theories by the functional integration along the gauge orbits, that is dealt with by the Faddeev–Popov procedure. The correct way of performing the quantization in the instanton background was discussed in [95]. The general instanton configuration will be parameterized as

$$\overline{A}^\Omega_\mu(x,\beta)=\Omega^\dagger\overline{A}_\mu\Omega+i\Omega^\dagger\partial_\mu\Omega\,,$$

where Ω denotes an arbitrary gauge transformation and β the set collective coordinates, 8 in the case of SU(2) and $K[A]{=}1$ and in general $4NK[A]$ in the K-instanton sector for a SU(N) gauge group. The basic idea is then to constrain *à la* Faddeev–Popov the quantum fluctuations Q_μ to be orthogonal to $\overline{A}^\Omega_\mu$ as Ω varies in G and β_i

in the moduli space $\mathcal{M}$. In the partition function of the system [2]

$$Z = \int [\mathcal{D}A] e^{-S[A]}$$

one inserts

$$1 = \Delta_{\rm FP} \int_G \prod_{a,x} [\mathcal{D}h^a(x)] \int_{\mathcal{M}} \prod_i d\beta_i \delta \left((A_\mu - \overline{A}^\Omega_\mu, \frac{\delta \overline{A}^\Omega_\mu}{\delta h^a}) \right) \delta \left((A_\mu - \overline{A}^\Omega_\mu, \frac{\partial \overline{A}^\Omega_\mu}{\partial \beta_i}) \right) , \tag{4.11}$$

where $\Omega(x)$ is parameterized as $\Omega(x) = e^{ih^a(x)T_a}$ and the scalar product in the arguments of the δ-functions is

$$(f_\mu, g^\mu) = \frac{1}{2} \int d^4x \, {\rm tr} \, (f_\mu(x) g^\mu(x)) \ . \tag{4.12}$$

Equation (4.11) defines the Faddeev–Popov determinant $\Delta_{\rm FP}$. Some algebraic steps allow write Z in the form

$$\begin{aligned} Z = & \int [\mathcal{D}A_\mu] \int \prod_i d\beta_i \, e^{-S[A]} \Delta_{\rm FP}(A, \beta) \cdot \\ & \cdot \delta \left({\rm tr} \left[T^a D_\mu(\overline{A}) \left(A^\mu - \overline{A} \right) (x) \right] \right) \delta \left((A_\mu - \overline{A}_\mu, \frac{\partial \overline{A}_\mu}{\partial \beta_i}) \right) \ . \end{aligned}$$

In this relation an infinite constant, coming from the integration over the group algebra, has been dropped, since in the following the above construction will be employed in the computation of normalized vev's of operators. The Faddeev–Popov determinant in the above expression can be computed much in the same way as in the case of the ordinary of gauge theories.

To compute expectation values in the semiclassical approximation one only needs $\Delta_{\rm FP}(A, \beta)$ on the classical solution $\overline{A}$. It can be shown that in this case $\Delta_{\rm FP}$ can be written as

$$\Delta_{\rm FP}(A, \beta) \Big|_{A=\overline{A}} = \det_{\substack{x,y \\ a,b}} \left(D^2_{ab}(\overline{A}) \delta(x-y) \right) \det_{i,j} \left(d_{ij}(A, \beta) \right) \ , \tag{4.13}$$

where

$$d_{ij}(A, \beta) = (\hat{a}^{(i)}_\mu(\beta), \hat{a}^{\mu(j)}(\beta))$$

[2] From now on $\hbar$ will be set equal 1.

The vectors $\hat{a}$ in the above equation are given by

$$\hat{a}_\mu^{(i)\,a}(x,\beta) = \frac{\partial \overline{A}_\mu^a}{\partial \beta_i} + \xi_\mu^{(i)\,a}(x)$$

and $\xi_\mu^{(i)\,a}$) are determined by the transversality condition

$$D_\mu^{ab}(\overline{A})\hat{a}_{\mu\,b}^{(i)}(x,\beta) = 0\,.$$

In conclusion introducing ghost fields, c and $\overline{c}$ to rewrite the determinants as Gaussian integrals the expectation value of a gauge invariant operator $\mathcal{O}[A]$ in the semiclassical approximation takes the form

$$\begin{aligned}
&\langle \mathcal{O}(x_1,\dots,x_n)\rangle_{\overline{A}} = e^{-\frac{8\pi^2}{g^2}K[\overline{A}]}\cdot \\
&\cdot \frac{\displaystyle\int [\mathcal{D}Q_\mu \mathcal{D}c\mathcal{D}\overline{c}] \prod_i d\beta_i \frac{\|\hat{a}\|}{\sqrt{2\pi}}\, e^{-\frac{1}{2}\int Q^\mu M^{\text{g.f.}}_{\mu\nu} Q^\nu - \int \overline{c} D^2(\overline{A}) c}\, \mathcal{O}[\overline{A}]}{\displaystyle\int [\mathcal{D}Q_\mu \mathcal{D}c\mathcal{D}\overline{c}] e^{-\frac{1}{2}\int Q^\mu M^{(0)\text{g.f.}}_{\mu\nu} Q^\nu - \int \overline{c}\Box c}} = \\
&= e^{-\frac{8\pi^2}{g^2}K[\overline{A}]} \int \prod_i d\beta_i \frac{\|\hat{a}\|}{\sqrt{2\pi}}\, \mathcal{O}[\overline{A}] \frac{\left(\det{}' M^{\text{g.f.}}_{\mu\nu}\right)^{-\frac{1}{2}} \det D^2(\overline{A})}{\left(\det{}' M^{(0)\text{g.f.}}_{\mu\nu}\right)^{-\frac{1}{2}} \det \Box}\,,
\end{aligned} \tag{4.14}$$

where the 'gauge-fixed' kinetic operators $M^{\text{g.f.}}_{\mu\nu}$ and $M^{(0)\text{g.f.}}_{\mu\nu}$ are

$$M^{\text{g.f.}}_{\mu\nu} = -D^2(\overline{A})\delta_{\mu\nu} - 2F_{\mu\nu}[\overline{A}]\,, \qquad M^{(0)\text{g.f.}}_{\mu\nu} = -\Box\delta_{\mu\nu}\,.$$

4.1.3 Bosonic zero-modes

To explicitly compute expectation values through equation (4.14) one needs to know the zero-mode vectors $\hat{a}_\mu^{(i)}$, whose norms enter the integration measure. Since the kinetic operator is modified by the gauge-fixing term the vectors $\hat{a}_\mu^{(i)}$ are given by

$$\hat{a}_\mu^{(i)}(x) = \frac{\partial \overline{A}_\mu}{\partial \beta_i}(x,\beta) + D_\mu(\overline{A})\Lambda^{(i)}(x)\,,$$

for a suitable function $\Lambda^{(i)}(x)$, such that

$$\hat{a}_\mu^{(i)}(x) \in \text{Ker}\,\left[M^{\text{g.f.}}_{\mu\nu}\right]\,.$$

To compute the zero-mode vectors it is convenient to refer to the singular instanton solution. For the case of a SU(2) gauge group, after rescaling $A_\mu \to \frac{1}{g}A_\mu$, it reads

$$\overline{A}_\mu(x,\beta) = -\frac{i}{g} e^{i\theta_a T^a} \left[\overline{\sigma}_{\mu\nu} \frac{\rho^2 (x-x_0)_\mu}{(x-x_0)^2[(x-x_0)^2+\rho^2]}\right] e^{-i\theta_a T^a}\,, \tag{4.15}$$

where the 8 moduli are $\beta_i = (x_0^\mu, \rho, \theta_a)$.

Zero-modes associated with translations.

In this case $\beta_i \to x_0^\mu$ and there are four zero-modes. The transverse vectors are

$$\hat{a}_{\mu(\nu)} = \frac{\partial}{\partial x_0^\nu}\overline{A}_\mu + D_\mu(\overline{A})\overline{A}_\nu = F_{\mu\nu}[\overline{A}]\,.$$

The computation of the norm is straightforward using (4.15), the result is

$$\|\hat{a}_{\mu(\nu)}\| = \frac{2\sqrt{2}\,\pi}{g}\,.$$

Zero-modes associated with dilatations.

The corresponding collective coordinate is the size of the instanton ρ. $\hat{a}_\mu^{(\mathrm{dil})}$ is simply given by

$$\hat{a}_\mu^{(\mathrm{dil})} = \frac{\partial\overline{A}_\mu}{\partial\rho} = -\frac{2i}{g}\overline{\sigma}_{\mu\nu}\frac{\rho y^\nu}{(y^2+\rho^2)^2}\,,$$

so that for the norm one obtains

$$\|\hat{a}_\mu^{(\mathrm{dil})}\| = \frac{4\pi}{g}\,.$$

Zero-modes associated with gauge rotations.

The relevant collective coordinates are the angles θ_a in (4.15) and one can put

$$\hat{a}_\mu^{(a)} = \frac{\partial\overline{A}_\mu}{\partial\theta_a} + D_\mu(\overline{A})\Lambda^{(a)} = D_\mu(\overline{A})\left[-\frac{i}{g}T^a + \Lambda^{(a)}\right] \equiv D_\mu(\overline{A})\phi^{(a)}\,,$$

where the scalar field $\phi^{(a)}$, defined by the last equality in the above equation, must satisfy

$$D^2(\overline{A})\phi^{(a)} = 0\,, \qquad \lim_{|x|\to\infty}\phi^{(a)} = -\frac{i}{g}T^{(a)}\,.$$

The solution is

$$\phi^{(a)} = \left(-\frac{i}{g}T^{(a)}\right)\frac{r^2}{r^2+\rho^2} \qquad \text{with} \quad r = |x|\,.$$

The norms are then

$$\|\hat{a}_\mu^{(a)}\| = \frac{2\pi\rho}{g}\,.$$

The above eight vectors can be shown to be orthogonal with respect to the scalar product (4.12).

Having determined the vectors $\hat{a}_\mu$ one can thus replace the integration over the zero-modes in (4.14) with an integration over the instanton collective coordinates and write the integration measure as

$$\begin{aligned} d\mu_{\rm B} &= \prod_i d\beta_i \frac{\|\hat{a}\|}{\sqrt{2\pi}} = \frac{1}{(2\pi)^4}\left(\frac{2\sqrt{2}\pi}{g}\right)^4 \frac{4\pi}{g}\left(\frac{2\pi\rho}{g}\right)^3 d^4x_0 d\rho d^3\theta = \\ &= \frac{2^7\pi^4}{g^8}\rho^3 d^4x_0 d\rho d^3\theta\,. \end{aligned} \tag{4.16}$$

The generalization to larger gauge groups, in particular to SU(N), is rather straightforward [96, 97]. First it can be shown [98], that the mapping of S^3 into a generic group G, which determines the topological classification of the instanton solutions, can be continuously deformed to a mapping into SU(2). Hence the classification of classical solutions into homotopy classes holds in general. Furthermore one can construct a solution in the case of a gauge group G by the embedding of a SU(2) instanton. In particular for G=SU(N) one can consider a configuration

$$\overline{A}_\mu = \frac{2i}{g}\frac{\overline{\eta}^a_{\mu\nu}x^\nu}{x^2(x^2+\rho^2)}\frac{\lambda_k}{2}\,,$$

where λ_k, $k=1,2,3$, are the first three generators of the fundamental representation of SU(N). Under the SU(2) subgroup of SU(N) $(\lambda_1,\lambda_2,\lambda_3)$ transforms as a triplet, whereas the other generators are organized into $2(N-2)$ doublets and the remaining are singlets. In conclusion, in addition to eight zero-modes of the SU(2) instanton, there are $4(N-2)$ additional zero-modes. The corresponding vectors take the form [97]

$$\hat{a}^r_\mu = D_\mu(\overline{A})\left[\frac{\lambda^r}{g}\left(\frac{x^2}{x^2+\rho^2}\right)^{\frac{1}{2}}\right]\,.$$

4.2 Instanton calculus in the presence of fermionic fields

In this section the semiclassical quantization in the background of an instanton configuration will be generalized to the case of a non Abelian gauge theory with interacting

fermions.

The Euclidean action of the model is

$$S[\psi,\overline{\psi},A] = S_{\rm YM}[A] + S_A[\psi,\overline{\psi}] = S_{\rm YM}[A] + \int d^4x\,\overline{\psi}\,[iD_\mu(\mathbf{r})]\,\overline{\sigma}^\mu\psi\,, \tag{4.17}$$

where $S_{\rm YM}$ is the Yang–Mills action (4.3) and $D_\mu(\mathbf{r})$ denotes the covariant derivative in the representation $\mathbf{r}$ of the gauge group G (see appendix A for the notations). In (4.17) ψ is a Weyl fermion; an equivalent description can be given in terms of a Dirac spinor Ψ, introducing a fictitious right chirality Weyl spinor and writing the action as

$$S_A[\Psi,\overline{\Psi}] = \int d^4x\,\overline{\Psi}\,\left[i\not{D}_L(\mathbf{r})\right]\,\Psi\,,$$

where

$$\not{D}_L(\mathbf{r}) = \begin{pmatrix} 0 & \sigma^\mu\partial_\mu \\ \overline{\sigma}^\mu(\partial_\mu + A_\mu) & 0 \end{pmatrix}.$$

4.2.1 Semiclassical approximation

In the full interacting theory one is interested in computing expectation values of fields $\mathcal{O}(x_1,\ldots,x_n)$ that are composite operators made up from the elementary fields ψ, $\overline{\psi}$ and A_μ. In the semiclassical approximation one considers fluctuations, $A_\mu = \overline{A}_\mu + Q_\mu$, around a classical solution $\overline{A}_\mu$; for the fermionic part of the action it is sufficient to restrict oneself to $S_{\overline{A}}[\psi,\overline{\psi}]$. In the spirit of the semiclassical approximation, in the computation of expectation values of operators depending on A_μ as well as on fermionic fields, the functional integration over the latter is performed first and then the result is substituted into (4.14). Thus as a starting point one considers

$$\langle\mathcal{O}[\Psi,\overline{\Psi}]\rangle_A = \int[\mathcal{D}\Psi\mathcal{D}\overline{\Psi}]e^{-S_A[\Psi,\overline{\Psi}]}\mathcal{O}[\Psi,\overline{\Psi}]$$

and evaluates this integral through a saddle point approximation.

According to the properties of Berezin integration (see appendix A) the last integral yields a non-vanishing result only if all the integrations over the fermionic variables are correctly saturated. Expanding Ψ and $\overline{\Psi}$ into eigenfunctions of the Dirac operator

$$i\not{D}_L(\mathbf{r})f^{(n)} = \lambda_n f^{(n)}\,, \qquad\qquad i\not{D}_R(\mathbf{r})g^{(n)} = \lambda_n^* g^{(n)}\,,$$

gives

$$\begin{aligned}\Psi &= \sum_{i=1}^{m} b_i^{(0)} f_i^{(0)} + \sum_n b_n f^{(n)} \\ \overline{\Psi} &= \sum_{j=1}^{\overline{m}} \overline{c}_j^{(0)} g_j^{(0)\dagger} + \sum_n \overline{c}_n g^{(n)\dagger}\,,\end{aligned} \tag{4.18}$$

where b and $\overline{c}$ are Grassmannian c-numbers. Substituting into the action one gets

$$S_A[\Psi,\overline{\Psi}] = \sum_{n\neq 0} \lambda_n \overline{c}_n b_n\,,$$

so that one obtains

$$\langle \mathcal{O} \rangle_A = \int \prod_{i=1}^{m} [db_i^{(0)}] \prod_{j=1}^{\overline{m}} [d\overline{c}_j^{(0)}] \prod_{n\neq 0} [db_n\, d\overline{c}_n]\, e^{-\sum_{n\neq 0}\lambda_n \overline{c}_n b_n}\, \mathcal{O}(b,\overline{c})\,. \tag{4.19}$$

This expression shows that the operator $\mathcal{O}$ must contain exactly the number of zero-modes of the Dirac operator that are present in the instanton background in order to have a non-vanishing vev.

Hence one must study the eigenvalue problem for the Dirac operator in the instanton background. The number, n_L, of left-chirality zero-modes of the Dirac operator (*i.e.* the number of zero-modes of $i\not{D}_L$) and the number of right-chirality zero-modes (*i.e.* the number of zero-modes of $i\not{D}_R$), n_R, are related by an index theorem [99]

$$n_L - n_R = 2 d_{\mathbf{r}} K[\overline{A}]\,. \tag{4.20}$$

Moreover for configurations with self-dual (anti self-dual) field strength one has n_R=0 (n_L=0). Thus from (4.19) in the K-instanton sector it immediately follows that the operator $\mathcal{O}$ must contain an excess of exactly $2Kd_{\mathbf{r}}$ factors of Ψ with respect to $\overline{\Psi}$ to yield a non-zero result.

For instance in the case of a theory containing two Weyl fermions, ψ and χ, in the fundamental representation of the gauge group SU(2) the relation

$$n_L = 2 d_{\mathbf{r}} K[\overline{A}]\,, \qquad n_R = 0 \qquad \text{when} \quad F_{\mu\nu}[A] = \tilde{F}_{\mu\nu}[A] \tag{4.21}$$

gives one single zero-mode, $f^{(0)}$ for ψ and χ in the K=1 sector, so that

$$\begin{aligned}\psi &= a_0 f^{(0)} + \sum_{n\neq 0} a_n f^{(n)} \\ \chi &= b_0 f^{(0)} + \sum_{n\neq 0} b_n f^{(n)}\,,\end{aligned}$$

while $\overline{\psi} = \sum_{n\neq 0} \overline{c}_n g^{(n)\dagger}$ and $\overline{\chi} = \sum_{n\neq 0} \overline{d}_n g^{(n)\dagger}$ do not possess zero-modes. An operator with non-vanishing vev is $\mathcal{O}(x,y) = \psi^\alpha(x)\chi_\alpha(y)$. The saddle point semiclassical approximation to the expectation value of $\mathcal{O}$ is

$$\begin{aligned}\langle\mathcal{O}\rangle_A &= \int [da_0\, db_0] \prod_{n\neq 0} [da_n\, db_n\, d\overline{c}_n\, d\overline{d}_n] e^{-\sum_{n\neq 0}\lambda_n(\overline{c}_n a_n + \overline{d}_n b_n)}\,. \\ &\cdot \left(\sum_{n=0}^{\infty} a_n f^{(n)}\right)\left(\sum_{n=0}^{\infty} b_n f^{(n)}\right) = \left[\det{}'(i\not{D}_L(\mathbf{2}))\right]^2 f^{(0)}(x) f^{(0)}(y)\,,\end{aligned}$$

where $\det'$ denotes the product of the non-zero eigenvalues. The generalization of this elementary example to more complicated cases is rather straightforward. In the following sections computations of vev's of composite operators in $\mathcal{N}$=4 supersymmetric Yang–Mills theory will be described. The $\mathcal{N}$=4 theory contains four Weyl spinors in the adjoint representation. In the case of a SU(2) gauge group equation (4.21) with $d_{\mathbf{3}}$=2 yields sixteen zero-modes in the one-instanton sector.

4.2.2 Fermionic zero-modes

The discussion of the previous subsection shows that in order to compute the instanton contribution to expectation values of fermionic fields one needs to know the zero-value eigenstates of the Dirac operator. In particular their norms are necessary to convert the integrations over the zero-modes into integrations over (Grassmannian) collective coordinates as in the case of the bosonic zero-modes. For a Weyl spinor in the fundamental representation of SU(2) there is one single zero mode which satisfies the equation

$$iD^{(\mathrm{f})}_\mu(\overline{A})\overline{\sigma}^{\mu\,\dot\alpha\alpha}\psi^{(0)}_\alpha = 0\,,$$

where $D^{(\mathrm{f})}_\mu(\overline{A})$ denotes the Dirac operator in the fundamental of SU(2) in the K=1 instanton sector. The solution is

$$\psi^{(0)}_{\alpha,s}(x) = \frac{\varepsilon_{\alpha s}}{[(x-x_0)^2\rho^2]^{3/2}}\,,$$

where s=1,2 is an index of the fundamental of SU(2). The norm of this zero eigenfunction is

$$\|\psi^{(0)}\| = \frac{\pi}{\rho}\,.$$

In the case of the adjoint representation the equation for the zero-modes becomes

$$\left[iD_\mu^{(\mathrm{adj})}(\overline{A})\right]_a{}^b\,\overline{\sigma}^{\mu\,\dot{\alpha}\alpha}\lambda_{\alpha b}^{(0)}=0\,. \tag{4.22}$$

This equation is more complicated and according to the Atiyah–Singer theorem [99] possesses four independent solutions in the case K=1 and G=SU(2). Instead of explicitly solving equation (4.22) the zero eigenfunctions can be obtained acting with the symmetries of the model [3] on the configuration

$$\lambda=\overline{\lambda}=0\,,\qquad A_\mu=\overline{A}_\mu\,,$$

which trivially satisfies (4.22). More precisely the zero-modes are obtained acting by the symmetries broken by the instanton background. Two of the four zero-modes are generated by a supersymmetry transformation

$$\delta_1\lambda=\frac{\sqrt{\rho}}{2}\overline{F}_{\mu\nu}\sigma^{\mu\nu}\eta$$

and are associated with the supersymmetry parameter η. The remaining two eigenfunctions are obtained by a superconformal transformation with parameter $\overline{\xi}$

$$\delta_2\lambda=\frac{1}{2\sqrt{\rho}}\overline{F}_{\mu\nu}\sigma^{\mu\nu}\left[(x-x_0)_\kappa\sigma^\kappa\overline{\xi}\right]\,.$$

The above four zero-modes can be written in the compact form

$$\lambda_i^{(0)}=\frac{1}{2}\overline{F}_{\mu\nu}\sigma^{\mu\nu}\zeta_i(x)\,, \tag{4.23}$$

where

$$\zeta_i^{(0)}(x)=\frac{1}{\sqrt{\rho}}\left[\rho\eta_i+(x-x_0)^\mu\sigma_\mu\xi_{i-2}\right]\,,\qquad i=1,2,3,4 \tag{4.24}$$

and

$$\eta_1=\overline{\xi}_1=\begin{pmatrix}1\\0\end{pmatrix}\,,\qquad \eta_2=\overline{\xi}_2=\begin{pmatrix}0\\1\end{pmatrix}\,,\qquad \eta_{3,4}=\overline{\xi}_{-1,-2}=0\,.$$

The powers of ρ in (4.24) are chosen in such a way as to give to $\zeta_i^{(0)}$ the correct dimension, $[\mathrm{mass}]^{-\frac{1}{2}}$, for a supersymmetry parameter. The zero-modes $\lambda_i^{(0)}$ are orthogonal and their norms are

$$\|\lambda_j^{(0)}\|=\frac{4\sqrt{2}\pi\sqrt{\rho}}{g}\,,\qquad j=1,2\,,\qquad \|\lambda_k^{(0)}\|=\frac{8\pi\sqrt{\rho}}{g}\,,\qquad k=3,4\,.$$

[3] Note that this construction applies to non-supersymmetric cases as well.

The extension of the analysis described here to the case of a SU(N) gauge group, in the K=1 sector, can be obtained without too much effort and is achieved by the embedding of a SU(2) instanton in the SU(N) field configuration, see [100]. The generalization to multi-instanton sectors on the contrary is quite involved and has been developed in [101, 102, 103, 104, 105].

4.3 Instanton calculus in supersymmetric gauge theories

The machinery of the previous sections allows to compute the instanton contribution to the vacuum expectation value of composite operators in supersymmetric gauge theories. The problem is reduced to the computation of the "primed" determinants of the kinetic operators, *i.e.* the product of non-zero eigenvalues, and then to the integration over the collective coordinates with the measure constructed in sections 4.1 and 4.2.

In non-supersymmetric theories the integration over the bosonic collective coordinates is IR-divergent because of the singular behaviour at $\rho \to 0$. In supersymmetric gauge theories, on the contrary, in all the known cases the integration turns out to be finite thanks to the balance of fermionic and bosonic degrees of freedom that controls the ρ-dependence. This exact balance produces another significant simplification: the product of non-zero eigenvalues of bosonic and fermionic kinetic operators exactly cancels out. As a result in various models instanton calculus allows to obtain finite and exact results (in the semiclassical approximation) for the vev's of composite operators, see [87] for a review.

Cancellation of determinants in supersymmetric theories.

Supersymmetry implies relations among the eigenvalues of the kinetic operators for scalars, vectors and spinors, that lead to an exact compensation in the ratio of determinants that enters the expression of the expectation values in the semiclassical approximation.

First notice that the chiral Dirac operator $i\not{D}_L(\mathbf{r})$ satisfies

$$\left|\det\left(i\not{D}_L(\mathbf{r})\right)\right|^2 = \det\left[\left(i\not{D}_L(\mathbf{r}) i\not{D}_R(\mathbf{r})\right)\right] = \det\left(i\not{D}(\mathbf{r})\right)\,\det(-\Box)\,,$$

where $i\not{D}(\mathbf{r}) = i\gamma^\mu D_\mu(\mathbf{r})$ is the kinetic operator for a Dirac spinor, so it is sufficient

to solve the eigenvalue problem for $i\not{D}(\mathbf{r})$. Let λ_n be the non zero solutions of the eigenvalue problem for a scalar field in the instanton background

$$D^2(\overline{A})\varphi^{(n)} = -\lambda_n^2\varphi^{(n)} \qquad\qquad \lambda_n \neq 0\,.$$

The two eigenvalue problems to be solved are

$$\begin{aligned} &i\not{D}(\overline{A})\Psi^{(n)} = \mu_n\Psi^{(n)} \\ &\left(-D^2(\overline{A})\delta_{\mu\nu} - 2F_{\mu\nu}[\overline{A}]\right)Q_\nu^{(n)} = \rho_n^2 Q_\mu^{(n)} \end{aligned}$$

and one can prove the following relations

$$\mu_n = \lambda_n \qquad \longleftrightarrow \qquad \Psi^{(n)}(x) = \begin{pmatrix} \xi^{(n)} \\ \overline{\chi}^{(n)} \end{pmatrix}\,,$$

with $\xi^{(n)} = \frac{i}{\lambda_n}D_\mu(\overline{A})\varphi^{(n)}\sigma^\mu\overline{\epsilon}$, $\overline{\chi}^{(n)} = \overline{\epsilon}\varphi^{(n)}$ and

$$\rho_n = \lambda_n \qquad \longleftrightarrow \qquad Q_\mu^{(n)} = \overline{\eta\sigma}_\mu\xi^{(n)}\,,$$

where $\overline{\eta}$ and $\overline{\epsilon}$ are two constant Weyl spinors.

The ratio of determinants that enters the expression for the vev of an operator in the semiclassical approximation is

$$\left[\frac{\det(M_{\mu\nu}^{\text{g.f.}}+\mu^2)}{\det' M_{\mu\nu}^{\text{g.f.}}}\right]^{\frac{1}{2}} \frac{\det D^2(\overline{A})}{\det(D^2(\overline{A})+\mu^2)} \frac{\det'\left(i\not{D}_L^{(\text{adj})}\right)}{\det\left(i\not{D}_L^{(\text{adj})}+i\mu\right)}\,,$$

where a regularization *à la* Pauli–Villars has been introduced. Now substituting the eigenvalues found in the previous sections one can show that there is a an exact cancellation between bosons and fermions, so that the contribution of the non-zero modes is 1! The total contribution of the zero-modes is

$$\mu^{n_B-\frac{1}{2}n_F}\,, \tag{4.25}$$

where n_B and n_F are respectively the number of bosonic and fermionic zero-modes. Notice that in the case of $\mathcal{N}$=4 super Yang–Mills theory (4.25) reduces to 1, as is expected, since the theory, being finite, cannot lead to a result depending on the renormalization scale.

For the calculation of vev's of operators $\mathcal{O}$ there are two different situations to be considered. If the operator $\mathcal{O}$ contains precisely the number of zero-modes to

saturate the fermionic integrations in the instanton background, one must simply substitute the fermionic fields in $\mathcal{O}$ with the zero-modes and perform the integration over the collective coordinates. This is the case that will be discussed in section 4.6. If instead the operator $\mathcal{O}$ does not contain a sufficient number of fermionic insertions to saturate the Grassmannian integrals, then the correlator is zero at the lowest order and one must consider in $\langle\mathcal{O}\rangle$ the insertion of terms lowered from the action, until the correct number of spinors is obtained. This case will be described in section 4.5 for the correlation function of four scalar fields.

4.4 Instanton calculations in $\mathcal{N}$=4 supersymmetric Yang–Mills theory

The formalism described in the preceding sections will now be applied to the computation of Green functions of gauge invariant composite operators in $\mathcal{N}$=4 supersymmetric Yang–Mills theory. In particular correlation functions of operators belonging to the multiplet of currents, see equations (2.17) and (2.18), will be calculated to lowest order in the one-instanton sector and with gauge group SU(2) [89]. The extension of the results presented here to the case of SU(N) and to the K-instanton sector in the large N limit was given in [90] and [91, 92] and will be briefly reviewed in the concluding section. The operators in the current multiplet that will be considered play a central rôle in the correspondence with type IIB superstring theory compactified on $\mathrm{AdS}_5 \times S^5$. This issue will be discussed in chapter 5, where the instanton calculations will be compared with D-instanton effects in the type IIB superstring on $\mathrm{AdS}_5 \times S^5$.

In [89] a closed form for the instanton contribution to a four-point function of composite scalar operators, $\mathcal{Q}^{ij}$ in the **20** of the SU(4) R-symmetry group has been given. In the next section the complete space-time dependence of this Green function will be computed. As will be discussed it will prove useful to employ for this calculation a description of the model in terms of $\mathcal{N}$=2 multiplets. Also, it will be shown that when a limit of coincident points is considered, in order to study the operator product expansion (OPE), the function develops a logarithmic singularity. Analogous results have been found by various groups [106, 107, 108, 109]. In these papers a similar behaviour was extracted without explicitly performing all the integrations, on the contrary in the computation presented here the singular behaviour is shown with no ambiguities after a complete evaluation of the integrals which yield

the Green function under consideration in the semiclassical approximation. A more detailed analysis of this subject and related questions will be presented elsewhere [110].

In section 4.6 the computation of correlation functions of sixteen fermionic currents $\hat{\Lambda}^A_\alpha$ and of eight gaugino bilinears $\mathcal{E}^{AB}$ will be reported. For these Green functions the integration over the collective coordinates will not be performed, nevertheless the unintegrated form presented here will be sufficient for the comparison with the results from type IIB superstring theory discussed in the final chapter.

4.5 A correlation function of four scalar supercurrents

The multiplet of superconformal currents of $\mathcal{N}$=4 super Yang–Mills theory has been given in section 2.2 for the Abelian case. The natural four-point function of superconformal currents to consider would be a correlator of four stress energy tensors. However, due to its complicated tensorial structure even the free-field expression for this correlator is awkward to express compactly and it is much simpler to consider correlations of four gauge-invariant composite scalar operators

$$\mathcal{Q}^{ij} = \varphi^i\varphi^j - \frac{1}{6}\delta^{ij}\varphi_k\varphi^k\,, \qquad i,j,k = 1,2,\ldots,6\,, \tag{4.26}$$

belonging to the representation **20** of the SU(4) R-symmetry group. As discussed in chapter 2 $\mathcal{Q}^{ij}$ is the lowest component of the composite twisted chiral current superfield [111] $\mathcal{W}^{ij}_{(2)}$

$$\mathcal{W}^{ij}_{(2)} = \mathrm{tr}\left(W^iW^j - \frac{\delta^{ij}}{6}W_kW^k\right)\,.$$

After calculating correlation functions of these components one can derive those of any other component in the $\mathcal{N} = 4$ supercurrent multiplet by making use of the superconformal symmetry [111]. A way to explicitly do it may be to resort to analytic superspace, recalling that

$$\mathcal{Q}^{ij}(x) \sim \mathcal{W}^{ij}_{(2)}(\Upsilon)\Big|_{\theta=\bar\theta=0}\,,$$

where Υ are the supercoordinates of analytic superspace [21]. Therefore, by computing correlators of $\mathcal{Q}^{ij}(x)$ and substituting $x \to \Upsilon$ any of the other correlation

functions can in principle be obtained by expanding in the fermionic as well as in the auxiliary bosonic coordinates of analytic superspace.

The correlation function that will be considered is therefore

$$\langle \mathcal{Q}^{i_1 j_1}(x_1)\mathcal{Q}^{i_2 j_2}(x_2)\mathcal{Q}^{i_3 j_3}(x_3)\mathcal{Q}^{i_4 j_4}(x_4)\rangle \,. \tag{4.27}$$

The value of this correlation function in the free field theory is determined from the expression for the free two-point scalar Green function which is

$$\langle \varphi^{ia}(x)\varphi^{jb}(y)\rangle_{\text{free}} = \frac{1}{(2\pi)^2}\frac{\delta^{ij}\delta^{ab}}{(x-y)^2}\,. \tag{4.28}$$

Hence, the free-field expression for the correlation function that follows by Wick contractions is

$$\begin{aligned}&\langle \mathcal{Q}^{i_1 j_1}(x_1)\mathcal{Q}^{i_2 j_2}(x_2)\mathcal{Q}^{i_3 j_3}(x_3)\mathcal{Q}^{i_4 j_4}(x_4)\rangle_{\text{free}} \qquad (4.29)\\ &= \frac{1}{(4\pi^2)^4}\left[N^4\frac{\delta^{i_1 i_3}\delta^{j_1 j_3}\delta^{i_2 i_4}\delta^{j_2 j_4}}{x_{13}^4 x_{24}^4} + N^2\frac{\delta^{j_4 i_1}\delta^{j_1 i_3}\delta^{j_3 i_2}\delta^{j_2 i_4}}{x_{41}^2 x_{13}^2 x_{32}^2 x_{24}^2} + \text{permutations}\right],\end{aligned}$$

where,

$$x_{ij} = x_i - x_j\,.$$

The first term in this expression is simply the product of two two-point functions and is known to be exact. The second term, which is the free-field contribution to the connected four-point function, certainly gets corrections from interactions.

As has been discussed in detail in the preceding sections, in $\mathcal{N}$=4 Yang–Mills theory there are sixteen zero-modes in the one-instanton sector in the case at hand of a SU(2) gauge group. In the background of the one-instanton solution, (4.9), (4.10), the correlation function

$$\langle \mathcal{Q}^{i_1 j_1}(x_1)\mathcal{Q}^{i_2 j_2}(x_2)\mathcal{Q}^{i_3 j_3}(x_3)\mathcal{Q}^{i_4 j_4}(x_4)\rangle_{K=1} \tag{4.30}$$

where the subscript $K = 1$ denotes the winding number of the background, is zero at the lowest order, because it does not contain any fermionic insertion to saturate the integrations over the Grassmannian collective coordinates of the sixteen zero-modes. However it is easy to check that the correlation function (4.30) soaks up these sixteen gaugino zero-modes when the perturbative corrections to the standard instanton configuration are considered. This can be seen either from the form of the

supersymmetry transformations, see equation (2.6), or from the equation of motion for the scalar fields φ^i. The expression (4.10) is annihilated by the conserved supersymmetry transformations (those associated with the parameter $\overline{\eta}^{\dot{\alpha}}{}_A$ in (2.6)), while the transformations corresponding to supersymmetries associated with η^A_α are broken and generate eight of the sixteen fermionic zero modes. The other eight zero-modes are generated by superconformal transformations. As a consequence acting with the broken supersymmetry and superconformal transformations on the scalar fields in (4.30) produces a configuration which possesses the required sixteen zero-modes. To obtain the first non-vanishing contribution to (4.30) one must consider insertions of terms lowered from the action until the correct number of fermionic fields is obtained. The first non vanishing correction comes from the insertion of a Yukawa term for each field φ^i (see the form (2.5) of the action)

$$\langle \mathcal{Q}^{i_1 j_1}(x_1)\mathcal{Q}^{i_2 j_2}(x_2)\mathcal{Q}^{i_3 j_3}(x_3)\mathcal{Q}^{i_4 j_4}(x_4)\left(\int d^4 z_1 \left[\frac{i}{4} g f^{abc}\overline{t}^i_{AB}\left(\lambda_a^{\alpha A}\lambda^B_{\alpha b}\varphi_{ic}\right)(z_1)\right]\right)\cdot$$
$$\ldots\cdot\left(\int d^4 z_8 \left[\frac{i}{4} g f^{abc}\overline{t}^i_{AB}\left(\lambda_a^{\alpha A}\lambda^B_{\alpha b}\varphi_{ic}\right)(z_8)\right]\right)\rangle_K\,. \tag{4.31}$$

Wick contractions among the scalar fields produce propagators, so that (4.31) becomes schematically

$$\langle\left(\int d^4 z_1\left[g\Delta(x_1-z_1)\,(\lambda\lambda)\,(z_1)\right]\right)\ldots\left(\int d^4 z_8\left[g\Delta(x_4-z_8)\,(\lambda\lambda)\,(z_8)\right]\right)\rangle, \tag{4.32}$$

where for simplicity of notation not all the indices have been indicated explicitly. Computing the integrations in (4.32) amounts to substitute each factor with the solution, $\varphi_i^{(0)}$, of an equation of the form

$$\left[D^2(\overline{A})\varphi^i\right](x) = J^i(x)\,, \tag{4.33}$$

with 'source' $J^i(x) = \overline{t}^i_{AB}\left(\lambda^A\lambda^B\right)(x)$.

In order to evaluate this instanton contribution in the most convenient fashion it will be convenient to use the $\mathcal{N}$=2 supersymmetric description in which, as explained in section (2.1), φ^i decomposes into the complex $\mathcal{N}$=2 singlet φ with U(1) charge +2 and a neutral $\mathcal{N}$=2 'quaternion' $q_{_T}$ in the $(\mathbf{2},\mathbf{2})_0$ representation of $\mathrm{SU}(2)_{\mathcal{V}}\times\ \mathrm{SU}(2)_{\mathcal{H}}$ which resides in the $\mathcal{N}$=2 hypermultiplet. The scalar component $\mathcal{Q}^{AB}{}_{CD}$ of the $\mathcal{N}$=4 current decomposes in the following way in terms of $\mathcal{N}$=2 fields,

$$\begin{aligned}
\mathcal{Q}^{S0} &= q^S\varphi \qquad \mathcal{Q}^{S0\,\dagger} = q^S\overline{\varphi} \qquad \mathcal{Q}^{ST} = q^S q^T - \text{trace} \\
\mathcal{Q}_{(+)} &= \varphi^2 \qquad \mathcal{Q}_{(-)} = \overline{\varphi}^2 \qquad \mathcal{Q}_{(0)} = \overline{\varphi}\varphi - \ \text{trace}\ ,
\end{aligned} \tag{4.34}$$

where the notations are those of section (2.1) and $q_{u\dot{u}} = q_S \sigma^S{}_{u\dot{u}}$ and $\sigma^S = (\mathbb{1}, \vec{\sigma})$, with $\vec{\sigma}$ the standard Pauli matrices (and $S, T = 1, \ldots, 4$ are $\mathrm{SU}(2)_{\mathcal{V}} \times \mathrm{SU}(2)_{\mathcal{H}}$ vector indices). The correlation function of two φ^2 and two $\overline{\varphi}^2$ will be discussed here. From (4.30) the free-field expression for this particular correlator is

$$\begin{aligned}
&\langle \varphi^2(x_1)\varphi^2(x_2)\overline{\varphi}^2(x_3)\overline{\varphi}^2(x_4)\rangle_{\text{free}} = \\
&= \frac{1}{(4\pi^2)^4}\left(\frac{4N^4}{x_{13}^4 x_{24}^4} + \frac{4N^4}{x_{14}^4 x_{23}^4} + \frac{16N^2}{x_{41}^2 x_{13}^2 x_{32}^2 x_{24}^2}\right) ,
\end{aligned} \tag{4.35}$$

where in the previous notation $\varphi^2 = \mathcal{Q}_{(+)} \equiv \mathcal{Q}^{55} - \mathcal{Q}^{66} + 2i\mathcal{Q}^{56}$.

The $\mathcal{N}$=2 formalism is particularly suitable to perform the integrations over the fermionic zero-modes because it follows from the structure of the Yukawa couplings (2.13) that φ only absorbs those zero modes of the $\mathcal{N}$=4 gauginos that belong to the $\mathcal{N}$=2 vector multiplet (λ^u_α), while $\overline{\varphi}$ absorbs the zero modes belonging to the $\mathcal{N}$=2 hypermultiplet ($\psi^{\dot{u}}_\alpha$). The expressions for the zero-modes that will be used are those suggested by the supersymmetry transformations of the instanton

$$\lambda^u_{(0)\alpha} = \frac{1}{2}\overline{F}_{\mu\nu}\sigma^{\mu\nu\,\beta}_{\alpha}\frac{1}{\sqrt{\rho_0}}\left(\rho_0\eta^u_\beta + (x - x_0)_\kappa \sigma^\kappa_{\beta\dot{\beta}}\bar{\xi}^{\dot{\beta}u}\right) . \tag{4.36}$$

and similarly for $\psi^{\dot{u}}_{(0)\alpha}$. Here and in the following the 'size' of the instanton will be denoted by ρ_0. The reason for this choice of notation will become clear when the results of this chapter will be reanalyzed in the context of the AdS/CFT correspondence in chapter 5. In this decomposition η_β and $x_{\beta\dot{\beta}}\overline{\xi}^{\dot{\beta}}$ are the parameters of the broken supersymmetry and special supersymmetry transformations respectively. One can assemble the fermionic collective coordinates into the spinors $\zeta^u_\pm(\rho_0, x)$ (where $\pm$ refers to the U(1) R-symmetry charge) to parameterize the fermionic zero modes

$$\zeta^u_{\pm\alpha}(\rho_0, x - x_0) = \frac{1}{\sqrt{\rho_0}}\left(\rho_0\eta^u_{\pm\alpha} + \sigma^\mu{}_{\alpha\dot{\alpha}}(x_\mu - x_{0\,\mu})\overline{\xi}^{\dot{\alpha}\,u}_{\pm}\right) , \tag{4.37}$$

Using the bosonic measure for SU(2) instantons derived in section 4.1 and the norms of the fermionic zero-modes given in section 4.2 and proceeding as described after equation (4.30) one finds that in the semi-classical approximation the one-instanton

contribution to the four bosonic current Green function is [4]

$$G_{\mathcal{Q}^4}(x_p) = \langle g^2\varphi^2(x_1) g^2\varphi^2(x_2) g^2\overline{\varphi}^2(x_3) g^2\overline{\varphi}^2(x_4)\rangle_{K=1} = \frac{g^8}{2^{32}\pi^{10}} e^{-\frac{8\pi^2}{g^2}+i\theta}$$
$$\int \frac{d\rho_0 d^4x_0}{\rho_0^5} d^4\eta_+ d^4\overline{\xi}_+ d^4\eta_- d^4\overline{\xi}_-\ \varphi^2_{(0)}(x_1)\varphi^2_{(0)}(x_2)\overline{\varphi}^2_{(0)}(x_3)\overline{\varphi}^2_{(0)}(x_4)\,, \qquad (4.38)$$

where (+) (or (−)) refers to the U(1) charges of the gauginos in the vector (or hyper) multiplet (see below). In equation (4.38) the explicit dependence on the vacuum angle, θ, has been indicated. The Green function $G_{\mathcal{Q}^4}$ receives a contribution also from the $K=-1$ sector that is the complex conjugate of (4.38).

In (4.38) the fields φ and $\overline{\varphi}$ have been replaced by the expressions

$$\varphi(x) \to \varphi_{(0)}(x) = \frac{1}{2\sqrt{2}}\varepsilon_{uv}\zeta_+^u \sigma^{\mu\nu}\zeta_+^v \overline{F}_{\mu\nu}$$
$$\overline{\varphi}(x) \to \overline{\varphi}_{(0)}(x) = \frac{1}{2\sqrt{2}}\varepsilon_{\dot{u}\dot{v}}\zeta_-^{\dot{u}} \sigma^{\mu\nu}\zeta_-^{\dot{v}} \overline{F}_{\mu\nu}\,, \qquad (4.39)$$

which satisfy an equation of the form of (4.33) and are the leading nonvanishing terms that result from Wick contractions in which Yukawa couplings are lowered from the exponential of the action until a sufficient number of fermion fields are present to saturate the fermionic integrals. Of course, these expressions can also be obtained directly from the supersymmetry transformations (2.6) by acting twice on $\overline{F}_{\mu\nu}$ with the broken supersymmetry generators. After some elementary Fierz transformations on the fermionic collective coordinates the fermionic integrations can be performed in a standard manner and the result is

$$G_{\mathcal{Q}^4}(x_p) = \frac{3^4}{4\pi^{10}}\, g^8\, e^{-\frac{8\pi^2}{g^2}+i\theta} \int \frac{d\rho_0 d^4x_0}{\rho_0^5}\, x_{12}^4 x_{34}^4 \prod_{p=1}^4 \left[\frac{\rho_0}{\rho_0^2+(x_p-x_0)^2}\right]^4. \qquad (4.40)$$

The integration in (4.40) resembles that of a standard Feynman diagram with momenta replaced by position differences and can be performed by introducing the Feynman parameterization,

$$G_{\mathcal{Q}^4}(x_p) = \frac{3^4\Gamma(16)}{4\pi^{10}\left(\Gamma(4)\right)^4}\, g^8\, e^{-\frac{8\pi^2}{g^2}+i\theta} \int \prod_p \alpha_p^3 d\alpha_p\, \delta\left(1-\sum_q \alpha_q\right) \cdot$$
$$\cdot \int \frac{d\rho_0 d^4x_0}{\rho_0^5} \frac{x_{12}^4 x_{34}^4\, \rho_0^{16}}{\left(\rho_0^2 + x_0^2 - 2x_0\cdot\sum_p \alpha_p x_p + \sum_p x_p^2\right)^{16}}\,. \qquad (4.41)$$

[4] Here and in the following computations of correlators of current bilinears in the fundamental Yang–Mills fields a factor of g^2 is included for each external insertion. It would be equivalent to rescale the fundamental Yang–Mills fields according to $A \to A' = gA$ and compute the Green functions of primed fields. This leads to a common overall dependence on g^2 for the three correlation functions that will be considered.

The five-dimensional integral yields

$$\begin{aligned} G_{\mathcal{Q}^4}(x_p) &= \frac{3^3\,\Gamma(11)}{2^7\,(\pi^3\Gamma(4))^4}\,g^8 e^{-\frac{8\pi^2}{g^2}+i\theta}\cdot \\ &\cdot\int\prod_p \alpha_p^3 d\alpha_p\,\delta\left(1-\sum\nolimits_q \alpha_q\right)\frac{x_{12}^4 x_{34}^4}{\left(\sum_p \alpha_p\alpha_q x_{pq}^2\right)^8}\,. \end{aligned} \tag{4.42}$$

This integral can be simplified by observing that it is essentially obtained by acting with derivatives on the box-integral with four massless external particles,

$$G_{\mathcal{Q}^4}(x_p) = \frac{3^3\,\Gamma(11)}{2^7\,(\pi^3\Gamma(4))^4}\,g^8 e^{-\frac{8\pi^2}{g^2}+i\theta}\,x_{12}^4 x_{34}^4\prod_{p<q}\frac{\partial}{\partial x_{pq}^2}B(x_{pq})\,, \tag{4.43}$$

where the box integral is

$$B(x_{pq}) = \int\prod_p d\alpha_p\,\delta\left(1-\sum_q \alpha_q\right)\frac{1}{\left(\sum_p \alpha_p\alpha_q x_{pq}^2\right)^2}\,. \tag{4.44}$$

The result, correcting a sign error in [89], may be expressed as a combination of logarithms and dilogarithms [112]

$$\begin{aligned} B(x_{pq}) &= \frac{4}{\sqrt{\Delta(x_{pq})}}\left[\frac{1}{2}\log\left(\frac{u_+u_-}{(1-u_+)^2(1-u_-)^2}\right)\log\left(\frac{u_+}{u_-}\right)+\right. \\ &\left.-\mathrm{Li}_2(1-u_+)+\mathrm{Li}_2(1-u_-)-\mathrm{Li}_2\left(1-\frac{1}{u_-}\right)+\mathrm{Li}_2\left(1-\frac{1}{u_+}\right)\right]\,, \end{aligned} \tag{4.45}$$

where

$$\Delta = \det_{4\times 4}((x_{pq}^2)) = X^2+Y^2+Z^2-2XY-2YZ-2ZX \tag{4.46}$$

and

$$u_\pm = \frac{Y+X-Z\pm\sqrt{\Delta}}{2Y}\,, \tag{4.47}$$

with $X = x_{12}^2x_{34}^2$, $Y = x_{13}^2x_{24}^2$ and $Z = x_{14}^2x_{23}^2$.

Notice that up to an overall dimensional factor needed for the correct scaling, $G_{\mathcal{Q}^4}$ turns out to be a function of the two independent superconformally invariant cross ratios X/Z and Y/Z. Although not immediately apparent, the expression $B(x_{pq})$ is symmetric under any permutation of the external legs, as can be seen by making use of the properties of the dilogarithms,

$$\begin{aligned} \mathrm{Li}_2(z)+\mathrm{Li}_2(1-z) &= \frac{\pi^2}{6}-\log(z)\log(1-z) \\ \mathrm{Li}_2(z)+\mathrm{Li}_2\left(\frac{1}{z}\right) &= -\frac{\pi^2}{6}-\frac{1}{2}[\log(-z)]^2\,, \end{aligned} \tag{4.48}$$

and observing that the relevant permutations correspond to permutations of $X = Yu_+u_-$, Y and $Z = Y(1-u_+)(1-u_-)$, that are generated by the two transformations: a) $u_+ \to 1/u_-$, $u_- \to 1/u_+$, $Y \to Yu_-u_+$, which is equivalent to the exchange of X and Y, leaving Z fixed (*i.e.* to the exchange of x_1 and x_4 or, equivalently, of x_2 and x_3) and b) $u_+ \to 1-u_-$, $u_- \to 1-u_+$ at fixed Y, which is equivalent to the exchange of X and Z (or the exchange of x_2 and x_4 or, equivalently, of x_1 and x_3).

To compute the explicit dependence of $G_{\mathcal{Q}^4}$ on x_p one must perform the six derivatives with respect to x_{pq}. These can be more conveniently calculated rewriting the differential operator in (4.43) as

$$\prod_{p<q} \frac{\partial}{\partial x_{pq}^2} = \mathcal{D}_Z \mathcal{D}_Y \mathcal{D}_X \,, \tag{4.49}$$

where

$$\mathcal{D}_T = \left(\frac{\partial}{\partial T} + T \frac{\partial^2}{\partial T^2} \right) , \qquad T = X, Y, Z \,.$$

Defining

$$Q(x_{pq}) = \prod_{p<q} \frac{\partial}{\partial x_{pq}^2} B(x_{pq})$$

after a lengthy computation one obtains

$$\begin{aligned}
Q(x_{pq}) = \frac{2}{\Delta^5}\Big(& 193X^6 - 114X^5Y - 1281X^4Y^2 + 2404X^3Y^3 - 1281X^2Y^4 + \\
& -114XY^5 + 193Y^6 - 114X^5Z + 4734X^4YZ - 4620X^3Y^2Z - 4620X^2Y^3Z + \\
& +4734XY^4Z - 114Y^5Z - 1281X^4Z^2 - 4620X^3YZ^2 + 15402X^2Y^2Z^2 + \\
& -4620XY^3Z^2 - 1281Y^4Z^2 + 2404X^3Z^3 - 4620X^2YZ^3 - 4620XY^2Z^3 + \\
& +2404Y^3Z^3 - 1281X^2Z^4 + 4734XYZ^4 - 1281Y^2Z^4 - 114XZ^5 - 114YZ^5 + \\
& +193Z^6\Big) + \frac{6}{\Delta^6}\Big\{\Big(- 33X^8 - 96X^7Y + 714X^6Y^2 - 1008X^5Y^3 + 1008X^3Y^5 + \\
& -714X^2Y^6 + 96XY^7 + 33Y^8 - 6X^7Z - 1770X^6YZ + 198X^5Y^2Z + 8298X^4Y^3Z + \\
& -8298X^3Y^4Z - 198X^2Y^5Z + 1770XY^6Z + 6Y^7Z + 435X^6Z^2 + 3312X^5YZ^2 + \\
& -13473X^4Y^2Z^2 + 13473X^2Y^4Z^2 - 3312XY^5Z^2 - 435Y^6Z^2 - 930X^5Z^3 + \\
& +1914X^4YZ^3 + 15396X^3Y^2Z^3 - 15396X^2Y^3Z^3 - 1914XY^4Z^3 + 930Y^5Z^3 + \\
& +645X^4Z^4 - 6384X^3YZ^4 + 6384XY^3Z^4 - 645Y^4Z^4 + 78X^3Z^5 + 3114X^2YZ^5 + \\
& -3114XY^2Z^5 - 78Y^3Z^5 - 279X^2Z^6 + 279Y^2Z^6 + 90XZ^7 - 90YZ^7\Big) \log\left[\frac{X}{Y}\right] +
\end{aligned}$$

$$
\begin{aligned}
&-\Big(11X^8-28X^7Y-52X^6Y^2+284X^5Y^3-430X^4Y^4+284X^3Y^5-52X^2Y^6+\\
&-28XY^7+11Y^8+62X^7Z+590X^6YZ-2142X^5Y^2Z+1490X^4Y^3Z+\\
&+1490X^3Y^4Z-2142X^2Y^5Z+590XY^6Z+62Y^7Z-331X^6Z^2+972X^5YZ^2+\\
&+4491X^4Y^2Z^2-10264X^3Y^3Z^2+4491X^2Y^4Z^2+972XY^5Z^2-331Y^6Z^2+\\
&+362X^5Z^3-4894X^4YZ^3+5132X^3Y^2Z^3+5132X^2Y^3Z^3-4894XY^4Z^3+\\
&+362Y^5Z^3+215X^4Z^4+3404X^3YZ^4-8982X^2Y^2Z^4+3404XY^3Z^4+\\
&+215Y^4Z^4-646X^3Z^5+1170X^2YZ^5+1170XY^2Z^5-646Y^3Z^5+383X^2Z^6+\\
&-1180XYZ^6+383Y^2Z^6-34XZ^7-34YZ^7-22Z^8\Big)\log\left[\frac{XY}{Z^2}\right]+\\
&+\frac{36}{\Delta^{13/2}}\Bigg\{\Big(X^9+3X^8(Y+Z)-6X^7\big(5Y^2-19YZ+5Z^2\big)+(Y-Z)^6\big(Y^3+9Y^2Z+\\
&+9YZ^2+Z^3\big)+X^6\big(62Y^3-144Y^2Z-144YZ^2+62Z^3\big)+3X(Y-Z)^4\cdot\\
&\cdot\big(Y^4+42Y^3Z+114Y^2Z^2+42YZ^3+Z^4\big)-6X^5\big(6Y^4+83Y^3Z-252Y^2Z^2+\\
&+83YZ^3+6Z^4\big)-6X^2(Y-Z)^2\big(5Y^5+34Y^4Z-189Y^3Z^2-189Y^2Z^3+34YZ^4+\\
&+5Z^5\big)-6X^4\big(6Y^5-175Y^4Z+223Y^3Z^2+223Y^2Z^3-175YZ^4+6Z^5\big)+\\
&+X^3\big(62Y^6-498Y^5Z-1338Y^4Z^2+3948Y^3Z^3-1338Y^2Z^4-498YZ^5+62Z^6\big)\Big)\cdot\\
&\cdot\Bigg[\frac{1}{2}\log\left(\frac{XY}{Z^2}\right)\log\left(1+\frac{(X+Y-Z)\sqrt{\Delta}+\Delta}{2XY}\right)-\mathrm{Li}_2\left(\frac{\mathrm{X-Y+Z-\sqrt{\Delta}}}{2\mathrm{X}}\right)+\\
&-\mathrm{Li}_2\left(\frac{\mathrm{-X+Y+Z-\sqrt{\Delta}}}{2\mathrm{Y}}\right)+\mathrm{Li}_2\left(\frac{\mathrm{X-Y+Z+\sqrt{\Delta}}}{2\mathrm{X}}\right)+\mathrm{Li}_2\left(\frac{\mathrm{-X+Y+Z+\sqrt{\Delta}}}{2\mathrm{Y}}\right)\Bigg]\Bigg\}.
\end{aligned}
$$

In conclusion the one-instanton contribution to the correlation function $G_{\mathcal{Q}^4}(x_p)$ is

$$G_{\mathcal{Q}^4}(x_p)=\frac{3^3\,\Gamma(11)}{2^5\,(\pi^3\Gamma(4))^4}\,g^8e^{-\frac{8\pi^2}{g^2}+i\theta}\,x_{12}^4x_{34}^4\,Q(x_{pq})\,. \tag{4.50}$$

Unlike correlation functions of elementary fields that are infra-red problematic and gauge-dependent, the above correlator is well defined at non-coincident points. Moreover it possesses the correct symmetry properties. This can be derived from the symmetry of $B(x_{pq})$ discussed above and the observation that the differential operator (4.49) is completely symmetric. Of course the symmetry properties can be checked directly on the final expression for $Q(x_{pq})$. Notice that the above form for $Q(x_{pq})$ is valid only for X,Y,Z in the 'physical domain', *i.e.* in the region obtainable for allowed choices of x_{pq}^2, that is defined by the condition

$$\Delta=\Delta(X,Y,Z)=X^2+Y^2+Z^2-2XY-2XZ-2YZ\leq 0\,.$$

In this region single terms in the above expression are complex, but the complete function is real. To describe the physical region it is convenient to rewrite $Q(x_{pq})$ in terms of two independent cross ratios,

$$r = \frac{X}{Y} = \frac{x_{12}^2 x_{34}^2}{x_{13}^2 x_{24}^2}\,, \qquad\qquad s = \frac{Z}{Y} = \frac{x_{14}^2 x_{23}^2}{x_{13}^2 x_{24}^2}\,,$$

extracting a factor $\frac{1}{Y^4}$ in $Q(x_{pq})$. Then the physical domain corresponds to the region

$$\delta(r,s) = \frac{\Delta}{Y^2} = 1 + r^2 + s^2 - 2r - 2s - 2rs \leq 0$$

in the (r, s) plane. This is the region inside the parabola in figure 4.1



Figure 4.1: Physical domain for $B(x_{pq})$

Having obtained a closed form for $G_{\mathcal{Q}^4}(x_p)$ one can study the behaviour when any two points are taken close to one another, $x_{pq} \to 0$. For instance $x_{12} \to 0$ corresponds to $r \to 0$, $s \to 1$ simultaneously. In this limit $B(x_{pq})$ develops a logarithmic singularity.

The logarithmic divergence in the Green function $\mathcal{G}_{\mathcal{Q}^4}$ can be shown more clearly considering the particular configuration in which the points x_p, $p = 1, 2, 3, 4$, are taken on a line. In this case the cross ratios r and s are not independent. Assuming $x_1 < x_2 < x_3 < x_4$ r and s are related by

$$\sqrt{r} + \sqrt{s} = 1\,.$$

Thus one can express the result in terms of a single variable η defined by

$$r = \eta^2 \,, \qquad s = (1-\eta)^2 \,. \tag{4.51}$$

In this limit the above computed function $Q(x_{pq})$ reduces to [5]

$$\begin{aligned}
\tilde{Q} = \tilde{Q}(\eta) = 3\,\frac{\log(\eta^2)}{(-1+\eta)^7}(100\,\eta^6 + 429 + 2431\,\eta^2 - 2717\,\eta^3 + 1794\,\eta^4 + \\
-650\,\eta^5 - 1287\,\eta) \; - 3\,\frac{\log[(-1+\eta)^2]}{\eta^7}\,(100\,\eta^6 + 50\,\eta^5 + 44\,\eta^4 + 41\,\eta^3 + \\
+44\,\eta^2 + 50\,\eta + 100) - 2\,\frac{(\eta^2-\eta+1)^2}{\eta^6\,(-1+\eta)^6}(300 - 900\,\eta + 307\,\eta^2 + 886\,\eta^3 + \\
+307\,\eta^4 - 900\,\eta^5 + 300\,\eta^6) \,.
\end{aligned} \tag{4.52}$$

and it can be shown that $\tilde{Q}(\eta)$ has a logarithmic singularity as $\eta \to 0$ or $\eta \to 1$.

This peculiar behaviour has been observed by various authors in four-point functions computed in the context of the AdS/CFT correspondence [106, 107, 108, 109], but here it is derived from an exact field theoretical computation and thus shown with no ambiguities. Notice that single terms in $Q(x_{pq})$ have pole-type singularities in the limit $x_{12} \to 0$, but in the sum the singularity is only logarithmic. Note in particular that the poles associated with operators of dimension lower or equal to $\mathcal{Q}^2$ in the OPE are absent. Moreover with non-vanishing Y the only singularities in $B(x_{pq})$ correspond to the points $(r = 0, s = 1)$ and $(r = 1, s = 0)$ marked in figure 4.1. Taking into account the pre-factor, $x_{12}^4 x_{34}^4$, in the complete expression of $G_{\mathcal{Q}^4}$ one finds that the limit in which two φ^2 operators are taken to coincide gives a vanishing result, while a logarithmic singularity is actually present when φ^2 approaches $\overline{\varphi}^2$.

The logarithms by themselves do not violate the superconformal symmetry of the theory, that is manifest in the final expression of the four-point function. In some respect, logarithmic behaviours in four-point functions should not sound unexpected in a superconformal theory, such as $\mathcal{N}$=4 supersymmetric Yang–Mills theory, that contains a large number of primary fields all of whose (protected) dimensions are integer. The singularity may be due to the presence of an infinite tower of stable BPS dyons that collapse to vanishing mass and size in the superconformal phase or to a gas of instantons of small size from the Euclidean viewpoint. This problem is under active investigation [110].

[5] This expression has been obtained by Yassen Stanev starting from a different but equivalent formula for $Q(x_{pq})$.

4.6 Eight- and sixteen-point correlation functions of current bilinears

In this section the one-instanton contribution to correlation functions of sixteen fermionic bilinears $\hat{\Lambda}^A_\alpha$ and of eight gaugino bilinears $\mathcal{E}^{AB}$ in the current supermultiplet will be calculated in the semiclassical approximation. The expressions obtained will be reconsidered in the next chapter and a comparison will be made with D-instanton effects in type IIB superstring theory compactified on $\mathrm{AdS}_5 \times S^5$.

4.6.1 The correlation function of sixteen fermionic currents

As could have been anticipated, it is particularly simple to analyze the contribution of the Yang–Mills instanton to the correlation function of sixteen of the fermionic superconformal current bilinears, $\hat{\Lambda}^A_\alpha = \mathrm{tr}\left(\sigma^{\mu\nu}{}_\alpha{}^\beta F^-_{\mu\nu}\lambda_\beta{}^A\right)$,

$$G_{\hat{\Lambda}^{16}}(x_p) = \langle \prod_{p=1}^{16} g^2 \hat{\Lambda}^{A_p}_{\alpha_p}(x_p) \rangle_{K=1} \ , \tag{4.53}$$

Since each factor of $\hat{\Lambda}$ in the product provides a single fermion zero mode it is necessary to consider the product of sixteen currents in order to saturate the sixteen Grassmannian integrals. To leading order in g, $G_{\hat{\Lambda}^{16}}$ does not receive contribution from anti-instantons. The leading term in the one-instanton sector is simply obtained by replacing each $F^-_{\mu\nu}$ with the instanton profile $\overline{F}_{\mu\nu}$ (equation (4.10)) and each $\lambda_\alpha{}^A$ with the corresponding zero mode, $\lambda^A_{(0)\alpha}$

$$\lambda^A_{(0)\alpha} = \frac{1}{2}\overline{F}_{\mu\nu}\sigma^{\mu\nu\,\beta}_\alpha \frac{1}{\sqrt{\rho_0}}\left(\rho_0\eta^A_\beta + (x-x_0)_\mu \sigma^\mu_{\beta\dot{\beta}}\overline{\xi}^{\dot{\beta}A}\right) . \tag{4.54}$$

The resulting correlation function thus has the form

$$G_{\hat{\Lambda}^{16}}(x_p) = \frac{2^{62}3^{16}}{\pi^{10}}\, g^8 e^{-\frac{8\pi^2}{g^2}+i\theta} \int \frac{d^4x_0\, d\rho_0}{\rho_0^5} \int d^8\eta d^8\overline{\xi}\, \cdot$$
$$\cdot \prod_{p=1}^{16}\left[\frac{\rho_0^4}{[\rho_0^2+(x_p-x_0)^2]^4}\,\frac{1}{\sqrt{\rho_0}}\left(\rho_0\eta^{A_p}_{\alpha_p} + (x_p-x_0)_\mu \sigma^\mu_{\alpha_p\dot{\alpha}_p}\overline{\xi}^{\dot{\alpha}_p A_p}\right)\right] . \tag{4.55}$$

The integration over the fermion zero modes leads to a sixteen-index invariant tensor, t_{16}, of the product of the SU(4) and Lorentz groups. Assembling the 16 fermionic collective coordinates into a sixteen-dimensional spinor t_{16} would simply read $t_{16}^{a_1a_2\ldots a_{16}} = \varepsilon^{a_1a_2\ldots a_{16}}$, with $a_i = 1, 2 \ldots 16$. Further integration over the instanton

moduli space would determine the dependence on the coordinates x_p. However, for the purpose of the comparison with the corresponding expression obtained in the type IIB string theory in $\mathrm{AdS}_5 \times S^5$, that will be discussed in the next chapter, it is sufficient to leave the expression in the unintegrated form (4.55).

4.6.2 The correlation function of eight gaugino bilinears

In a similar fashion it is easy to deduce the one-instanton contribution to other related processes, such as the eight-point correlation function,

$$G_{\mathcal{E}^8}(x_p) = \langle g^2 \mathcal{E}^{A_1 B_1}(x_1) \dots g^2 \mathcal{E}^{A_8 B_8}(x_8) \rangle_{K=1} \,, \tag{4.56}$$

which also saturates the sixteen fermionic zero-modes present in the SU(2) one-instanton background. To leading order in g, $G_{\mathcal{E}^8}$ does not receive contribution from anti-instantons. The complete non-abelian expression for $\mathcal{E}^{AB}$ was given in equation (2.19) and reads

$$\mathcal{E}^{AB} = \lambda^{\alpha a A} \lambda_{\alpha a}{}^{B} + g f_{abc}\, t^{(AB)+}_{ijk} \phi^{ia} \phi^{jb} \phi^{kc} \,, \tag{4.57}$$

but at leading order in the gauge coupling constant only the term proportional to the gaugino bilinear is relevant. In the instanton background the gaugino bilinear is given by

$$\lambda^{\alpha a A}_{(0)} \lambda_{(0)\alpha a}{}^{B} = \frac{3 \cdot 2^6}{g^2} \frac{\rho_0^4}{\left(\rho_0^2 + (x - x_0)^2\right)^4} \, \zeta^{\alpha A} \zeta_{\alpha}{}^{B} \,. \tag{4.58}$$

Then it follows

$$\begin{aligned} G_{\mathcal{E}^8}(x_p) = {} & \frac{3^8 2^{14}}{\pi^{10}} \, g^8 \, e^{-\frac{8\pi^2}{g^2} + i\theta} \int \frac{d^4 x_0 \, d\rho_0}{\rho_0^5} \int d^8\eta d^8\overline{\xi} \cdot \\ & \cdot \prod_{p=1}^{8} \left[\frac{\rho_0^4}{\left(\rho_0^2 + (x_p - x_0)^2\right)^4} \frac{1}{\sqrt{\rho_0}} \left(\rho_0 \eta^{A_p}_{\alpha_p} + (x_p - x_0)_\mu \sigma^{\mu}_{\alpha_p \dot{\alpha}_p} \overline{\xi}^{A_p \dot{\alpha}_p} \right) \cdot \right. \\ & \left. \cdot \varepsilon^{\alpha_p \beta_p} \frac{1}{\sqrt{\rho_0}} \left(\rho_0 \eta^{B_p}_{\beta_p} + (x_p - x_0)_\nu \sigma^{\nu}_{\beta_p \dot{\beta}_p} \overline{\xi}^{B_p \dot{\beta}_p} \right) \right] . \end{aligned} \tag{4.59}$$

The integration over the fermion zero modes leads to an SU(4) invariant contraction of the sixteen-index tensor t_{16} defined after (4.55) and further integration over the instanton moduli space would determine the exact dependence on the coordinates x_p. Again the unintegrated expression (4.59) is sufficient for comparison with the D-instanton contribution to the corresponding $\mathrm{AdS}_5 \times S^5$ amplitude that will be considered in the next chapter.

4.7 Generalization to arbitrary N and to any K for large N

In the previous sections instanton calculus in $\mathcal{N}$=4 supersymmetric Yang–Mills theory with gauge group SU(2) in the K=1 sector have been presented following [89]. These results have been extended to the case of SU(N) for generic N in [90] and to arbitrary instanton number in the large N limit in [91, 92]. These extensions are of fundamental importance for the re-interpretation of the results of this chapter in the context of the AdS/CFT correspondence that will be discussed in chapter 5. In this section the main steps of the extensions of [90, 91, 92] will be briefly reviewed. The generalization to $N > 2$ is rather straightforward, on the contrary instanton calculus in sectors $K > 1$ presents an enormous increase of computational complexity. The formalism for multi-instanton calculations (ADHM formalism) has been developed in [101, 102, 103, 104, 105] and a review of the application of these techniques to SU(N) supersymmetric gauge theories can be found in [113]. As has been explained in [91, 92] there are remarkable simplifications in the ADHM formalism in the large N limit.

Since for a gauge group SU(N) there are $2N \cdot \mathcal{N}$ gaugino zero modes in the one-instanton background it might appear from a superficial analysis that the $\hat{\Lambda}^{16}$-correlation function should vanish for $N > 2$. However it has been argued in [89] and then clearly shown in [90] that only the sixteen 'geometric' (8 supersymmetric + 8 superconformal) zero-modes are actually relevant in the general case of SU(N), since all the remaining ones, μ_i^A, $\overline{\mu}_i^A$, are 'lifted' by a four-fermion term that is generated in the one-instanton action. The instanton action computed in [90] has the form

$$S_{\text{inst}} = \frac{8\pi^2 K}{g^2} + S_{\text{4F}} \,, \tag{4.60}$$

where S_{4F} contains a quartic fermionic interaction built up with the $8(N-2)$ additional fermionic collective coordinates. The exact form of S_{4F} in the K=1 sector is

$$S_{\text{4F}} = \frac{\pi^2}{16\rho^2 g^2}\, \varepsilon_{ABCD} \left[\sum_{i=1}^{N-2} \overline{\mu}_i^A \mu_i^B\right] \left[\sum_{j=1}^{N-2} \overline{\mu}_j^C \mu_j^D\right] . \tag{4.61}$$

S_{4F} is supersymmetric and lifts all but the 16 'geometric' gaugino zero modes.

For definiteness in the following the attention will be focused mainly on the $\hat{\Lambda}^{16}$ correlator, that is studied in [90]. In the case of SU(N) there are additional zero-modes contributing to ${\lambda_{(0)\alpha}}^A$ in (4.54). However these are lifted by (4.61) and the

lowest order contribution in the one-instanton background is still obtained by substituting (4.54) in each factor of $\hat{\Lambda}$. As a result the expression for the $\hat{\Lambda}^{16}$-correlator reviewed above for the case N=2 is actually true for all N, apart from an overall N-dependent factor

$$G^N_{\hat{\Lambda}^{16}}(x_p) = C_N\, g^8 e^{-\frac{8\pi^2}{g^2}+i\theta} \int \frac{d^4x_0\, d\rho_0}{\rho_0^5} \int d^8\eta d^8\overline{\xi} \int \prod_{A=1}^{4} \prod_{i=1}^{N-2} d\mu_i^A d\overline{\mu}_i^A\, e^{-S_{\rm 4F}} \cdot$$
$$\cdot \prod_{p=1}^{16} \left[\frac{\rho_0^4}{[\rho_0^2 + (x_p - x_0)^2]^4} \frac{1}{\sqrt{\rho_0}} \left(\rho_0 \eta^{A_p}_{\alpha_p} + (x_p - x_0)_\mu \sigma^\mu_{\alpha_p \dot{\alpha}_p} \overline{\xi}^{\dot{\alpha}_p A_p} \right) \right] , \qquad (4.62)$$

where C_N is a N-dependent constant determined by the norms of the bosonic and fermionic zero-modes. To completely determine the dependence on N one must still compute the integrals over μ_i^A and $\overline{\mu}_i^A$

$$I_N = \int \prod_{A=1}^{4} \prod_{i=1}^{N-2} d\mu_i^A d\overline{\mu}_i^A e^{-S_{\rm 4F}} . \qquad (4.63)$$

In computing the last integral it is convenient to perform a Hubbard-Stratonovich transformation of the fermion bilinears and represent $S_{\rm 4F}$ as a Gaussian integral over auxiliary bosonic 'collective coordinates' χ^i_{AB} [90]. In conclusion one finds the following overall N dependence [90]

$$G^N_{\hat{\Lambda}^{16}}(x_p) \sim \sqrt{N}\, G_{\hat{\Lambda}^{16}}(x_p) , \qquad (4.64)$$

where $G_{\hat{\Lambda}^{16}}(x_p)$ is the expression valid in the N=2 case. This factor of $\sqrt{N}$ will be of relevance in the next chapter for the comparison with D-instanton effects in type IIB superstring theory.

The generalization to multi-instanton sectors in the large N limit, as derived in [91], is much more involved and would require an extended discussion, so only the basic ideas will be summarized here. A detailed report on the construction can be found in [92]. The moduli space of a SU(N) K-instanton configuration, known as ADHM moduli space, has a very complicated structure for generic N. In [91, 92] it has been proved that in the large N limit it is dominated by K instantons living in K mutually orthogonal SU(2) subgroups of SU(N). Then a saddle point approximation is used to show that the geometry is actually described by $(\mathrm{AdS}_5 \times S^5)^K$, where in each factor AdS_5 is parameterized by the positions and sizes of the K instantons, $(x^i_{0\,\mu}, \rho^i)$, $i = 1, \ldots, K$, and S^5 by the auxiliary bosonic coordinates, χ^i_{AB}, introduced in the

Hubbard–Stratonovich transformation. Moreover the integrations in the vicinity of the large N saddle point generate an attractive potential which actually forces all the K instantons to a single point reducing the moduli space to a single copy of $\mathrm{AdS}_5 \times S^5$ [92]. Finally the small fluctuations about this reduced moduli space describe the dimensional reduction of SU(K) $\mathcal{N}$=1 super Yang–Mills from d=10 to d=0+0, so that the K-instanton measure in the large N limit factorizes into the product of the measure on $\mathrm{AdS}_5 \times S^5$ times the partition function $\mathcal{Z}_K$ for the SU(K) (0+0)-dimensional theory. In conclusion, after evaluating $\mathcal{Z}_K$, see [114, 115, 116, 117], one obtains for a n-point correlation function like those considered in this chapter an expression of the form [92]

$$G_n^{(K)}(x_p) = g^8 \sqrt{N} K^{n-\frac{7}{2}} e^{-\frac{8\pi^2 K}{g^2}} \sum_{d|K} \frac{1}{d^2} F_n(x_p)\,, \tag{4.65}$$

where the sum is over the divisors of K and the function $F_n(x_p)$ is independent of K.

Chapter 5

AdS/SCFT correspondence

The remarkable proposal suggested by Maldacena in [38] of a correspondence relating type IIB superstring theory in $\mathrm{AdS}_5 \times S^5$ to $\mathcal{N}$=4 super Yang–Mills theory has led to a great renewal of interest in the latter. According to the formalization of Maldacena's original idea given by Gubser, Klebanov and Polyakov [118] and by Witten [119] $\mathcal{N}$=4 supersymmetric Yang–Mills theory in the superconformal phase is dual to type IIB superstring theory in $\mathrm{AdS}_5 \times S^5$ in the sense that correlation functions of Yang–Mills gauge invariant composite operators can be derived from the computation of amplitudes in the compactification of the type IIB superstring on $\mathrm{AdS}_5 \times S^5$. The precise interpretation of this duality will be discussed later.

More generally the main idea of [38] is that type IIB superstring theory in $\mathrm{AdS}_{d+1} \times M$, where M is a compact manifold with positive curvature, should be equivalent to a (super) conformal field theory living on the d-dimensional boundary of AdS_{d+1}. In this respect the proposed correspondence is holographic in the sense of [120, 121].

Moreover in the case of superconformal field theories with gauge group $\mathrm{SU}(N)$ Maldacena has argued that the large N limit should be equivalent to the supergravity approximation to the AdS compactification of the type IIB superstring. This consideration makes the proposed duality particularly intriguing because it has long been believed that the large N behaviour should be crucial in the understanding of the non-perturbative dynamics of non Abelian gauge theories [122].

Non trivial checks of the suggested correspondence have been obtained in the large

N limit, *i.e.* employing the supergravity approximation, both at the perturbative and at the non-perturbative level. However very little has been said about the extension to finite N and the rôle of string corrections, although arguments have been developed supporting the validity of the correspondence for finite N as well.

The computation of four-point functions in the context of the AdS/SCFT correspondence has also allowed to point out peculiar behaviours in Green functions of the (super) conformal field theories under consideration at short distances. Four-point functions develop logarithmic singularities in the limit of coincident points. This kind of behaviour has already been shown in the instanton contribution to a four-point function in $\mathcal{N}$=4 super Yang–Mills theory in section 4.5. Analogous results have been observed by various authors in Green functions computed from supergravity amplitudes in $\mathrm{AdS}_5 \times S^5$.

Most of the work on the AdS/CFT correspondence has concentrated on the case of $\mathrm{AdS}_5 \times S^5$ which is related to $\mathcal{N}$=4 super Yang–Mills. However considering the compactification of the type IIB superstring on $\mathrm{AdS}_{d+1} \times M_{9-d}$, with M_{9-d} a different compact manifold, one can study superconformal field theories with less supersymmetry. This is a very interesting subject for future developments. In particular the possibility of generalizing the results to non-supersymmetric theories would be of great relevance.

Some recent reviews on the subject can be found in [123].

The chapter is organized as follows. In section 5.1 a very brief introduction to type IIB superstring theory and to its low energy supergravity limit is given. The concept of D-branes, that has played a crucial rôle in Maldacena's construction, is recalled in section 5.2. The original conjecture of [38] together with the successive formalizations are reviewed in section 5.3. The results by various authors in the computation of two- and three-point functions are reported in section 5.4, while section 5.5 contains a discussion on four-point functions. The remaining sections report on original results in the comparison between instanton effects in $\mathcal{N}$=4 super Yang–Mills theory and D-instanton contributions to amplitudes in type IIB superstring theory, following [89].

5.1 Type IIB superstring theory: a bird's eye view

After their introduction in the late sixties in the context of "dual models" for the description of hadronic processes, string theories have emerged as a prime (and up

to now unique) consistent candidate for a unified theory of fundamental interactions including gravity.

A form of the action describing the motion of a string in d-dimensional Minkowski space-time that is suitable for quantization has been proposed in [124, 125]

$$S_{\rm P} = \frac{1}{4\pi\alpha'} \int_\Sigma d\sigma d\tau \, \sqrt{\gamma}\gamma^{ab}\partial_a X^\mu \partial_b X_\mu \,, \tag{5.1}$$

where $X^\mu = X^\mu(\sigma,\tau)$ are the coordinates of the string which give the embedding of the two-dimensional surface spanned by the string (worldsheet) into the d-dimensional space-time (target space), α' is a parameter with dimension $(\text{mass})^{-2}$, related to the string tension T by $T = \frac{1}{2\pi\alpha'}$, and γ^{ab} is the two-dimensional metric.

Roughly speaking the quantization of the bosonic string is achieved by expanding in normal modes the solution of the classical equations of motion which follow from (5.1) and interpreting the coefficients of the expansion as creation and annihilation operators. For open strings this gives rise to a spectrum made of a tachyon with mass $m^2 = -\frac{1}{\alpha'}$ and a massless vector plus an infinite tower of massive states, with masses proportional to $\frac{1}{\sqrt{\alpha'}}$. For closed strings one obtains, beyond the tachyon and a tower of massive states, a rank two tensor, which decomposes into a symmetric traceless tensor, an antisymmetric tensor and a scalar.

The presence of tachyons in the spectrum of bosonic string theories is very problematic. An interesting extension is represented by superstring theories, in which worldsheet supersymmetry is obtained by supplementing the bosonic fields $X^\mu(\sigma,\tau)$ with their superpartners, $\Psi^\mu(\sigma,\tau)$, that are two dimensional Majorana spinors and space-time vectors. Superstring theories are consistently formulated in ten space-time dimensions and, as will be briefly discussed, possess a spectrum of states including a finite set of massless states plus an infinite tower of massive states, but no tachyons.

The observation that two-dimensional gravity is conformally invariant at the classical level is the key point that allows to exploit the powerful tools of conformal field theory to describe the dynamics of the string. This is achieved by mapping the string worldsheet to the complex plane through $(\sigma,\tau) \to z = e^{\tau+i\sigma}$ and introducing bosonic and fermionic coordinates for the string as functions of z and $\overline{z}$. The quantization is then performed employing a BRS formalism to identify the physical states in the Hilbert space generated by "vertex operators". However since the aim of this section is to briefly review the basic ingredients of superstring theories (particularly the type IIB superstring) the more intuitive formalism of oscillators will be used.

The supersymmetric generalization of (5.1) is

$$S = \frac{1}{4\pi\alpha'}\int_\Sigma d\sigma d\tau \left\{\sqrt{\gamma}\gamma^{ab}\left(\partial_a X^\mu \partial_b X_\mu + i\overline{\Psi}^\mu \rho_a \partial_b \Psi_\mu\right) + \right.$$
$$\left. + \overline{\chi}_a \rho^b \rho^a \left(X_\mu \partial_b X^\mu + \frac{1}{2}\overline{\Psi}^\mu \Psi_\mu \chi_b\right)\right\} ,$$

where ρ^a are worldsheet Dirac matrices. χ^a is a Rarita–Schwinger field, the worldsheet gravitino (superpartner of γ_{ab}), which is a space-time scalar. Exploiting worldsheet superconformal symmetry one can put the action in the form

$$S = \frac{1}{4\pi\alpha'}\int d^2\sigma \left[\partial_+ X^\mu \partial_- X_\mu - i\psi_L \cdot \partial_+ \psi_L - i\psi_R \cdot \partial_- \psi_R\right] , \tag{5.2}$$

where + and − indices refer to light-cone coordinates, $\sigma_\pm = \tau \pm \sigma$. The equations of motion coming from (5.2) lead to an expansion of X^μ and ψ^μ in terms of independent left- and right-moving oscillation modes for closed strings. For open strings imposing Neumann (free) boundary conditions makes left- and right-moving degrees equivalent, so that roughly speaking the open string has half the degrees of freedom of a closed string.

The coordinate σ is taken in the range $[0, 2\pi]$ and is periodic; for the fields $\psi_{L,R}$ there are two consistent boundary conditions, either periodic, giving rise to the so called *Ramond sector* (R), or antiperiodic, corresponding to the *Neveu–Schwarz sector* (NS).

The solution of the equations of motion can be written schematically in the form

$$\partial X^\mu_L = \sum_n \alpha^\mu_{-n} e^{in(\tau+\sigma)} \tag{5.3}$$

$$\psi^\mu_L = \sum_n \psi^\mu_{-n} e^{in(\tau+\sigma)} \tag{5.4}$$

and analogously for the right-moving part. The sum in (5.3) is over integer n, while for the fermions in (5.4) n takes integer or semi-integer values in the R and NS sectors respectively. The zero-modes in the expansion of $\partial X^\mu_{L,R}$ correspond to the momentum of the center of mass of the string, $P_L = P_R = P$. For the fermions there are only zero-modes in the R sector that satisfy a Clifford algebra. The ground state of the quantum theory is constructed acting with the creation operators associated with the zero-modes on the trivial vacuum. It is a SO(9,1) spinor in the R sector and a (tachyonic) scalar in the NS sector. A generic state in the spectrum of the string is of the form

$$\alpha^{L\mu_1}_{-n_1} \ldots \psi^{L\mu_k}_{-n_k} \ldots |P_L, a\rangle \otimes \alpha^{R\nu_1}_{-m_1} \ldots \psi^{R\nu_h}_{-m_h} \ldots |P_R, b\rangle ,$$

where conventionally the negative modes are taken as creation operators and a and b refer to the choice of the spin ground state in the R sector. The states are built considering separately the NS and R sectors for left- and right-moving degrees of freedom and then combining them. For the left-movers for instance this construction leads to a tachyonic ground state in the NS sector, while the first higher mass state is a massless vector. In the R sector the ground state corresponds to massless fermions. The correct way of dealing with tachyons was worked out in [36], where a prescription, known as *GSO projection*, was given which leads to space-time supersymmetric string theories, containing no tachyonic state. In the NS sector the GSO projection eliminates the tachyon and leaves with a massless vector, while in the R sector it produces a spinor with definite chirality. Denoting, after the GSO projection, by $\mathbf{v}$ the NS ground state and with $\mathbf{s}$ and $\mathbf{s}'$ the spinors with opposite chirality of the R sector one has the following possibilities. If the GSO projections on L and R moving degrees of freedom are different one gets the type IIA superstring theory, which has space-time supersymmetry $\mathcal{N}$=(1,1) and is non chiral, whereas using the same projection on the L and R parts gives the type IIB string theory, which has $\mathcal{N}$=(2,0) supersymmetry and is chiral. In particular for the type IIB theory, on which the attention will be focused in the following, the massless bosonic degrees of freedom are obtained combining

$$NS \otimes NS \quad \longleftrightarrow \quad \mathbf{v} \otimes \mathbf{v}$$

or

$$R \otimes R \quad \longleftrightarrow \quad \mathbf{s} \otimes \mathbf{s}\,,$$

and correspond to the following field content

$$\begin{aligned} NS \otimes NS \quad &\longleftrightarrow \quad (g_{\mu\nu}, B_{\mu\nu}, \phi) \\ R \otimes R \quad &\longleftrightarrow \quad \left(\chi = C^{(0)}, B'_{\mu\nu}, C^{(4)}\right)\,, \end{aligned}$$

where $g_{\mu\nu}$ is a rank two symmetric traceless tensor (the graviton), $B_{\mu\nu}$ and $B'_{\mu\nu}$ antisymmetric tensors, ϕ a scalar (the dilaton), χ a pseudoscalar (the axion), and $C^{(4)}$ a four-form with self-dual field strength, $F^{(5)} = dC^{(4)} = (*F)^{(5)}$. From the mixed sectors ($R \otimes NS$ and $NS \otimes R$) one obtains the fermionic superpartners of the previous fields.

The low energy effective description of type IIB superstring theory, constructed in terms of the above fields, is type IIB supergravity in ten dimensions, which has chiral

$\mathcal{N}$=(2,0) supersymmetry. There are subtleties in writing down a covariant action for the model because of the self duality constraint on the four form field strength. The covariant equations of motion of the theory have been derived in [126]. Combining the $NS \otimes NS$ scalar and the $R \otimes R$ pseudoscalar into a complex field τ,

$$\tau = \tau_1 + i\tau_2 = \chi + ie^{-\phi}\,, \tag{5.5}$$

and denoting by H_1 and H_2 the field strengths of the $NS \otimes NS$ and $R \otimes R$ two forms, the low energy action of type IIB supergravity can be written, in the Einstein frame, in the form

$$\begin{aligned} S_{\rm IIB} &= \frac{1}{2\kappa_0^2}\int d^{10}X\,\sqrt{g}\left\{R - \frac{1}{2\tau_2^2}\partial_\Lambda\tau\partial^\Lambda\overline{\tau} + \left(F^{(5)}\right)^2 - \right. \\ &\left. -\frac{1}{12\tau_2}\left[\tau H_1 + H_2\right]_{\Lambda\Gamma\Pi}\left[\overline{\tau}H_1 + H_2\right]^{\Lambda\Gamma\Pi}\right\} + \dots\,, \end{aligned} \tag{5.6}$$

where κ_0 is the ten-dimensional Newton constant ($\kappa_0 \sim (\alpha')^2$). In (5.6) the dots stand for higher derivative terms and the fermionic fields have not been displayed. Here and in the following capital Greek letters refer to flat ten-dimensional space-time. In (5.6) the notation $\left(F^{(5)}\right)^2$ is schematic, since a covariant kinetic term for the $R \otimes R$ 4-form cannot be written down naively because of the self-duality constraint. The correct way to deal with this problem was described in [127]. The classical action is invariant under SL(2,$\mathbb{R}$) transformations that act projectively on the complex scalar τ. Under the SL(2,$\mathbb{R}$) group the $NS \otimes NS$ and $R \otimes R$ antisymmetric tensors form a doublet. The scalar τ parameterizes the coset space SL(2,$\mathbb{R}$)/U(1)$_B$, where the U(1)$_B$ is an anomalous R-symmetry acting on the fermions. In particular the left chirality gravitino carries charge $+\frac{1}{2}$ under this symmetry and the right-chirality dilatino present in the type IIB spectrum has charge $+\frac{3}{2}$. At the quantum level the continuous SL(2,$\mathbb{R}$) symmetry is expected to be broken to SL(2,$\mathbb{Z}$) [128].

For the calculations presented in the following the only relevant terms in the effective action are those involving only derivatives of the metric apart from some overall function of the complex scalar. For future purposes it is useful to write these terms in the string frame that is related to the Einstein frame by a Weyl rescaling

$$g^{(s)}_{\mu\nu} = e^{\phi/2}\, g^{(E)}_{\mu\nu}\,.$$

The leading terms in the derivative expansion that will be considered take the form [114]

$$(\alpha')^{-4}\int d^{10}X\sqrt{g}\left[e^{-2\phi}R + \kappa(\alpha')^3 f_4(\tau,\bar{\tau})e^{-\phi/2}\mathcal{R}^4\right]\,. \tag{5.7}$$

where κ is a numerical constant. The Riemann curvature enters the $\mathcal{R}^4$ factor in a manner that may be most compactly described by writing it as an integral over a sixteen-component Grassmann spinor,

$$\mathcal{R}^4 \equiv \int d^{16}\Theta (R_{\Theta^4})^4 \,,$$

where

$$R_{\Theta^4} = \overline{\Theta}\Gamma^{\Lambda_1\Lambda_2\Lambda}\Theta\,\overline{\Theta}\Gamma^{\Lambda_3\Lambda_4}{}_{\Lambda}\Theta\, R_{\Lambda_1\Lambda_2\Lambda_3\Lambda_4} \,,$$

which only includes the Weyl tensor piece of the Riemann tensor. Here, $\Gamma^{\Lambda_1\Lambda_2\Lambda_3}$ are the totally antisymmetric products of three ten-dimensional Γ-matrices and the Grassmann parameter Θ^a $(a = 1, \ldots, 16)$ is a chiral spinor of the ten-dimensional theory. This expresses $\mathcal{R}^4$ as an integral over half of the on-shell type IIB superspace.

Beyond these terms there is a sixteen-dilatino term in the effective action that is related to the $\mathcal{R}^4$ term by supersymmetry and will play a central rôle in the comparison with the results of instanton calculus presented in the previous chapter. The sixteen fermion interaction reads [129, 130]

$$(\alpha')^{-1} \int d^{10}X \sqrt{g}\, e^{-\phi/2} f_{16}(\tau, \overline{\tau}) \Lambda^{16} + \text{c.c.} \,, \tag{5.8}$$

where the complex chiral dilatino has been denoted by Λ and the interaction is (totally) antisymmetric in the sixteen spinor indices. The properties of the modular functions $f_4(\tau, \overline{\tau})$ and $f_{16}(\tau, \overline{\tau})$ in equations (5.7) and (5.8) will be discussed later. They receive both perturbative corrections and non-perturbative contributions from D-instantons [114]. The latter will be related to instanton effects in $\mathcal{N}$=4 super Yang–Mills theory.

5.2 D-branes

The perturbative spectrum of type II (A and B) superstring theories does not contain states charged with respect to the fields in the $R \otimes R$ sector [131]. However the flurry of work on string dualities during the last years has lead to a much deeper understanding of non-perturbative aspects of string theories, changing dramatically the scenario which was established at the perturbative level.

From a low energy viewpoint the existence of solitonic solutions of the supergravity equations of motion has been known for a long time [132]. These solitonic

configurations describe extended objects, *p-branes*, that are naturally coupled to the $R \otimes R$ forms and their duals. The rôle of these non perturbative states in the fundamental string theory has been clarified by Polchinski [133] through the concept of *D-branes*.

A p-brane is p-dimensional object, whose world-volume is $p+1$-dimensional. It naturally couples to a $(p+1)$-form, $C^{(p+1)}$, through an action of the form

$$\int_{V_{p+1}} C^{(p+1)} \,,$$

where V_{p+1} is the world-volume of the brane. In d space-time dimensions the corresponding "electric" charge would be

$$Q_e = \int_{S^{d-p-2}_\infty} *dC^{(p+1)} \,.$$

The magnetic dual of a p-brane is a $(d-p-4)$-brane, which couples to a $(d-p-3)$-form and carries "magnetic" charge

$$Q_m = \int_{S^{p+2}_\infty} dC^{(p+1)} \,.$$

In particular in the case of the type IIB theory in the $R \otimes R$ sector there is a scalar, the axion χ, *i.e.* a zero-form $C^{(0)} \equiv \chi$, a rank two anti-symmetric tensor, $B'_{\mu\nu}$, *i.e.* a two-form $C^{(2)}$ and a four form $C^{(4)}$. Coupled to $C^{(0)}$ one has a (-1)-brane which is an object localized in space-time, *i.e.* an instanton in Euclidean space-time. Effects of D-instantons in type IIB superstring theory will be discussed later. The two-form couples to a one-dimensional object, a D-string that should not be confused with the fundamental string which couples to the $NS \otimes NS$ two-form. Finally there is a 3-brane charged with respect to the four-form $C^{(4)}$. Moreover one can consider the corresponding magnetically dual objects.

Starting from the type II supergravity action, extremal (*i.e.* saturating a BPS bound) solitonic configurations with the previously discussed properties can be constructed assuming that only the metric $g_{\mu\nu}$, the dilaton ϕ and a $p+1$ form are non-vanishing. The ansatz that one considers for these fields is

$$\begin{aligned} ds^2 &= e^{2A(r)} d\vec{x}^2 + e^{2B(r)} d\vec{y}^2 \\ \phi &= \phi(r) \\ C^{(p+1)}_{01\ldots p} &= -e^{C(r)} \,, \end{aligned}$$

where the ten-dimensional coordinates X^Λ are split into longitudinal (x^μ, $\mu = 0, 1, \ldots, p$) and transverse (y^i, $i = p+1, \ldots, 9$) coordinates and $r^2 = y_i y^i$. The solution of the supergravity equations can be written in terms of a single harmonic function $H_p(r)$, *i.e.* a solution of the Laplace equation in the $9-p$ transverse dimensions,

$$H_p(r) = 1 + \frac{a_p}{r^{7-p}} \,,$$

where a_p is a constant with dimension L^{7-p}, related to the charge carried by the p-brane. The p-brane solution is (in the Einstein frame)

$$\begin{aligned} ds^2 &= [H_p(r)]^{\frac{p-7}{8}}\, d\vec{x}^2 + [H_p(r)]^{\frac{p+1}{8}}\, d\vec{y}^2 && (5.9)\\ C^{(p+1)}_{01\ldots p}(r) &= [H_p(r)]^{-1} - 1 && (5.10)\\ e^{2\phi(r)} &= [H_p(r)]^{\frac{3-p}{2}} \,. && (5.11) \end{aligned}$$

Notice in particular that in the case of a 3-brane the dilaton ϕ reduces to a constant. The solution simplifies in the string frame where the metric becomes

$$ds^2 = [H_p(r)]^{-\frac{1}{2}}\, d\vec{x}^2 + [H_p(r)]^{\frac{1}{2}}\, d\vec{y}^2 \,.$$

It can be proved that these solitonic solutions are BPS saturated and preserve one half of the supersymmetries of the original flat background, *i.e.* 16 supercharges out of 32. This implies for instance that there exist a relation between the mass and charge of the solitons.

Many of the recent developments in string theory have been made possible by the suggestion [133] of a way to incorporate the above solitonic objects into type II string theories, see [134] for a review.

The solitonic p-branes are introduced in the microscopic string theory in relation with a (Dirichlet) open string subsector. More precisely one introduces *Dirichlet p-branes* (or Dp-branes) as p-dimensional hyperplanes where open strings can end, satisfying Neumann (standard) boundary conditions on the p longitudinal coordinates X^μ, $\mu = 0, \ldots, p-1$ and Dirichlet (fixed) conditions on the remaining transverse coordinates X^m, $m = p, \ldots, 9$. D-branes defined in this way can be seen as 'topological defects' where open strings can end. These hyperplanes are actually dynamical objects, whose elementary excitations are described by the open strings attached to them. These open strings can be consistently introduced in type II superstring theories, without considering 'standard' open strings propagating independently in the target space. There are various arguments that lead to identify D-branes with

the solitonic p-branes of type II supergravities. For instance considering two parallel D-branes their interaction is mediated by the exchange of closed strings and one can prove that this results in a vanishing total force just like for example is found for a monopole pair. Furthermore it can be shown that the boundary conditions imposed on the open strings ending on the D-brane break half the supersymmetries, so that the D-branes are BPS saturated objects.

The dynamics of D-branes, in a microscopic string theory perspective, is described by the open strings attached to the brane. Comparing the effective world-volume dynamics of the D-brane with the dynamics of solitonic p-brane solutions in the moduli space approximation one can find further evidence for the identification of the two.

The massless spectrum of the subsector of open strings attached to a p D-brane is a maximally supersymmetric U(1) gauge theory in $p+1$ dimensions. The effective world-volume degrees of freedom of a p D-brane correspond to the dimensional reduction of a $\mathcal{N}$=1 vector multiplet from d=10 to $d = p+1$. This gives rise to a supersymmetric Maxwell theory (with sixteen supercharges) including $9-p$ scalars, one vector and their superpartners.

Considering N D-branes one can have also open strings stretched between them. They describe the exchange of massive vector states. There are N^2 ways of stretching the strings among the branes. In the limit in which the N D-branes are taken to coincide these N^2 different states become massless. The symmetry is enhanced to U(N) and one is left with an effective U(N) Yang–Mills theory in which the N^2 vectors form the adjoint. In general the relative separations among the branes determine the expectation values of the $9-p$ scalars in the effective theory.

In particular in the case of a collection of N coincident D3-branes the world-volume effective action is a U(N) supersymmetric Yang–Mills theory with sixteen supercharges in d=4, *i.e.* the $\mathcal{N}$=4 super Yang–Mills theory with gauge group U(N) [1].

[1] Actually there is a U(1) factor that describes the center of mass motion and decouples, so that the gauge group is SU(N) [135].

5.3 The AdS/SCFT correspondence conjecture

In the case of D3-branes the solution for the metric yields (in the string frame)

$$ds^2 = \left[1 + \frac{L^4}{r^4}\right]^{-\frac{1}{2}} d\vec{x}^2 + \left[1 + \frac{L^4}{r^4}\right]^{\frac{1}{2}} d\vec{y}^2 \,, \tag{5.12}$$

where (x^μ, y^i) are the previously introduced Cartesian coordinates and L is a length scale. As already noticed in the case of a D3-brane the dilaton is constant, $\phi = \phi_0$, and determines the string coupling constant

$$g_s = e^{\phi_0} \,.$$

Considering a D3-brane configuration with N units of $R \otimes R$ 5-form field strength, *i.e.* a configuration of N coincident D3-branes, the length L is

$$L^4 = 4\pi g_s N \alpha'^2 \,. \tag{5.13}$$

The key observation which led Maldacena to the formulation of the AdS/CFT correspondence conjecture [38] is that the metric (5.12), in the "near horizon" limit $r \to 0$, reduces to

$$ds^2 = \frac{L^2}{\rho^2}\left(dx \cdot dx + d\rho^2\right) + d\omega_5^2 \,, \tag{5.14}$$

where $\rho^2 = \frac{L^4}{r^2}$ and $d\omega_5^2$ is the spherically symmetric constant curvature metric on a 5-sphere, which is the metric of the $\mathrm{AdS}_5 \times S^5$ space. In (5.14) L is the radius of curvature of both the 5-sphere and the AdS factor. The D3-brane metric can be viewed as a solitonic configuration interpolating between two maximally supersymmetric backgrounds, the near horizon $\mathrm{AdS}_5 \times S^5$ and flat ten dimensional Minkowski space-time at infinity.

Maldacena has argued, on the basis of previous results in the context of the description of black holes by D-branes [136], that the region that should be identified with the Yang–Mills low energy description is the "throat", *i.e.* the region $r \ll L$, in the D3-brane metric (5.12). See [137] for related work. This leads to the conjecture that $\mathcal{N}$=4 supersymmetric Yang–Mills theory should be dual to type IIB superstring theory compactified on $\mathrm{AdS}_5 \times S^5$. The original conjecture relates type IIB supergravity to SU(N) $\mathcal{N}$=4 super Yang–Mills in the large N limit. This is related to the fact that the limit $g_s \ll 1$ at large length scale with respect to the

string scale, $\frac{\alpha'}{L^2} \ll 1$, in which one can trust the supergravity approximation, requires $N \to \infty$ with $4\pi g_s N$ fixed and large. The correspondence relates the complexified Yang–Mills coupling constant to the one of the type IIB superstring in $\mathrm{AdS}_5 \times S^5$ by

$$g_s = \frac{g^2_{\mathrm{YM}}}{4\pi}\,, \qquad \chi_0 = \frac{\theta_{\mathrm{YM}}}{2\pi}\,, \tag{5.15}$$

where χ_0 is the constant $R \otimes R$ scalar and from now on the Yang–Mills coupling constant and vacuum-angle will be denoted respectively by g_{YM} and θ_{YM}.

This implies that the large N limit of relevance here coincides with the large N limit considered by 't Hooft in [122], with the effective coupling $\hat{g} = g_{\mathrm{YM}} N^2$ fixed at a large value.

The AdS_5 space has constant negative curvature and possesses a boundary that is the four dimensional Minkowski space [2]. According to Maldacena's proposal, the boundary of AdS_5, which corresponds to $\rho \to 0$, *i.e.* $r \to \infty$ in the notation of (5.14), is exactly the location of the $\mathcal{N}$=4 theory and the boundary values of the bulk supergravity fields act as sources that couple to gauge-invariant composite operators in $\mathcal{N}$=4 super Yang–Mills theory. The correct interpretation of this statement has been explained in [118, 119] and will be discussed soon.

The $\mathrm{AdS}_5 \times S^5$ background is characterized by the non-vanishing fields,

$$\begin{aligned} F_{_{MNPQR}} &= \frac{1}{L}\varepsilon_{_{MNPQR}} \qquad R_{_{MNPQ}} = -\frac{1}{L^2}(g_{_{MP}}g_{_{NQ}} - g_{_{MQ}}g_{_{NP}}) \\ F_{mnpqr} &= \frac{1}{L}\varepsilon_{mnpqr} \qquad R_{mnpq} = +\frac{1}{L^2}(g_{mp}g_{nq} - g_{mq}g_{np})\,, \end{aligned}$$

where upper case Latin indices, $M, N, \cdots = 0, 1, 2, 3, 5$, span the AdS_5 coordinates and lower case Latin indices, $m, n, \cdots = 1, 2, 3, 4, 5$ span the S^5 coordinates. The only non-vanishing components of the Ricci tensor are

$$R_{_{MN}} = -\frac{4}{L^2}g_{_{MN}} \qquad R_{mn} = +\frac{4}{L^2}g_{mn}\,. \tag{5.16}$$

Upon contracting (5.16) with the metric tensor it follows that the total (5+5)-dimensional scalar curvature vanishes. The Weyl tensor, defined in d-dimensional space-time as

$$\begin{aligned} C_{\mu\nu\rho\sigma} &= R_{\mu\nu\rho\sigma} - \frac{1}{d-2}\left[R_{\mu\nu}g_{\nu\sigma} + (3\text{ terms})\right] + \\ &+ \frac{1}{(d-1)(d-2)}R\left(g_{\mu\rho}g_{\nu\sigma} - g_{\mu\sigma}g_{\nu\rho}\right)\,, \end{aligned}$$

[2] Actually this is better understood considering the Euclidean version of the AdS_5 space, see [119] for a detailed description.

vanishes as well in $\mathrm{AdS}_5 \times S^5$ because of the conformal flatness of the metric.

This background is maximally supersymmetric (just like the Minkowski vacuum) so there are 32 conserved supercharges, that transform as a complex chiral spinor of the tangent-space group, SO(4,1)×SO(5). In the basis where the ten dimensional Γ_Λ matrices are given by $\Gamma_M = \sigma_1 \otimes \gamma_M \otimes \mathbb{1}$ and $\Gamma_m = \sigma_2 \otimes \mathbb{1} \otimes \gamma_m$, the supersymmetries are generated by the Killing spinors that satisfy

$$D_\Lambda \epsilon - \frac{1}{2L}(\sigma_1 \otimes \mathbb{1} \otimes \mathbb{1})\Gamma_\Lambda \epsilon = 0\,, \tag{5.17}$$

which follows from the requirement that the gravitino supersymmetry transformation should vanish. In this basis the complex chiral supersymmetry parameters read

$$\epsilon_\pm = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \zeta_\pm \otimes \kappa_\pm\,,$$

where $\zeta_\pm$ are complex four-component SO(4,1) spinors and $\kappa_\pm$ complex four-component SO(5) spinors, satisfying

$$\begin{aligned} D_M \zeta_\pm \mp \frac{1}{2L}\gamma_M \zeta_\pm &= 0 \\ D_m \kappa_\pm \mp i\frac{1}{2L}\gamma_m \kappa_\pm &= 0\,. \end{aligned}$$

Strong support to the conjecture comes from symmetry considerations. AdS_5 has a SO(2,4) isometry group which is to be identified with the four-dimensional conformal group in the boundary theory. The isometry group of the S^5 factor is SO(6)∼SU(4) and is related to the SU(4) R-symmetry group of the $\mathcal{N}$=4 Yang–Mills theory. Including the associated fermionic generators the super-isometry group of the type IIB background (5.14) is exactly the superconformal group SU(2,2|4) of $\mathcal{N}$=4 super Yang–Mills in four dimensions. Moreover the SL(2,$\mathbb{Z}$) S-duality group that is conjectured to be preserved in the type IIB superstring at the quantum level is identified with the (conjectured) exact S-duality symmetry group of four dimensional SU(N) $\mathcal{N}$=4 Yang–Mills theory. The former is connected to the existence of an infinite set of stable dyonic BPS string solitons [128], the latter to the existence an infinite set of stable dyonic BPS states [72], as discussed in chapter 2.

The spectrum of excitations of the maximally supersymmetric compactification of the type IIB superstring on the $\mathrm{AdS}_5 \times S^5$ background, first considered in [126], was studied in great detail in [138, 139]. The map that associates fields in the multiplet of type IIB supergravity to the corresponding operators in $\mathcal{N}$=4 super Yang–Mills

has been worked out in [140, 141, 142]. States in the spectrum of d=5 $\mathcal{N}$=8 gauged supergravity admit a natural classification in terms of spherical harmonics on the 5-sphere. In the conjectured correspondence the lowest lying states, assembled into the $\mathcal{N}$=8 gauged supergravity multiplet, couple to the supermultiplet of super Yang–Mills currents, equations (2.17) and (2.18), that are "bilinear" in the elementary Yang–Mills fields. Higher ℓ spherical harmonics, *i.e.* Kaluza–Klein (KK) descendants of the supergravity fields, couple to other "chiral primary fields", that correspond to "ℓth-order" gauge-invariant polynomials in the elementary super Yang–Mills fields. Moreover, stable string excitations that couple to non-chiral primary fields are needed for the closure of the operator algebra [143]. Despite some interesting progress [144], a useful worldsheet description for the type IIB superstring in $\mathrm{AdS}_5 \times S^5$ is still lacking. Nevertheless the stringy truncation of the KK spectrum to $\ell \leq N$ seems to be a robust result that matches with the naive expectations on the super Yang–Mills side.

The explicit connection between the bulk theory and the boundary theory has been given a precise formulation in [118, 119]. The partition function of the type IIB superstring, computed with suitable boundary conditions on the four-dimensional boundary of AdS_5, plays the rôle of generating functional for connected Green functions of gauge-invariant composite operators in $\mathcal{N}$=4 Yang–Mills theory. More precisely one identifies

$$Z_{\text{IIB}}[J] = \int [DA]\, \exp(-S_{\text{YM}}[A] + \mathcal{O}[A]J)\,, \tag{5.18}$$

where $Z_{\text{IIB}}[J]$ is the partition function of the type IIB superstring, evaluated in terms of the bulk string fields, which is computed with prescribed boundary values J for the fields. Elementary fields of the $\mathcal{N}$=4 super Yang–Mills theory on the boundary are denoted by A and $\mathcal{O}(A)$ are gauge-invariant composite operators to which $J(x)$'s couple.

The left hand side of (5.18) is computed performing the functional integration over the bulk fields $\Phi(z;\omega)$, where ω are the coordinates on S^5 and $z^M \equiv (x^\mu, \rho)$ ($M = 0, 1, 2, 3, 5$ and $\mu = 0, 1, 2, 3$) the coordinates on AdS_5 ($\rho \equiv z_5$ is the coordinate transverse to the boundary)

$$Z_{\text{IIB}}[J] = \int [D\Phi]_J \exp(-S_{\text{IIB}}[\Phi])\,. \tag{5.19}$$

In (5.19) the notation $\Phi(z;\omega)$ refers collectively to the 'massless' supergravity fields,

their Kaluza–Klein descendents, and possibly the string excitations. The functional integral depends on the boundary values, $J(x)$, of the bulk fields.

The exact map which associates fields in the type IIB compactification with operators in $\mathcal{N}$=4 super Yang–Mills theory is dictated by the matching of quantum numbers corresponding to the previously discussed global symmetries. Furthermore the conformal dimension, Δ, of a Yang–Mills operator $\mathcal{O}$ is related to the AdS 'mass' m of the corresponding supergravity field. For instance in the case of scalar operators the relation is

$$(mL)^2 = \Delta(\Delta - 4)\,, \tag{5.20}$$

where only the positive branch of $\Delta_\pm = 2 \pm \sqrt{4 + (mL)^2}$ is relevant for the lowest-'mass' supergravity multiplet.

As already remarked in the Kaluza–Klein reduction of type IIB supergravity on S^5 the states are classified expanding the ten-dimensional fields in spherical harmonics on the 5-sphere. This procedure generates an infinite tower of states with spins up to 2. The masses of the states associated with single modes and their behaviour under the SO(2,4) isometry group of AdS_5 are determined by solving the corresponding linearized equations of motion [139].

Alternatively the Kaluza–Klein excitations of all the fields in the type IIB gauged supergravity can be constructed, using the Killing spinors defined in equation (5.17), from the modes of the massless singlet dilaton e^ϕ. The construction is rather involved and will be briefly sketched only for fields that will be relevant in the following.

One obtains a classification of the states according to the representations of the isometry group of $AdS_5 \times S^5$, which adding the associated fermionic generators becomes SU(2,2|4). In particular the mass of the states is identified with the eigenvalue of one of the generators of SO(2,4). This allows to construct a map which relates each field in the supergravity multiplet to the corresponding composite operator in $\mathcal{N}$=4 super Yang–Mills theory by matching the SU(2,2|4) quantum numbers (recall that on the super Yang–Mills side SU(2,2|4) is the $\mathcal{N}$=4 superconformal group). Notice that since the states in the supergravity multiplet as well as their KK descendants have spins not larger than 2, they belong to short multiplets of SU(2,2|4).

The massless dilaton is associated with the constant mode on S^5, *i.e.* with the scalar spherical harmonic $Y_\ell(\omega)$ with $\ell = 0$. Also at the $\ell = 0$ level there is the five-dimensional graviton, describing the excitations of the AdS_5 metric. The other scalars in the supermultiplet are associated with excitations on the 5-sphere, *i.e.* they

have $\ell > 0$. In particular there are real scalars Q^{ij} with mass $m^2 = -\frac{4}{L^2}$ in the $\mathbf{20}_{\mathbb{R}}$ of the $SO(6)$ isometry group of S^5. They result from a combination of the trace of the internal metric and the self-dual $R \otimes R$ five-form field, $F^{(5)}$, with $\ell = 2$ (Q^{ij} are quadrupole moments of S^5). The trace part of the metric fluctuations can be written in the form

$$\delta g_{MN} = \frac{16}{15} f \delta_{MN} \qquad \delta g_{mn} = f \delta_{mn} \,,$$

where $f = f(z, \omega)$. Analogously for the $R \otimes R$ 4-form one gets

$$A_{MNPQ} = -\frac{L}{4} \varepsilon_{MNPQR} \nabla^R f \qquad A_{mnpq} = -\frac{L}{4} \varepsilon_{mnpqr} \nabla^r f \,.$$

The relation with the scalar fields $Q_{ij}(z)$ is of the form

$$f(z, \omega) = Q_{ij}(z) Y^{ij}_{\ell=2}(\omega) \,.$$

Similarly the complex scalars E^{AB} with mass $m^2 = -\frac{3}{L^2}$ and their conjugates are associated with the pure two-form fluctuations with $\ell = 1$ of the complexified antisymmetric tensor, constructed from the $R \otimes R$ and $NS \otimes NS$ two-forms, in the internal directions. The massless vectors $V_M^{[ij]}$ in the **15**, that gauge the SO(6) isometry group, are in one-to-one correspondence with the Killing vectors of S^5 and result from a linear combination with $\ell = 1$ of the mixed components of the metric and the internal three-form components of the $R \otimes R$ four-form potential, $C^{(4)}$. The **6** complex antisymmetric tensors $B_{MN}^{[AB]}$ with $m^2 = \frac{1}{L^2}$, that have peculiar first order equations of motion, result from scalar spherical harmonics with $\ell = 1$. The analysis of the fermions is similar. The dilatini Λ^A in the **4** of SO(6) are proportional to the internal Killing spinors κ_+ and have mass $m = -\frac{3}{2L}$. They play a central rôle in the comparison between the effects of instantons in $\mathcal{N}$=4 Yang–Mills and D-instantons in type IIB string theory that has been carried out in [89] and will be reviewed in section 5.6. The $\mathbf{20}_{\mathbb{C}}$ spinors χ^A_{BC} with mass $m = -\frac{1}{2L}$ correspond to internal components of the gravitino with $\ell = 1$. Finally the supergravity multiplet is completed by the massless $\overline{\mathbf{4}}$ gravitinos Ψ_{MA} which are proportional to the internal Killing spinors κ_-.

The above fields are the ones that act as sources for the superconformal currents (2.17), (2.18). More precisely the graviton and dilaton are associated with the SU(4) singlets in the $\mathcal{N}$=4 current multiplet, the stress-energy tensor $\mathcal{T}_{\mu\nu}$ and the scalar $\mathcal{C}$ respectively. The other scalars, E^{AB} and Q^{ij}, correspond to $\mathcal{E}^{AB}$ and $\mathcal{Q}^{ij}$ in (2.18) respectively. The massless vectors $V_M^{[ij]}$ are dual to the SU(4) currents $\mathcal{J}^A_{\mu B}$

and the anti-symmetric tensors $B^{[AB]}_{MN}$ to $\mathcal{B}^{AB}_{\mu\nu}$. The identification can be carried out analogously for the fermions. The dilatini Λ^A and the spinors χ^A_{BC} are associated respectively to the fermions $\hat{\Lambda}^A$ and $\hat{\chi}^A_{BC}$ in (2.18). The gravitinos $\Psi_{M\,A}$ correspond to the supersymmetry currents $\Sigma_{\mu A}$.

Higher Kaluza–Klein modes correspond to higher values of ℓ. For example, there are other scalar modes with $\ell > 2$. Each of these can be put in one-to-one correspondence with a gauge singlet composite operator $W_{(\ell)}$ that starts with

$$W^{(i_1\ldots i_\ell)}\big|_{\theta=0} = \mathrm{tr}\left(\phi^{(i_1}\ldots\phi^{i_\ell)}\right) - \text{traces}\,, \tag{5.21}$$

which has dimension $\Delta = \ell$ and belongs to the ℓ-fold symmetric traceless tensor representation of $SO(6)$ with Dynkin labels $(0,\ell,0)$. The multiplet contains $256\ell^2(\ell^2-1)/12$ states with different Δ and SU(4) quantum numbers.

The correct recipe for calculating correlations follows from (5.18) and amounts to computing the truncated Green functions in the bulk and attaching *bulk-to-boundary propagators* to the external legs [118, 119, 145]. The precise forms of the latter depend on the spin and 'mass' of the field. For instance, the bulk-to-boundary Green function for a bulk scalar field Φ_m associated to a dimension Δ operator $\mathcal{O}_\Delta$ on the boundary reads

$$K_\Delta(x^\mu,\rho;x'^\mu,0) = c_{_\Delta}\frac{\rho^\Delta}{(\rho^2+(x-x')^2)^\Delta}\,, \tag{5.22}$$

where $c_{_\Delta}$ is a constant given by $c_{_\Delta} = \Gamma(\Delta)/[\pi^2\Gamma(\Delta-2)]$. For simplicity the ω dependence has been suppressed, so that the expression (5.22) is appropriate for an 'S-wave' process in which there are no excitations in the directions of the five-sphere, S^5. In terms of K_Δ the bulk field

$$\Phi_m(z;J) = \int d^4x' K_\Delta(x,\rho;x',0)J_\Delta(x') \tag{5.23}$$

satisfies the boundary condition $\Phi_m(x,\rho;J) \approx \rho^{4-\Delta}J_\Delta(x)$ as $\rho\to 0$, since $\rho^{\Delta-4}K_\Delta$ reduces to a δ-function on the boundary. As observed in [145] the bulk-to-boundary propagator must be modified for Δ=2. In this case the previous expression vanishes since $c_{_\Delta}$=0 for Δ=2. In the special case Δ=2 one must require $\Phi_m(x,\rho;J)\to\rho^2\log\rho\,J(x)$, which yields $c_{_{\Delta=2}} = \frac{\Gamma(2)}{2\pi^2}$. Before discussing specific calculations, that will be related to instanton effects in $\mathcal{N}$=4 supersymmetric Yang–Mills theory, this prescription for evaluating Green functions will be illustrated in more detail in the case of two- and three-point functions in the next section.

The generalization of this construction to fermionic fields has been considered in [146] and will be relevant later when the calculation of a correlation function of sixteen fermionic dilatini will be presented.

The original conjecture involves the supergravity approximation and holds in the large N limit. However in [38] it has been suggested that the correspondence can be generalized to finite N by taking in account string theory corrections. This claim has been made more explicit in [119]. For finite N the rôle of generating functional is played by the full partition function of the string in $\mathrm{AdS}_5 \times S^5$, computed with suitable boundary conditions. Moreover for finite N, *i.e.* for finite radius, one expects a truncation to $\ell \leq N$ of the Kaluza–Klein spectrum, that should result by taking into account full-fledged stringy geometry as in any string compactifications.

Perturbative checks of the conjecture at finite N for two- and three-point functions have been given in [147, 148] and will be discussed in the next section. The non-perturbative computations of [89] suggest that the correspondence remain valid at finite N also for four- or higher-point functions of some protected operators at least.

Witten has also suggested an intuitive way to represent diagrammatically the above described construction through graphs like those depicted in figure 5.1.



Figure 5.1: Witten's representation of three point functions in the AdS/CFT correspondence.

The circles in figure 5.1 represents the four-dimensional boundary of AdS_5 and the interior is the bulk. Lines connecting an internal point with a point on the boundary are the above discussed bulk-to-boundary propagators (see *e.g.* equation (5.22)). In the supergravity approximation, *i.e.* in the limit of large N and large and fixed $g^2_{\mathrm{YM}}N$, only tree diagrams like the first one in figure 5.1 contribute, while the loop-diagrams become relevant for finite N, when string effects must be taken into account. This

can be understood as the loops correspond to closed string exchange which gives a contribution that is suppressed in the large N limit.

Attempts have also been made to generalize the correspondence to conformal theories with less supersymmetry and hopefully to non-supersymmetric theories. This is achieved by studying the compactification of the type IIB superstring on $\mathrm{AdS}_5 \times X_5$, where X_5 is a compact Einstein manifold, *i.e.* a positively curved compact manifold with metric satisfying $R_{\mu\nu} = \lambda g_{\mu\nu}$, with $\lambda > 0$. The simplest examples are obtained considering $X_5 = S^5/\Gamma$, where Γ is a discrete subgroup of SO(6). In this case X_5 presents orbifold type singularities, but it has locally the geometry of S^5. In [149, 150] this models have been discussed in detail and it has been shown that taking N coincident D3-branes placed at an orbifold singularity one can construct $\mathcal{N}=2$ and $\mathcal{N}=1$ theories. Such theories have product gauge groups $\mathrm{SU}(N)^k$ and contain matter fields in the fundamental and adjoint representations.

A possible approach to the study of non conformal field theories through the AdS correspondence has been suggested in [151]. The model proposed in [151] is based on the observation that field theories at finite temperature can be described by near-extremal p-brane solutions. Non supersymmetric theories are then obtained by imposing anti-periodic boundary conditions for the fermions on the compactified direction, corresponding to the raising of temperature, which explicitly break supersymmetry. Qualitative agreement with the expectations for QCD at strong coupling has been found following this approach. In particular the results from supergravity calculations include the area low behaviour of Wilson loops, the presence of a mass gap for glueball states, the relation between confinement and dual superconductivity and the existence of heavy quark baryonic states [151, 152]. Numerical calculations of glueball masses have been performed in [153] applying the same method. The results appear to be in agreement with those obtained from lattice simulations. The limit of this approach is that for the supergravity approximation to be reliable one must take the effective 't Hooft coupling to be large, while the weak coupling region is the one relevant for the continuum limit on the lattice. As a consequence to actually compare the results from AdS with lattice extrapolations to the continuum, it seems to be necessary to extend the correspondence to finite N, which requires a full string theory description in AdS.

More recently the near-horizon geometry of a stack of D3-branes in type 0 theories has been proposed as a candidate for a non supersymmetric version of the above discussed correspondence. In these models the *"tachyon should ... peacefully*

condense in the bulk" [154]. The basic idea suggested by Polyakov in [154] is that non-supersymmetric theories should have as their duals string theories displaying worldsheet supersymmetry, but no space-time supersymmetry. This is the property characterizing type 0A and 0B string theories, which have been introduced in [155]. This theories are constructed by a non-chiral GSO projection, which explicitly breaks space-time supersymmetry and leads to a spectrum containing no space-time fermions. The bosonic spectrum is the same as in the corresponding type II theories in the $NS\otimes NS$ sector and contains a doubled set of $R\otimes R$ fields. The interest in these theories has been renewed in the systematic construction of their open string descendants [156]. The inclusion of D-branes in such theories has been studied in [156, 157] and allows to construct non-supersymmetric gauge theories as world-volume theories, which do not contain open string tachyons [156, 154, 158]. In particular in [158] a SU(N) Yang–Mills theory coupled to six real massless scalars has been constructed as world-volume theory of N coincident D3-branes in type 0B theory. In this model the 5-form field strength is not constrained to be self dual and so one can consider both electrically and magnetically charged D3-branes. Moreover in the D3-brane solution the dilaton is not constant, but acquires a spatial (radial) dependence as a consequence of the presence of the $NS\otimes NS$ tachyon. In [158] it is argued that this radial dependence may be related to a dependence of the coupling constant on the energy scale in the gauge theory description. In [159, 160] a field theory with asymptotically free behaviour is constructed from a D3-brane type 0 background with $\mathrm{AdS}_5\times S^5$ geometry.

This very interesting issues will not be further addressed here, in the following the attention will be focused on the correspondence between type IIB string theory in $\mathrm{AdS}_5\times S^5$ and $\mathcal{N}$=4 super Yang–Mills theory.

5.4 Calculations of two- and three-point functions

In this section the prescription for the calculation of correlation functions will be discussed in more detail. First the general idea in the simplest case of scalar fields and for spinors will be reviewed considering two-point functions, then some general considerations on the properties of correlators of R-current operators and some non-renormalization theorems will be reported. Finally the computation of two- and three-point functions of generic chiral primary operators will be discussed.

The computation of two-point functions according to the prescription of the pre-

vious section goes as follows. The Euclidean action for a real massive scalar field in AdS_5 is

$$S = \frac{1}{2}\int d^5z\,\sqrt{g}\left[g^{MN}(\partial_M\Phi)(\partial_N\Phi) + m^2\Phi^2\right]\,. \tag{5.24}$$

As has been explained in [118, 119] in order to compute the two-point Green function of an operator $\mathcal{O}[A]$, related by the correspondence to Φ and coupled to a source J in the boundary theory, in the supergravity approximation valid for large N, one must solve the equation of motion obtained from (5.24) with the condition that $\Phi(x,\rho)$ approach $J(x)$ as $\rho \to 0$. The equation of motion is

$$\frac{1}{\sqrt{g}}\partial_M\left(\sqrt{g}g^{MN}\partial_N\Phi\right) - m^2\Phi = 0\,.$$

Then, after substituting into (5.24) the solution with the prescribed behaviour in the limit $\rho \to 0$, taking the functional derivatives with respect to J one obtains the Green function. The relevant solution is of the form of equation (5.23). As a result one gets for the two-point functions

$$\begin{aligned}\langle\mathcal{O}(x)\mathcal{O}(y)\rangle \;\;=\;\; -\int \frac{d^4x'd\rho}{\rho^5}\;\big[&\partial_M K_\Delta(x',\rho;x,0)\rho^2\partial^M K_\Delta(x',\rho;y,0)+\\ &+m^2K_\Delta(x',\rho;x,0)K_\Delta(x',\rho;y,0)\big]\,.\end{aligned} \tag{5.25}$$

Then integration by parts and use of the equation of motion yields [3]

$$\langle\mathcal{O}(x)\mathcal{O}(y)\rangle = \frac{\Gamma(\Delta+1)}{\pi^2\Gamma(\Delta-2)}\frac{1}{(x-y)^{2\Delta}}\,. \tag{5.26}$$

This computation of two point functions has led Witten to establish the relation (5.20) between the AdS 'mass' m of the field Φ and the conformal dimension Δ of the corresponding operator $\mathcal{O}$. In [119] the same analysis has been carried out for massive vector fields. In general for a p-form field the relation between the AdS 'mass' and the scaling dimension of the dual composite operator becomes $(\Delta-4+p)(\Delta-p) = (mL)^2$, which yields

$$\Delta = 2 \pm \sqrt{4 + (mL)^2 + p(p-4)}\,.$$

The case of the graviton g_{MN}, related to the stress-energy tensor, has been considered in [118, 161].

[3] Again this expression must be modified in the case Δ=2, see the previous section.

Repeating the above construction for a massive spinor in AdS_5 provides the bulk to boundary propagator for fermionic fields [146]. One starts from the action for 'massive' spinors in AdS_5

$$S = \int d^5z \sqrt{g} \left[\overline{\Psi}(\not{D} - m)\Psi\right] , \tag{5.27}$$

which yields the Dirac operator acting on spin-$\frac{1}{2}$ fields in AdS_5

$$\not{D}\Psi = e_{\hat{L}}{}^{M}\gamma^{\hat{L}} \left(\partial_M + \frac{1}{4}\omega_M^{\hat{M}\hat{N}}\gamma_{\hat{M}\hat{N}}\right)\Psi = (\rho\gamma^{\hat{5}}\partial_5 + \rho\gamma^{\hat{\mu}}\partial_\mu - 2\gamma^{\hat{5}})\Psi , \tag{5.28}$$

where $e_{\hat{L}}{}^{M}$ is the vielbein, $\omega_M^{\hat{M}\hat{N}}$ the spin connection (hatted indices refer to the tangent space) and $\gamma^{\hat{\mu}}$ are the four-dimensional Dirac matrices. Just like in the case of scalar fields in order to compute correlation functions one must solve the equations of motion with assigned boundary conditions. However the action (5.27) vanishes for any field configuration which satisfies the equations of motion. This implies that the partition function is actually independent of the boundary condition on the field Ψ, so that it cannot be used as a generating functional. To solve this problem a suitable modification of the action (5.27) by the addition of a boundary term has been proposed in [146]. The uniqueness of such a boundary term has been recently discussed in [162].

The suggestion of [146] is to supplement the action (5.27) by a term of the form

$$S_{\rm b} = \lim_{\epsilon\to 0}\int_{M^\epsilon} d^4x\sqrt{g^\epsilon}\overline{\Psi}\Psi , \tag{5.29}$$

where M^ϵ is a closed four-dimensional submanifold of AdS_5, which approaches the boundary of AdS_5 as $\epsilon \to 0$. In (5.29) g^ϵ is the metric on M^ϵ induced by the one of AdS_5. This boundary term does not modify the equations of motion in the bulk and furthermore is invariant under the isometry group of AdS_5. As a consequence of the vanishing of the bulk action on field configurations satisfying the equations of motion the entire contribution to the partition function, to be employed as generating functional in the AdS/CFT correspondence, comes from the boundary term (5.29).

This allows to construct the bulk to boundary propagator $K^F_\Delta(x', \rho; y, \rho_0)$ for spinors of mass m. For fermionic fields, as discussed in [146], the relation between the AdS 'mass' of a supergravity field and the scaling dimension, Δ, becomes

$$\Delta = 2 - mL .$$

The case of the dilatino, Λ, with AdS mass $m = -\frac{3}{2L}$, corresponding to an operator of dimension $\Delta = \frac{7}{2}$ will be studied in section 5.6.

In the construction of [146] there is a subtlety related to the apparent arbitrariness of the coefficient of the boundary term (5.29) to be added to to the bulk action. This point has been recently clarified in [162]. In the derivation of the equations of motion from a variational principle only the negative chirality part of the boundary limit of the spinor Ψ is fixed. The variation of the positive chirality component of Ψ on the boundary generates the boundary term in the action, which is uniquely fixed by the requirement that the total variation of the action must be zero on the solution of the equations of motion in the class of histories defined by the boundary conditions.

To compute three-point functions one must consider interaction terms in the supergravity action and solve the corresponding equations of motion, with a suitable prescription for the behaviour on the boundary. For instance, in the case of scalar fields, interaction terms of the form

$$\begin{aligned} \mathcal{L}^{(A)} &= \Phi_1\Phi_2\Phi_3 \\ \mathcal{L}^{(B)} &= g^{MN}\Phi_1(\partial_M\Phi_2)(\partial_N\Phi_3) \end{aligned}$$

give the following contributions to three point functions

$$\begin{aligned} \langle \mathcal{O}_1(x)\mathcal{O}_2(y)\mathcal{O}_3(z)\rangle^{(A)} &= -\int \frac{d^4x'd\rho}{\rho^5}\, K_{\Delta_1}(x',\rho;x,0)K_{\Delta_2}(x',\rho;y,0)\cdot \\ &\qquad \cdot K_{\Delta_3}(x',\rho;z,0) \\ \langle \mathcal{O}_1(x)\mathcal{O}_2(y)\mathcal{O}_3(z)\rangle^{(B)} &= -\int \frac{d^4x'd\rho}{\rho^5}\, K_{\Delta_1}(x',\rho;x,0)\partial_M K_{\Delta_2}(x',\rho;y,0)\cdot \\ &\qquad \cdot \rho^2\partial^M K_{\Delta_3}(x',\rho;z,0)\,, \end{aligned} \tag{5.30}$$

where Δ_1, Δ_2 and Δ_3 are the dimensions of the operators coupled to Φ_1, Φ_2 and Φ_3 respectively. Equations (5.30) are obtained, analogously to the expressions for two-point functions, by solving the equations of motion in the presence of the above interaction terms and then substituting back into the action, whose exponential is used as generating functional. To derive (5.30) one must solve the equations recursively as a series in the coupling constant g_s. To generate three-point functions it is sufficient to stop to the lowest order, so that the bulk to boundary propagators, K_{Δ_i} in (5.30) are the same entering two-point functions.

The contributions to three-point functions can be represented through a diagram like the one depicted in figure 5.2, where the derivatives, read from the interaction, eventually act on the bulk to boundary propagators.



Figure 5.2: AdS diagram for the correlator of three scalar operators

By writing explicitly (5.30) one immediately sees that these correlation functions have the correct form required by conformal symmetry. Actually, as will be discussed in the case of R-currents, two-point functions in the boundary conformal field theory are determined by superconformal symmetry up to a constant, so that the comparison with the results obtained from AdS_5 only allows to establish the map between the fields on the two sides of the correspondence and to fix the normalizations. Analogously three-point functions are strongly constrained. Their supergravity calculation in AdS_5 and comparison with field theory only provides a check of the normalization constants (OPE coefficients). A completely non-trivial check of the proposed correspondence requires the computation of four- or higher-point functions, that will be examined in subsequent sections.

The R-symmetry conserved currents of $\mathcal{N}$=4 super Yang–Mills theory were given in equation (2.17). The two-point function $\langle \mathcal{J}^a_\mu \mathcal{J}^b_\nu \rangle$ is completely fixed by superconformal invariance. The dimension of the operators $\mathcal{J}^a_\mu$ is Δ=3 and the form of the correlator of two currents is [145]

$$\langle \mathcal{J}^a_\mu(x) \mathcal{J}^b_\nu(y) \rangle = c \frac{\delta^{ab}}{(2\pi)^4} \left(\delta_{\mu\nu}\Box - \partial_\mu\partial_\nu\right) \frac{1}{(x-y)^4} \,,$$

where c is a positive constant, which is only determined by one-loop corrections thanks to a non-renormalization theorem. c is the central charge of the $\mathcal{J}\mathcal{J}$ OPE and equals the trace anomaly. Since the latter is related by $\mathcal{N}$=1 supersymmetry to the R-current anomaly which is one-loop exact c is determined by the one-loop result as well. In the case of $\mathcal{N}$=4 super Yang–Mills with gauge group SU(N) one finds [145]

$$c = \frac{1}{2}(N^2 - 1) \,.$$

The three point function of currents is also strongly constrained by superconformal symmetry; it contains a normal parity part and an abnormal parity part [145]

$$\begin{aligned}\langle \mathcal{J}^a_\mu(x)\mathcal{J}^b_\nu(y)\mathcal{J}^c_\lambda(z)\rangle &= \langle \mathcal{J}^a_\mu(x)\mathcal{J}^b_\nu(y)\mathcal{J}^c_\lambda(z)\rangle_+ + \langle \mathcal{J}^a_\mu(x)\mathcal{J}^b_\nu(y)\mathcal{J}^c_\lambda(z)\rangle_- = \\ &= f^{abc}\left[k_1^{(+)}D_{\mu\nu\lambda}(x,y,z) + k_2^{(+)}C_{\mu\nu\lambda}(x,y,z)\right] + k^{(-)}d^{abc}M_{\mu\nu\lambda}(x,y,z)\,,\end{aligned}$$

where $D_{\mu\nu\lambda}$, $C_{\mu\nu\lambda}$ and $M_{\mu\nu\lambda}$ are known tensor functions and the f^{abc} and d^{abc} SU(N) symbols are defined by $\mathrm{tr}\left(T^aT^bT^c\right) = \frac{1}{4}(if^{abc}+d^{abc})$. $k_1^{(+)}$, $k_2^{(+)}$ and $k^{(-)}$ are numerical constants; $k_1^{(+)}$ is fixed by the Ward identity connecting two- and three-point functions, which relates it to c by $k_1^{(+)}=\frac{c}{16\pi^2}$, $k_2^{(+)}$ is independent and it is not protected by general non-renormalization theorems. The third constant, $k^{(-)}$, is controlled by the Adler–Badeen theorem which implies that it is one-loop exact. The same non-renormalization properties of the R-currents should of course hold true for the other composite operators in the current multiplet.

The computation of two- and three-point functions of chiral primary operators (CPO's) will now be reviewed following [148, 145]. The chiral primary operators that will be considered are of the form

$$\mathcal{O}^I_\ell = g^\ell_{\mathrm{YM}} t^I_{i_1\ldots i_\ell}\mathrm{tr}\left(\varphi^{i_1}(x)\ldots\varphi^{i_\ell}(x)\right)\,, \tag{5.31}$$

where φ^i are the six real matrix-valued scalar fields of $\mathcal{N}$=4 supersymmetric Yang–Mills theory and $t^I_{i_1\ldots i_\ell}$ is a totally symmetric (traceless) rank ℓ tensor of SU(4)∼SO(6), whose normalization is chosen such that $t^{I_1}_{i_1\ldots i_\ell}t^{I_2\,i_1\ldots i_\ell} = \delta^{I_1I_2}$. The free propagator for the fields φ^i is

$$\langle \varphi^i_a\varphi^j_b\rangle = \frac{\delta^{ij}\delta_{ab}}{(2\pi)^2}\frac{1}{(x-y)^2}\,.$$

Chiral primary operators (5.31) belong to representations of the SU(4) R-symmetry group characterized by Dynkin labels $(0,\ell,0)$, they are Lorentz scalars, *i.e.* they belong to the representation j_1=j_2=0 of the Euclidean Lorentz group SO(4)=SU(2)× SU(2), and their conformal dimension is Δ=ℓ. The class of CPO's comprises the operators $\mathcal{Q}^{ij}$ and in particular $\mathcal{Q}$ and $\overline{\mathcal{Q}}$ considered in section 4.5.

In general operators involved in the AdS/CFT correspondence have a conformal dimension Δ that is not an integer. Only operators belonging to short supermultiplets of the SU(2,2|4) superconformal group, which are dual to the massless AdS supergravity fields and their KK descendants, have a dimension that is protected under renormalization [163]. In supersymmetric field theories there are chiral primary

operators, like hose considered here, whose dimensions are integer and protected, even if they are not conserved currents. In the case of $\mathcal{N}$=4 super Yang–Mills theory under consideration the conformal dimension of a generic operator satisfies the bound

$$\Delta \geq q = \ell \,, \tag{5.32}$$

where q is the charge of the field under a U(1) subgroup of the SU(4) R-symmetry group. (5.32) is a "BPS-type" bound, which follows from the superconformal algebra (2.16).

As already discussed in the previous section the supergravity fields and the Kaluza–Klein states in d=5 $\mathcal{N}$=8 gauged supergravity belong to short multiplets of the superisometry group SU(2,2|4). The same is true for their counterparts in $\mathcal{N}$=4 Super Yang–Mills theory (where SU(2,2|4) is the superconformal group). The CPO's introduced above play a central rôle in the construction of these multiplets. As explained in [163] all these multiplets can be obtained starting from scalar chiral primary operators like those in (5.31) and resorting to analytic superspace. The generic short multiplet is described by a composite twisted chiral superfield [111] of the form

$$\mathcal{W}_{(\ell)} = \mathrm{tr}\left(W^{(i_1}\dots W^{i_\ell)}\right) - \text{ traces}\,,$$

where the superfields $W^i = \frac{1}{2}t^i_{AB}W^{[AB]}$ are the same introduced in chapter 2, equations (2.7) and (2.8).

Massive string states on the contrary are in long multiplets as well as their duals, which are non chiral primary operators. An example of operators of this kind is represented by the $\mathcal{N}$=4 embedding of the Konishi multiplet [164, 163], which, in terms of $\mathcal{N}$=1 superfields, is

$$\Sigma = \Phi^I e^V \Phi^\dagger_I \,, \tag{5.33}$$

where Φ^I, $I = 1, 2, 3$ are the $\mathcal{N}$=1 chiral multiplets and V the $\mathcal{N}$=1 vector multiplet of $\mathcal{N}$=4 super Yang–Mills theory.

In [148] two- and three-point Green functions of CPO's have been computed using type IIB supergravity in AdS_5 and Witten's prescription. The result is

$$\begin{aligned}
\langle \mathcal{O}^{I_1}_\ell(x)\mathcal{O}^{I_2}_\ell(y)\rangle &= \hat{g}^{2\ell}\frac{\ell}{(2\pi)^{2\ell}}\delta^{I_1I_2}\frac{1}{(x-y)^{2\ell}} \\
\langle \mathcal{O}^{I_1}_{\ell_1}(x)\mathcal{O}^{I_2}_{\ell_2}(y)\mathcal{O}^{I_3}_{\ell_3}(z)\rangle &= \frac{\hat{g}^\Sigma}{N}\frac{\ell_1\ell_2\ell_3}{(2\pi)^\Sigma}t^{I_1I_2I_3}\frac{1}{(x-y)^{2\alpha_3}(y-z)^{2\alpha_1}(x-z)^{2\alpha_2}}\,,
\end{aligned}$$

where $\Sigma{=}\ell_1+\ell_2+\ell_3$ and the tensor $t^{I_1I_2I_3}$ is the unique invariant that can be constructed contracting $\alpha_1=\frac{\ell_2+\ell_3-\ell_1}{2}$ indices between $t^{I_2}_{i_1\ldots i_{\ell_2}}$ and $t^{I_3}_{i_1\ldots i_{\ell_3}}$ and $\alpha_2=\frac{\ell_1+\ell_3-\ell_2}{2}$ and $\alpha_3=\frac{\ell_1+\ell_2-\ell_3}{2}$ respectively between the other two pairs. After rescaling the CPO's according to

$$\mathcal{O}^I_\ell \longrightarrow \tilde{\mathcal{O}}^I_\ell = \frac{(2\pi)^\ell}{\hat{g}^\ell\sqrt{\ell}}\mathcal{O}^I_\ell\,,$$

the above correlators are found to agree perfectly with the free field calculation in $\mathcal{N}{=}4$ super Yang–Mills.

This result is expected to be valid only in the large N limit at fixed and large $\hat{g}$, when the Yang–Mills coupling $g_{\rm YM}$ goes to zero. However in [147] it was proved that the one-loop corrections to the above correlators are zero. As a consequence the AdS calculation agrees with the *exact* perturbative contribution to two- and three-point functions of CPO's for arbitrary $g_{\rm YM}$. This strongly suggests that the conjectured correspondence should be valid at finite N as well. In [147] a component formulation in terms of $\mathcal{N}{=}1$ multiplets is employed. The chiral primary operators that are considered are of the form

$$\mathrm{tr}\big(X^\ell\big) \;=\; \mathrm{tr}\left(\varphi^{I_1}\ldots\varphi^{I_\ell}\right) \tag{5.34}$$

$$\mathrm{tr}\big(X^{\dagger\ell}\big) \;=\; \mathrm{tr}\left(\varphi^\dagger_{I_1}\ldots\varphi^\dagger_{I_\ell}\right)\,, \tag{5.35}$$

where φ^I and $\varphi^\dagger_I$, with $I=1,2,3$, are the complex scalar fields belonging to the chiral multiplets. The action of the model in this formulation was given in equation (2.11).

The proof of the vanishing of the one-loop corrections to correlators of CPO's given in [147] is based on a non-renormalization 'theorem' [165] for the two-point function

$\langle \mathrm{tr}\big(X^2\big)\mathrm{tr}\big(X^{\dagger 2}\big)\rangle =$ 

and on clever colour combinatorics. The non-renormalization theorem says that the one loop correction to the above correlator vanishes. This is a consequence of the following relations. The one-loop corrections to $\langle \mathrm{tr}\big(X^2\big)\mathrm{tr}\big(X^{\dagger 2}\big)\rangle$ come from the diagrams



(5.36)



$$= 2c\frac{A(x,y)}{(x-y)^4} \tag{5.37}$$

Notice that in the formulation of (2.11), that is used in [147], the Wess–Zumino gauge is exploited, so that the one loop corrections to the propagator, represented by the bubbles in (5.37), are not zero. Moreover the second term in (5.36) is the contribution of the 'D-term', while the 'F-term' does not enter. Both $B(x,y)$ and $A(x,y)$ are logarithmically divergent (see section 3.1), but one can prove the non-renormalization theorem

$$B(x,y) + 2A(x,y) = 0\,. \tag{5.38}$$

The one-loop correction to the two-point function $\langle \mathrm{tr}\big(X^\ell\big)\mathrm{tr}\big(X^{\dagger\ell}\big)\rangle$



comes from the sum of diagrams of the following types







One must consider the sum of contributions in which the 'two-particle' interaction structures (5.36) and (5.37) are inserted between each pair of lines. In [147] it is shown, through a combinatoric analysis, that this sum leads to a total one-loop correction of the form

$$\langle \mathrm{tr}\big(X^k X^{\dagger k}\big) \sim kN \frac{[B(x,y)+2A(x,y)]}{(x-y)^{2k}}\,,$$

which vanishes thanks to the non-renormalization theorem for the k=2 case.

Analogously one can prove the vanishing of the one-loop correction to the three point function $\langle \mathrm{tr}\big(X^{\ell_1}\big)\mathrm{tr}\big(X^{\ell_2}\big)\mathrm{tr}\big(X^{\dagger\ell_3}\big)\rangle$



To summarize the expressions of two- and three-point functions of CPO's in perturbation theory are given by the free-field result and, as shown in [148], this coincides with what is found in the supergravity calculation in AdS_5. As already remarked supersymmetry allows to generalize this result to all the composite operators of the current multiplet. In [147] this has been explicitly verified for a three-point function involving the operator $\mathcal{E}^{AB}$ of equation (2.18), which is a descendant [4] of CPO's. Notice that in this computation it is crucial to take into account the modification (2.19) of the operator $\mathcal{E}^{AB}$ in the non Abelian case [89].

[4] $\mathcal{E}^{AB}$ is a descendant under superconformal transformations, but a primary field from the viewpoint of conformal symmetry.

5.5 Four-point functions

Unlike two- and three-point functions which, as discussed in the previous section, are completely determined by superconformal symmetry up to constants, four-point functions are only determined up to an arbitrary function of two independent cross-ratios, for instance those introduced in section 4.5

$$r = \frac{x_{12}^2 x_{34}^2}{x_{13}^2 x_{24}^2}\,, \qquad\qquad s = \frac{x_{14}^2 x_{23}^2}{x_{13}^2 x_{24}^2}\,. \tag{5.39}$$

As a consequence the comparison of four-point amplitudes computed in supergravity/superstring theory in AdS_5 with super Yang–Mills correlation functions would represent a truly 'dynamical' and non-trivial check of the correspondence. Such a check has been carried out, in an indirect way, for the non-perturbative correction to a four-point function in [89] using the exact match of a correlation function of sixteen fermionic operators in the same supersymmetry multiplet and will be reviewed in the next section.

The calculation of four-point Green functions of composite operators in $\mathcal{N}$=4 supersymmetric Yang–Mills is rather complicated. The application of $\mathcal{N}$=1 superspace techniques to this problem will be examined later.

On the supergravity/superstring side the computation of four- and higher-point amplitudes with the approach of section 5.4 presents a new difficulty, with respect to the case of two- and three–point functions, related to the existence of two kinds of contributions, associated with "contact" and "exchange" diagrams.



Figure 5.3: Contact 4-point amplitude



Figure 5.4: Exchange 4-point amplitude

The former, see figure 5.3, are completely analogous to the above discussed contributions to two- and three-point functions, while the latter correspond to diagrams like the one in figure 5.4 and, as a new feature, involve also *bulk to bulk propagators* in

AdS_5. The diagram of figure 5.4 represents an s-channel amplitude, analogously one must take into account t and u channels. Moreover one must consider the possible exchange of all the states, for finite N also string states, coupled to the external vertices in the supergravity/superstring action. The corresponding amplitude has the form

$$A(x_1,x_2,x_3,x_4)=\int \frac{d^4x'd\rho'}{\rho'^5}\frac{d^4x''d\rho''}{\rho''^5}\ [K(x',\rho';x_1,0)\cdot \\ \cdot K(x',\rho';x_2,0)G(x',\rho';x'',\rho'')K(x'',\rho'';x_3,0)K(x'',\rho'';x_4,0)]\ ,$$

where $G(z';z'')$ denotes the bulk to bulk propagator and, for compactness of notation, indices and possible derivatives acting on the various lines, depending the explicit form of the trilinear couplings, have been suppressed.

The first computations of this kind have been presented in [166, 106] for scalar four-point functions. The amplitudes considered are

$$\begin{aligned} &A_{\phi\phi\phi\phi}(x_1,x_2,x_3,x_4)\\ &A_{\phi\chi\phi\chi}(x_1,x_2,x_3,x_4)\\ &A_{\chi\chi\chi\chi}(x_1,x_2,x_3,x_4)\,, \end{aligned} \tag{5.40}$$

where ϕ and $\chi{=}C^{(0)}$ are the the two massless scalars of the type IIB superstring, the dilaton and the axion respectively. The operator corresponding to ϕ is the "glueball field" $\mathcal{G}=\mathrm{tr}\,(F^2)$, whereas χ corresponds to $\mathcal{C}=\mathrm{tr}\left(F\tilde{F}\right)$, so that the amplitudes (5.40) should give the Green functions

$$\begin{aligned} &\langle\mathcal{G}(x_1)\mathcal{G}(x_2)\mathcal{G}(x_3)\mathcal{G}(x_4)\rangle\\ &\langle\mathcal{G}(x_1)\mathcal{C}(x_2)\mathcal{G}(x_3)\mathcal{C}(x_4)\rangle\\ &\langle\mathcal{C}(x_1)\mathcal{C}(x_2)\mathcal{C}(x_3)\mathcal{C}(x_4)\rangle \end{aligned} \tag{5.41}$$

in $\mathcal{N}{=}4$ super Yang–Mills.

The relevant part of the type IIB supergravity action in AdS_5 is

$$S=\frac{1}{2\kappa_0^2}\int d^5z\sqrt{g}\left[R-\frac{1}{2}\left(\partial\phi\right)^2-\frac{1}{2}e^{2\phi}\left(\partial\chi\right)^2\right]\,. \tag{5.42}$$

Notice that since the scalar fields ϕ and χ correspond to zeroth-order spherical harmonics, one can forget about the S^5 factor and consider $\phi=\phi(z)$ and $\chi=\chi(z)$ as depending only on the AdS_5 coordinate.

In the case of the $A_{\phi\phi\phi\phi}$ amplitude the only exchange diagrams that contribute involve the graviton propagator and at this order there are no contact terms, in the $A_{\phi\chi\phi\chi}$ amplitude both the graviton and the axion propagators enter, while $A_{\chi\chi\chi\chi}$ receives contribution from the exchange of gravitons and dilatons. Furthermore contact diagrams contribute to $A_{\phi\chi\phi\chi}$ and $A_{\chi\chi\chi\chi}$.

The fundamental difficulty in the computation of the above amplitudes is the construction of the propagators in the AdS_5 space. An interesting result has been proved in [106]. It has been shown that the graphs corresponding to the χ-exchange in the amplitude $A_{\phi\chi\phi\chi}$ actually coincide with the contact graph. This is a consequence of the fact that the scalar bulk to bulk propagator $G(z;z')$ satisfies an equation of the form

$$\Delta_z G(z;z') = \delta(z-z')$$

in AdS_5. Integrating by parts in the exchange amplitude, one obtains exactly a factor $\Delta_z G(z;z')$ and hence one of the two integrations over AdS_5 can be performed using the δ-function. However this result does not hold in general and so the knowledge of the propagators for all the relevant fields in AdS_5 is necessary.

The AdS_5 propagator for scalar fields has been computed in [167], while the propagator for gauge fields was given in [168]. The graviton propagator in AdS_5 has been computed only very recently [169].

As already remarked in section 5.1 the type IIB supergravity action contains terms that receive also non-perturbative corrections from D-instantons. In particular D-instantons contribute to the $\mathcal{R}^4$ term (5.7) and to the Λ^{16} term (5.8). As argued in [170, 89] the corresponding contributions to supergravity/superstring amplitudes in AdS_5 should be compared with instanton contributions to the correlation functions of the associated composite operators in $\mathcal{N}$=4 supersymmetric Yang–Mills theory. This comparison has been achieved in [89] for the correlation functions presented in the previous chapter (see sections 4.5 and 4.6). The results of [89] will be reviewed in the following sections. Notice that for the non-perturbative contribution to scattering amplitudes, with the "minimal" number of insertions leading to a non-vanishing result, *only contact terms are relevant* [5]. As a consequence it is easy to single out the contributions to be compared with the Yang–Mills computations. In particular notice that there is no ambiguity related to possible cancellations among contact and

[5] Two- and three-point functions do not receive non-perturbative corrections at lowest order.

exchange integrals. Moreover in general the contact non-perturbative contribution to an amplitude has schematically the form

$$\int \frac{d^4x d\rho}{\rho^5} K_{\Delta_1} K_{\Delta_2} \dots K_{\Delta_n} ,$$

which exactly coincides with the form expected for an instanton contribution in $\mathcal{N}$=4 Yang–Mills theory after the integrations over the fermionic collective coordinates have been performed, since the one-instanton moduli space is exactly AdS_5 [89, 90, 91, 92] and the instanton 'profile' $\mathrm{tr}\left(\overline{F}^2_{\mu\nu}\right)$ is identical to the bulk to boundary propagator K_Δ for $\Delta = 4$. This implies that the comparison with the Yang–Mills calculation can be performed *before* computing the integrals. These considerations will be exploited in the next section, where specific examples will be considered.

Some remarks on the computation of perturbative contributions to four-point Green functions will now be made. The calculation of correlation functions of composite operators in $\mathcal{N}$=4 super Yang–Mills theory can be simplified using superspace techniques. Although the explicit calculations have not yet been completed the application of this method to chiral primary operators will be briefly sketched.

The approach followed in [147] and reviewed in the previous section for the computation of three-point functions of CPO's can be generalized to correlation functions of $\mathcal{N}$=1 superfields whose lowest components are chiral primary operators of the form of equation (5.35). The superfields to be considered are of the form

$$\mathrm{tr}\left(\Phi^{I\,\ell}\right) , \qquad \mathrm{tr}\left(\Phi_I^{\dagger\,\ell}\right) , \qquad I = 1, 2, 3 . \tag{5.43}$$

The simplest case, ℓ=2, will be discussed here. Exploiting the SU(3) symmetry, that is manifest in the $\mathcal{N}$=1 formulation, one can restrict to Green functions with fixed values of the indices I, which allows to simplify the calculation. The chiral primary operators with $\ell = 2$ are the scalars $\mathcal{Q}^{ij}$ in (2.18). The operators $\mathcal{Q}^{ij}$ belong to the $\mathbf{20}_{\mathbb{R}}$ of the SU(4) R-symmetry group, which decomposes with respect to the SU(3)×U(1) subgroup that is manifest in the $\mathcal{N}$=1 formulation according to

$$\mathbf{20}_{\mathbb{R}} \supset \overline{\mathbf{6}}_{-\frac{4}{3}} + \mathbf{6}_{\frac{4}{3}} + \mathbf{8}_0 , \tag{5.44}$$

where the subscript on the right hand side refers to the U(1) charge. The corresponding decomposition of $\mathcal{Q}^{ij}$ reads

$$\begin{aligned} &\mathcal{O}^{IJ}_{\mathbf{6}} = \mathrm{tr}\left(\Phi^I \Phi^J\right) \in \mathbf{6} , \qquad \mathcal{O}_{\overline{\mathbf{6}}\,IJ} = \mathrm{tr}\left(\Phi^\dagger_I \Phi^\dagger_J\right) \in \overline{\mathbf{6}} , \\ &\mathcal{O}^I_{\mathbf{8}\,J} = \mathrm{tr}\left(\Phi^I e^V \Phi^\dagger_J - \frac{\delta^I{}_J}{3} \Phi^K e^V \Phi^\dagger_K\right) \in \mathbf{8} . \end{aligned} \tag{5.45}$$

For the sake of simplicity the attention will be restricted to Green functions of operators $\mathcal{O}^{IJ}_{\mathbf{6}}$ and $\mathcal{O}_{\overline{\mathbf{6}}\,IJ}$.

Hence one type of four-point functions that one must consider is of the form

$$G^{(a)}_I = \langle \mathrm{tr}\left(\Phi^{\dagger\,2}_I\right) \mathrm{tr}\left(\Phi^{I\,2}\right) \mathrm{tr}\left(\Phi^{\dagger\,2}_I\right) \mathrm{tr}\left(\Phi^{I\,2}\right)\rangle \,,$$

with fixed I. There are two different tree-level contributions, corresponding to the diagrams (the notations are the same as in chapter 3)



The connected diagram is proportional to N^2 while the disconnected one goes like N^4. The only non-vanishing corrections to this Green function come from the diagrams



plus the analogous diagrams in which the VV internal propagator is inserted between different pairs of external lines. Notice in particular that the following correction,

obtained from the disconnected diagram,



is trivially zero because of colour contractions. Moreover working in the superfield formalism and choosing the Fermi–Feynman gauge, there are no corrections to the propagators to be included. Also, because of the non-renormalization properties of two-point functions discussed in the previous section, there is no non-vanishing disconnected diagram contributing to the four-point function at first or higher order.

A second kind of four-point functions is of the form

$$G^{(b)}_{IJ} = \langle \mathrm{tr}\left(\Phi_I^{\dagger\,2}\right) \mathrm{tr}\left(\Phi^{I\,2}\right) \mathrm{tr}\left(\Phi_J^{\dagger\,2}\right) \mathrm{tr}\left(\Phi^{J\,2}\right)\rangle\,,$$

with $I \neq J$. In this case the only tree-level diagram is disconnected



and goes as N^4. There is a unique non vanishing first order correction corresponding

to the connected diagram



where the internal propagator corresponds to the exchange of a chiral superfield with index $K \neq I \neq J$. Just like in the previous case there is no disconnected first order correction because of the non-renormalization properties of two-point functions.

The previous diagrams can be evaluated using the techniques of chapter 3. Since the points where the composite operators are inserted are external and not interaction points the result will not be local in the fermionic variables of $\mathcal{N}$=1 superspace. This is not in agreement with the results of a calculation presented in [171].

An analysis similar to the one described here has been carried out in [172] employing a formulation of the model in $\mathcal{N}$=2 harmonic superspace. In [172] four-point functions are explicitly computed in terms of polylogarithm functions. The expressions obtained resemble the one that was obtained in section 4.5 from the instanton contribution to the correlator $\langle \mathcal{Q}\mathcal{Q}\overline{\mathcal{Q}\mathcal{Q}}\rangle$.

It has been observed by various authors [166, 106, 109, 107, 108, 173, 174] that the four-point functions computed in the AdS/CFT correspondence develop logarithmic singularities, when the limit of two coincident points is considered. This kind of behaviour has already been pointed out in section 4.5. In all the papers [166, 106, 109, 107, 173, 174] the short-distance behaviour has been shown taking the limit in the unintegrated expression obtained from the supergravity amplitude in AdS_5. This is rather subtle because the exchange of limit and integration might be problematic, moreover, as observed in [106], there could be compensations among various AdS diagrams. However this kind of cancellations cannot take place in the cases considered in [107, 108]. In particular in [107] some non-perturbative contributions from D-instantons to the four-point functions of scalar operators corresponding the dilaton and axion fields has been considered. The results of section 4.5, where the instanton contribution to the four-point function $G_{\mathcal{Q}^4}$ has been given explicitly, unconfutably show the logarithmic behaviour. A satisfactory explanation of the origin of these

singularities as well as of the reason for the lack of the expected pole-type singularities has not been proposed yet. In [109] an interesting analysis relating the contribution of contact and exchange amplitudes in AdS to the OPE of the corresponding operators in $\mathcal{N}$=4 super Yang–Mills has been given, using the technique of 'conformal partial wave expansion', but the possible cancellation of logarithms between the two types of contributions has not been excluded.

From the point of view of the boundary conformal field theory the central point appears to be the presence of anomalous dimensions for some operators that does not violate the conformal symmetry. This issue is under active investigation [110] and will be briefly addressed in the concluding remarks. Notice however that the behaviour of the correlation function $G_{\mathcal{Q}^4}$ for coincident points does not violate the non-renormalization theorems for three-point functions of the previous section, so that such non-renormalization properties could be generalizable to the non-perturbative level, at least for protected operators. In fact the limit in which two $\mathcal{Q}$ or two $\overline{\mathcal{Q}}$ operators are taken to coincide in $G_{\mathcal{Q}^4}$, that would correspond to one of the three-point functions that are expected to be tree-level exact, gives a vanishing result thanks to the pre-factor in equation (4.43).

5.6 D-instanton effects in d=10 type IIB superstring theory

As already discussed the low energy type IIB supergravity action contains a $f_4(\tau,\overline{\tau})\mathcal{R}^4$ term as well as a supersymmetry related $f_{16}(\tau,\overline{\tau})\Lambda^{16}$ term. Both of these terms receive contributions from D-instantons of the type IIB theory.

The AdS/CFT correspondence described in the previous section allows to associate to these terms well defined contributions to correlation functions of composite operators in the boundary Yang–Mills theory. It is natural to expect that the D-instanton generated terms should be related to instanton effects in $\mathcal{N}$=4 super Yang–Mills.

The type IIB D-instantons are Dirichlet $p=-1$ branes, *i.e.* they are localized objects in ten-dimensional space-time, which means that they have a zero-dimensional world-volume. The charge-K D-instanton in flat ten-dimensional space-time is a finite-action solution of the classical Euclidean equations of motion in which the metric (in the Einstein frame) is flat and the complex scalar, $\tau=\chi+ie^{-\phi}$, has a non-trivial profile with a singularity at the position of the D-instanton, whereas the

other fields vanish [175]. Moreover the D-instanton is a BPS saturated configuration preserving half of the supersymmetries. In the following the constant expectation values of the dilaton and the axion in the zero instanton sector will be denoted by $\tilde{\phi}$ and $\tilde{\chi}$ respectively, while hatted symbols will refer to the classical D-instanton solution.

To compute the scalar part of the type IIB action in the D-instanton background it is useful to dualize the $R \otimes R$ scalar, $\chi = C^{(0)}$, to an eight-form potential, $C^{(8)}$. Since the metric is trivial, the action (up to constants) can be put in the form

$$\begin{aligned} S &= \frac{1}{4}\int \left(d\phi \pm e^{-\phi} * dC^{(8)}\right) \wedge * \left(d\phi \pm e^{-\phi} * dC^{(8)}\right) \pm \frac{1}{2}\int e^{-\phi} d\phi \wedge dC^{(8)} = \\ &= \frac{1}{4}\int \left(d\phi \pm e^{-\phi} * dC^{(8)}\right) \wedge * \left(d\phi \pm e^{-\phi} * dC^{(8)}\right) \mp \frac{1}{2}\int_{\partial M} e^{-\phi} dC^{(8)}\,, \end{aligned} \tag{5.46}$$

where ∂M denotes the boundary of space-time, that in the presence of the D-instanton is understood as made of the sphere at infinity and an infinitesimal sphere around the location of the instanton. The first term in (5.46) is semi-positive definite, so that it follows that the action satisfies a Bogomol'nyi bound

$$S \geq \frac{1}{2}\left|\int_{\partial M} e^{-\phi} dC^{(8)}\right|\,. \tag{5.47}$$

Minimizing the action yields a BPS condition relating ϕ and $C^{(8)}$, which, expressed in terms of χ, becomes [175]

$$\partial_\Sigma \chi = \pm i \partial_\Sigma e^{-\phi}\,. \tag{5.48}$$

The D-instanton solution is constructed through the ansatz

$$\chi = \tilde{\chi} + i f(r)\,,$$

where $r = |X - X_0|$ and $f(r) \to 0$, as $r \to \infty$. After imposing (5.48) the equation for the dilaton outside a sphere centered at the D-instanton location becomes

$$e^{-\phi} g^{\Lambda\Sigma} \nabla_\Lambda \nabla_\Sigma e^{\phi} = 0\,. \tag{5.49}$$

The solution is the harmonic function [175]

$$e^{\hat{\phi}^{(10)}} = g_s + \frac{3K\alpha'^4}{\pi^4 |X - X_0|^8}\,. \tag{5.50}$$

where X^Λ is the ten-dimensional coordinate and X_0^Λ is the location of the D-instanton. In (5.50) g_s is the asymptotic value of the string coupling and the normalization of the second term has a quantized value [175] by virtue of a condition analogous to the Dirac–Nepomechie–Teitelboim condition that quantizes the charge of an electrically charged p-brane and of its magnetically charged p'-brane dual [176]. It is notable that the solution in (5.50) is simply the Green function for a scalar field to propagate from X_0 to X subject to the boundary condition that $e^\phi = g_s$ at $|X| \to \infty$ or $|X_0| \to \infty$. The solution for χ follows from (5.48) and is of the form

$$\chi = \tilde{\chi} + i\left(A - \frac{1}{g_s} + e^{-\hat{\phi}}\right) , \tag{5.51}$$

where A is an arbitrary constant.

The action for a single D-instanton receives contribution only from the boundary term in (5.46), *i.e.* from the integration over the sphere at infinity and over an infinitesimal sphere around X_0. The arbitrariness in the constant A can be used to simplify the computation. In particular with the choice A=0 in (5.51) the entire D-instanton action comes from the boundary of the infinitesimal sphere. Substituting for f from (5.51) gives

$$S_K = \frac{2\pi|K|}{g_s} . \tag{5.52}$$

On the other hand, with the choice $A = \frac{1}{g_s}$ in (5.51) the expression (5.46) reduces to an integral over the boundary at infinity, but the total action remains the same as S_K in (5.52).

The bosonic zero-modes associated with the D-instanton solution are parameterized by the coordinates X_0^Λ of the center of the D-instanton. As already remarked the D-instanton background breaks half of the supersymmetries. The fermionic zero-modes can be generated by acting with the broken supersymmetries on the scalar solution.

The D-instanton contributions to the effective action of type IIB supergravity can be deduced by studying scattering amplitudes in the instanton background. In this way one obtains in particular the corrections to the $\mathcal{R}^4$ and Λ^{16} terms in equations (5.7) and (5.8) [114, 130].

In [170] it was pointed out that the $\mathcal{R}^4$ term vanishes when computed in the $\mathrm{AdS}_5 \times S^5$ background, because it involves a fourth power of the (vanishing) Weyl tensor. The first, second and third functional derivatives of $\mathcal{R}^4$ vanish as well, and as

a result one finds no corrections to zero, one, two and three-point amplitudes when the $\mathcal{R}^4$ term acts as source for Yang–Mills operators on the boundary. However, there is a non-zero four-graviton amplitude arising from this term. The boundary values of these gravitons are sources for various bosonic components of the Yang–Mills current supermultiplet. For example, the components of the metric in the AdS_5 directions couple to the stress tensor, $\mathcal{T}^{\mu\nu}$, whereas the traceless components polarized in the S^5 directions couple to massive Kaluza-Klein states. A linear combination of the trace of the metric on S^5 and the fluctuation of the $R \otimes R$ four-form potential couples to the scalar components, $\mathcal{Q}^{ij}$. As a consequence the instanton contribution to the four-point correlation function $G_{\mathcal{Q}^4}$ computed in the previous chapter should be related to the appropriate D-instanton correction extracted from $f_4(\tau,\overline{\tau})\mathcal{R}^4$. However it is extremely laborious to single out the exact terms to be compared with the Yang–Mills computation, so that it will be convenient to show the matching of the D-instanton correction to the Λ^{16} term with the $G_{\hat{\Lambda}^{16}}$ Green function of section 4.6.

All the other terms of the same dimension, that are related to $\mathcal{R}^4$ by supersymmetry can be picked out, for example, associating the physical fields with the components of an on-shell type IIB superfield [177]. As already remarked included among these is the sixteen-fermion interaction (5.8). Supersymmetry arguments [178, 179] imply that the functions f_{16} and f_4 are related by the action of SL(2,$\mathbb{Z}$) modular covariant derivatives

$$f_{16}(\tau,\overline{\tau}) = (\tau_2\mathcal{D})^{12} f_4(\tau,\overline{\tau})\,, \tag{5.53}$$

where $\mathcal{D}=\frac{\partial}{\partial\tau}+2i\frac{q}{\tau_2}$ is the covariant derivative that maps a (q,p) modular form into a $(q+2,p)$ form (where the notation (q,p) labels the holomorphic and anti-holomorphic SL(2,$\mathbb{Z}$) weights of the form). Whereas f_4 transforms with modular weight $(0,0)$, the function f_{16} has weight $(12,-12)$ and therefore transforms with a non-trivial phase under SL(2,$\mathbb{Z}$). This is precisely the phase required to compensate for the anomalous $U(1)_B$ transformation of the 16 Λ's, so that the full expression (5.8) is invariant under the SL(2,$\mathbb{Z}$) S-duality group.

In the small coupling limit, all of these terms can be expanded in the form

$$\begin{aligned} e^{-\phi/2} f_n \;&=\; a_n\zeta(3)e^{-2\phi} + b_n + \\ &+ \sum_{K=1}^{\infty} \mu_K (Ke^{-\phi})^{n-7/2} e^{2\pi i K\tau} \left(1 + \sum_{k=1}^{\infty} c_{k,n}^K (Ke^{-\phi})^{-k}\right)\,. \end{aligned} \tag{5.54}$$

The first two terms have the form of string tree-level and one-loop contributions,

while the last one has the appropriate weight of $e^{2\pi i K\tau}$ to be associated with a charge-K D-instanton effect. Anti-D-instanton contributions have not been displayed. The coefficients $c^K_{k,n}$ as well as the 'D-instanton partition functions' μ_K are explicitly given in [130]. The coefficient of the leading term in the series of perturbative fluctuations around a charge-K D-instanton is independent of which particular interaction term is being discussed and reads

$$\mathcal{Z}_K = \mu_K (K e^{-\phi})^{-7/2} e^{2\pi i K\tau} \, .$$

It should be identified with the contribution of a charge-K D-instanton to the measure in string frame which, up to an overall numerical factor, reads

$$d\Omega^{(s)}_K = (\alpha')^{-1} d^{10}X \, d^{16}\Theta \, \mathcal{Z}_K \, . \tag{5.55}$$

Actually there are subtleties in doing this, since the full series is not convergent, it is rather an asymptotic approximation to a Bessel function.

5.6.1 Testing the AdS/CFT correspondence: type IIB D-instantons vs. Yang–Mills instantons

As observed in [170] the charge-K D-instanton action that appears in the exponent in (5.6) coincides with the action of a charge-K Yang–Mills instanton in the boundary theory, which indicates a correspondence between these sources of non-perturbative effects. This idea is reinforced by the correspondence between other factors. For example, after substituting $e^{\phi} = \frac{g^2_{\rm YM}}{4\pi}$ and $\alpha' = L^2 N^{-1/2} g^{-1}_{\rm YM}$, the measure (5.55) contains an overall factor of the coupling constant in the form $g^8_{\rm YM}$. Indeed, this is exactly the power expected on the basis of the AdS/CFT correspondence, since the one-instanton contribution to the Green functions in the $\mathcal{N} = 4$ Yang–Mills theory, considered in chapter 4, also has a factor of $g^8_{\rm YM}$ arising from the combination of the bosonic and fermionic zero-mode norms. In this section the leading instanton contributions to type IIB superstring amplitudes will be compared with the corresponding $\mathcal{N}$=4 current correlators considered in chapter 4. In the next section the same analysis will be carried out studying the semiclassical fluctuations around the $\mathrm{AdS}_5 \times S^5$ D-instanton solution.

To compare the Yang–Mills instanton calculation with an amplitude obtained according to Maldacena's prescription one must expand the function $f_{16}(\tau, \overline{\tau})$ in (5.53)

to extract the one-instanton term. In order to compare with the Yang–Mills sixteen-point correlator one needs to consider the situation in which all sixteen fermions propagate to well defined configurations on the boundary.

Equation (5.28) leads to the normalized bulk-to-boundary propagator of the fermionic field Λ of AdS 'mass' $m = -\frac{3}{2L}$, associated to the composite operator $\hat{\Lambda}$ of dimension $\Delta = \frac{7}{2}$, see equation (2.18),

$$K^{\rm F}_{7/2}(\rho_0, x_0; x) = K_4(\rho_0, x_0; x)\frac{1}{\sqrt{\rho_0}}\left(\rho_0\gamma_{\hat{5}} + (x_0 - x)^\mu\gamma_{\hat{\mu}}\right) . \tag{5.56}$$

Suppressing all spinor indices, one obtains

$$\Lambda_J(x_0, \rho_0) = \int d^4x K^{\rm F}_{7/2}(\rho_0, x_0; x) J_\Lambda(x) , \tag{5.57}$$

where $J_\Lambda(x)$ is a left-handed boundary value of Λ and acts as the source for the composite operator $\hat{\Lambda}$ in the boundary $\mathcal{N}$=4 Yang–Mills theory. As a result, the classical action for the operator $(\Lambda)^{16}$ in the type IIB supergravity on $\mathrm{AdS}_5 \times S^5$ is

$$\begin{aligned} S_\Lambda[J] = {} & e^{-2\pi(\frac{1}{g_s}+i\chi)} g_s^{-12} V_{S^5} \int \frac{d^4x_0 d\rho_0}{\rho_0^5} \\ & t_{16}\prod_{p=1}^{16}\left[K_4(\rho_0, x_0; x_p)\frac{1}{\sqrt{\rho_0}}\left(\rho_0\gamma^{\hat{5}} + (x_0 - x_p)^\mu\gamma_{\hat{\mu}}\right) J_\Lambda(x_p)\right] , \end{aligned} \tag{5.58}$$

where $e^\phi = g_s$ and $\chi = \tilde{\chi}$ (since the scalar fields are taken to be constant in the $\mathrm{AdS}_5 \times S^5$ background) and $V_{S^5} = \pi^3$ is the S^5 volume. The 16-index invariant tensor t_{16} is the same as the one defined after (4.55). The overall power of the coupling constant comes from the factor of $(e^{-\phi})^{-\frac{25}{2}} = \left(\frac{1}{g_s}\right)^{\frac{25}{2}}$ in the expansion (5.54) and the factor of $\frac{1}{\alpha'}$ in front of the Λ^{16} term in the type IIB action, which using the relation (5.13) yields

$$\frac{1}{\alpha'}\left(\frac{1}{g_s}\right)^{\frac{25}{2}} \sim \left(\frac{1}{g_s}\right)^{12} ,$$

where an overall numerical constant has been dropped. Using the dictionary (5.15) and differentiating with respect to the chiral sources this result agrees with the expression (4.55) of section 4.6 obtained in the Yang Mills calculation. In particular the overall power of $g_{\rm YM}$ in (4.55) is reproduced after introducing the same rescaling of each Λ in (5.58) by a factor of $4\pi g_s$ as in equation (4.53). Actually the overall numerical constant has not been checked, see however [92]. Notice that to match

the result of the Yang–Mills calculation it is crucial that in the product of sixteen matrices $K^{\rm F}_{7/2}$ exactly eight factors of $\sqrt{\rho_0}\gamma^5$ and eight factors of $\frac{(x-x_0)_\mu}{\sqrt{\rho_0}}\gamma^\mu$ are picked out. This is ensured by the contraction with t_{16}, which selects the correct number of factors of each kind. In the next section this expression will also be motivated directly by semi-classical quantization around a D-instanton field configuration that is a Euclidean solution of the type IIB supergravity in $\mathrm{AdS}_5 \times S^5$.

The agreement of the Λ^{16} amplitude with the corresponding sixteen-current correlation function of chapter 4 is sufficient to guarantee that the instanton contributions to all the other Yang–Mills correlation functions, that are related to this term by $\mathcal{N}$=4 supersymmetry, will also agree with their type IIB superstring counterparts. For example, the correlation function that has been considered in most detail in chapter 4 was the one with four superconformal scalar currents $\mathcal{Q}^{ij}$ which are in the $\mathbf{20}_{\mathbb{R}}$ of SU(4), and have dimension Δ=2 and AdS 'mass' $m^2 = -\frac{4}{L^2}$. As already discussed the supergravity field, Q^{ij}, that couples to $\mathcal{Q}^{ij}$ is a linear combination of the fluctuation of the trace of the metric on S^5, $h_m{}^m$, and of the four-form field potential, $C^{(4)}_{MNPQ} = \epsilon_{MNPQR}\nabla^R f$. Therefore, contributions to the correlation of four of these composite scalars in the one D-instanton background should correspond to the leading parts of the K=1 terms in the expansion of the $\mathcal{R}^4$ interaction (5.7) as well as terms of the form $\mathcal{R}^2\left(\nabla F^{(5)}\right)^2$ and $\left(\nabla F^{(5)}\right)^4$. These last two terms involve $F^{(5)}$, the self-dual field strength of the antisymmetric four-form potential, and are related by supersymmetry to the $\mathcal{R}^4$ term.

It follows from the structure of (4.40) that the Yang–Mills instanton contribution to each factor of $\mathcal{Q}^{ij}$ is of the form $\nabla\nabla K_2(x^\mu_0, \rho_0; x^\mu, 0)$, where the two derivatives are not necessarily contracted. But this is the expected form for a propagator from the $\mathrm{AdS}_5 \times S^5$ bulk to the boundary for a scalar field of dimension 2. Therefore, at least the general form of the expression obtained from the K=1 terms in the expansion of $\mathcal{R}^4$ and the related $F^{(5)}$ interactions agrees with the four-$\mathcal{Q}^{ij}$ correlation function in a Yang–Mills instanton background. In order to see this agreement in more detail it would first be necessary to determine the precise form of the $\left(\nabla F^{(5)}\right)^2$ and $\left(\nabla F^{(5)}\right)^4$ interactions that contain the fluctuations of $F^{(5)}$. The K=1 contribution to the amplitude of four fluctuations of the appropriate combination of $h_m{}^m$ and f can then be extracted from these interaction terms. Although this computation has not been performed explicitly, the result is guaranteed to reproduce the K=1 expression obtained from the Yang–Mills theory in chapter 4 since it is related by supersymmetry to the Λ^{16} amplitude.

The analogous comparison of the correlation function of eight $\mathcal{E}^{AB}$'s with the amplitude for eight E^{AB}'s in the type IIB superstring theory proceeds in much the same way. The supergravity fields, E^{AB}, that couple to $\mathcal{E}^{AB}$ arise from the internal components of the complex type IIB antisymmetric tensor. Supersymmetry relates an H^8 term, where H is the complex combination of the type IIB three-form field-strengths, H_1 and H_2, introduced in (5.6), to the $\mathcal{R}^4$ term in the type IIB effective action. One thus expects interactions schematically of the form $(\nabla K_3)^8$, with various contractions of the derivatives. Using the explicit form of the spinor collective coordinates $\zeta^A = (\rho_0\eta^A + (x-x_0)_\mu\sigma^\mu\bar{\xi}^A)/\sqrt{\rho_0}$ one may check that the gaugino bilinears $\mathcal{E}^{AB}$ in the one-instanton background exactly give rise to K_3 and derivatives thereof, as expected for the scalar propagator of a scalar field of dimension Δ=3 and AdS mass $m^2 = -\frac{1}{L^2}$. Although a precise matching of the resulting amplitudes has not been performed, one may appeal to supersymmetry arguments to determine the complete structure of these terms.

Notice that the above considered non-perturbative terms in the type IIB supergravity effective action when expanded around the $\mathrm{AdS}_5 \times S^5$ background give rise to both derivative and non-derivative interaction terms. The matching of the non-derivative "mass-related" terms considered here with corresponding terms in the Yang–Mills Green functions is rather straightforward, but clearly it is only a hint to the conjectured correspondence.

5.7 The D-instanton solution in $\mathbf{AdS}_5 \times S^5$

The results of the previous section will now be reinforced by studying how the $(\alpha')^{-1}$ terms in the type IIB effective action can be determined by semi-classical type IIB supergravity field theory calculation in a D-instanton background.

The solution of the equations of motion of the type IIB theory in Euclidean $\mathrm{AdS}_5 \times S^5$ will be now discussed. The BPS condition for a D-instanton in this background is identical to (5.48) and together with the equation of motion for the dilaton leads to

$$g^{\Lambda\Sigma}\nabla_\Lambda\nabla_\Sigma e^\phi = 0 \tag{5.59}$$

and determines the complex scalar τ. In the Einstein frame the Einstein equations are unaltered by the presence of the D-instanton, because the associated Euclidean stress energy tensor vanishes. As a consequence $\mathrm{AdS}_5 \times S^5$ remains a solution. Equation

(5.59) is identical to the equation for the Green function of a massless scalar propagating between the location of the D-instanton (x_0^μ, y_0^i) and the point (x^μ, y^i), which is the bulk-to-bulk propagator (subject to the boundary condition that it is constant in the limits $\rho \to 0$ and $\rho \to \infty$). This is easy to solve using the conformal flatness of $\mathrm{AdS}_5 \times S^5$. In Cartesian coordinates $X^\Lambda = (x^\mu, y^i)$, (5.14) reads $ds^2 = L^2 \rho^{-2} dX \cdot dX$, where $\rho^2 = y \cdot y$ and the solution of (5.59) with a constant asymptotic behavior is of the form

$$e^{\hat{\phi}} = g_s + \frac{\rho_0^4 \rho^4}{L^8} \left(e^{\hat{\phi}^{(10)}} - g_s \right) , \tag{5.60}$$

where $\rho_0 = |y_0|$ and $e^{\hat{\phi}^{(10)}}$ is the harmonic function that appeared in the flat ten-dimensional case, (5.50). In evaluating D-instanton dominated amplitudes one is only interested in the case in which the point (x^μ, y^i) approaches the boundary ($\rho \equiv |y| \to 0$), in which case it is necessary to rescale the dilaton profile (just as it is necessary to rescale the scalar bulk-to-bulk propagator, [118, 119]), so that the combination

$$\rho^{-4} \left(e^{\hat{\phi}} - g_s \right) = \frac{3K(\alpha')^4}{L^8 \pi^4} \frac{\rho_0^4}{((x - x_0)^2 + \rho_0^2)^4} , \tag{5.61}$$

will be of relevance in the $\rho \to 0$ limit.

As mentioned earlier, the correspondence with the Yang–Mills instanton follows from the fact that $\rho_0^4/((x - x_0)^2 + \rho_0^2)^4 = K_4$ is proportional to the instanton number density, $(\overline{F}_{\mu\nu})^2$, in the $\mathcal{N}$=4 Yang–Mills theory. Strikingly, the scale size of the Yang–Mills instanton is replaced by the distance ρ_0 of the D-instanton from the boundary. This is another indication of how the geometry of the Yang–Mills theory is encoded in the type IIB superstring. Note, in particular, that as the D-instanton approaches the boundary $\rho_0 \to 0$, the expression for $\rho^{-4} e^{\hat{\phi}}$ reduces to a δ function that corresponds to a zero-size Yang–Mills instanton.

The BPS condition implies that one can write the solution for the $R \otimes R$ scalar as

$$\hat{\chi} = \tilde{\chi} + i f(x, y) , \tag{5.62}$$

where $\tilde{\chi}$ is the constant real part of the field (which corresponds to $\frac{\theta_{\mathrm{YM}}}{2\pi}$, see equation (5.15)) and just like in the flat ten-dimensional case

$$f = A - \frac{1}{g_s} + e^{-\hat{\phi}} . \tag{5.63}$$

Since the action is independent of constant shifts of χ, actually it does not depend on the arbitrary constant, A. In a manner that follows closely the flat ten-dimensional case considered in the appendix of [114] and reviewed in the preceding section, the action for a charge K D-instanton can be written as

$$S_K = -\frac{L^{10}}{(\alpha')^4}\int \frac{d\rho d^4x d^5\omega}{\rho^5} g^{\Lambda\Sigma}\nabla_\Lambda(e^{2\hat{\phi}} f\partial_\Sigma f)\,, \tag{5.64}$$

which reduces to an integral over the boundaries of $\mathrm{AdS}_5\times S^5$ and the surface of an infinitesimal sphere centered on the D-instanton at $x = x_0$, $y = y_0$. The result is exactly the same as in the flat case

$$S_K = \frac{2\pi|K|}{g_s}\,. \tag{5.65}$$

By choosing $A = \frac{1}{g_s}$ in (5.63) the expression (5.64) reduces to an integral over the boundary at $\rho = 0$. Remarkably, in this case the boundary integrand is *identical* to the action density of the standard four-dimensional Yang–Mills instanton.

Whereas the $\mathrm{AdS}_5\times S^5$ metric remains unchanged by the presence of the D-instanton in the Einstein frame, it is radically altered in the string frame where the instanton is manifested as a space-time wormhole (as in the flat ten-dimensional case [175]). For finite values of K the dilaton becomes large in the Planck-scale neck and the classical solution is not reliable in that region. However, for very large instanton number, the neck region becomes much larger than the Planck scale so, by analogy with the D-brane examples studied in [38], it should be very interesting to study the implications of the modified $\mathrm{AdS}_5\times S^5$ geometry in the large-K limit of the large-N theory.

The D-instanton contribution to the amplitude with sixteen external dilatinos, Λ^A_α, may now be obtained directly by semi-classical quantization around the classical D-instanton solution in $\mathrm{AdS}_5\times S^5$. The leading instanton contribution can be determined by applying supersymmetry transformations to the scalar field which has an instanton profile given by (5.61). Since the D-instanton background breaks half the supersymmetries the relevant transformations are those in which the supersymmetry parameter corresponds to the Killing spinors for the sixteen broken supersymmetries. These Killing spinors have $\mathrm{U}(1)_B$ charge $\frac{1}{2}$ and are defined by a modified version of (5.17) that includes the non-trivial composite $\mathrm{U}(1)_B$ connection, Q_M [126], that is made from the type IIB scalar field [175]

$$\mathcal{D}_M\zeta \equiv \left(D_M - \frac{i}{2}Q_M\right)\zeta = \frac{1}{2L}\gamma_M\zeta\,. \tag{5.66}$$

Substituting the Euclidean D-instanton solution into the expression for the composite connection gives

$$Q_M = \frac{i}{2} e^{-\hat{\phi}} \partial_M e^{\hat{\phi}} \,, \tag{5.67}$$

with $\hat{\phi}$ defined by (5.61). The solution of (5.66) is

$$\zeta_\pm = e^{-\frac{\hat{\phi}}{4}} \frac{z_M \gamma^{\hat{M}}}{\sqrt{\rho_0}} \zeta_\pm^{(0)} \,, \tag{5.68}$$

where $\zeta_\pm^{(0)}$ is a constant spinor satisfying $\gamma_5 \zeta_\pm^{(0)} = \pm \zeta_\pm^{(0)}$.

The sixteen broken supersymmetry transformations associated with $\zeta_-^{(0)}$ give rise to the dilatino zero-modes,

$$\Lambda_{(0)} = \delta\Lambda = (\gamma^M \hat{P}_M) \zeta_- \,, \tag{5.69}$$

where $\hat{P}_M$ is the expression for $P_M \equiv \frac{i}{2\tau_2} \partial_M \tau^*$ in the D-instanton background [114], *i.e.*

$$\hat{P}_M = e^{-\hat{\phi}} \partial_M e^{\hat{\phi}} \,. \tag{5.70}$$

Using the Killing spinor equation and the D-instanton equation $\mathcal{D}^M \hat{P}_M = 0$ it is easy to check (recalling that P_M has $\mathrm{U}(1)_B$ charge 2) that

$$\gamma^M \mathcal{D}_M \Lambda_{(0)} = -\frac{3}{2L} \Lambda_{(0)} \,, \tag{5.71}$$

so that $\Lambda_{(0)}$ is a solution of the appropriate massive Dirac equation. The relevant amplitudes are those with external states located on the boundary, in which case one may use the fact that for $\rho \sim 0$

$$P_M \sim \frac{1}{g_s} \partial_M e^{\hat{\phi}} \tag{5.72}$$

in (5.69), which leads to

$$\Lambda_{(0)} \sim \frac{4}{g_s} (e^{\hat{\phi}} - g_s) \zeta_- \,. \tag{5.73}$$

This means that near $\rho = 0$ the dilatino profile in the D-instanton background is proportional to $\rho^4 K_4(x_0, \rho_0; x, 0)$.

As a result the leading contribution to the sixteen-dilatino amplitude again reproduces the corresponding sixteen-current correlator in $\mathcal{N} = 4$ supersymmetric Yang–Mills theory. Explicitly, the D-instanton approximation to the amplitude with sixteen

external dilatinos, Λ^A_α, at points on the $\rho = 0$ boundary is (up to an overall constant factor)

$$\langle \prod_{p=1}^{16} \Lambda^{A_p}_{\alpha_p}(x_p,0)\rangle_J = g_s^{-12} e^{-2\pi K(\frac{1}{g_s}+iC^{(0)})} V_{S^5} \int \frac{d^4x_0 d\rho_0}{\rho_0^5} \int d^{16}\zeta^{(0)}_- \cdot \prod_{p=1}^{16} \left[K_4\left(x_0,\rho_0;x_p\right) \frac{1}{\sqrt{\rho_0}} \left(\rho_0 \eta^{A_p}_{\alpha_p} + \left(x_p - x_0\right)_\mu \sigma^\mu_{\alpha_p \dot{\alpha}_p} \overline{\xi}^{\dot{\alpha}_p A_p} \right) J_\Lambda(x_p) \right], \tag{5.74}$$

where $J_\Lambda(x_p)$ is the wave-function of the dilatino evaluated at the boundary point $(x_p, 0)$ and the Grassmann spinor $\zeta^{(0)}_-$ is defined in terms of η and $\bar{\xi}$ by

$$\zeta(\rho_0, x - x_0) = \left(\frac{1-\gamma^5}{2} \right) \frac{z^M \gamma_M \zeta^{(0)}}{\sqrt{\rho}_0} ,$$

where

$$\zeta^{(0)} = \begin{pmatrix} \eta_\alpha \\ \\ \overline{\xi}^{\dot{\alpha}} \end{pmatrix} , \quad z^M = (x^\mu - x^\mu_0, \rho_0) .$$

The power of g_s^{-12} has been inserted in (5.74) from the expression given in section (5.1) although this power should also follow directly by considering the normalization of the bosonic and fermionic zero modes. Up to the overall constant factor, the amplitude (5.74) agrees with (5.58) and therefore with (4.55).

In similar manner the instanton profiles of all the fields in the supergravity multiplet follow by applying the broken supersymmetries to P_M the appropriate number of times, just as in the flat ten-dimensional case [114]. The single D-instanton contributions to any correlation function can then be determined. This is guaranteed to agree with the corresponding term in the expansion of the effective type IIB action as well as with the corresponding $\mathcal{N}$=4 Yang–Mills correlation function.

Conclusions

This dissertation reports the study of various aspects of $\mathcal{N}$=4 supersymmetric Yang–Mills theory.

The perturbative finiteness properties of the theory have been known for a long time. However the analysis presented here allows to point out several problems in the perturbative computation of Green functions of elementary (super) fields. The calculations reviewed in chapter 3 show that the gauge-fixing procedure is quite problematic and presents subtleties both in the description in terms of component fields and using the $\mathcal{N}$=1 superfield formalism. The appearance of UV divergences in off-shell propagators, when the Wess–Zumino gauge in components is exploited, has been discussed and it has been shown that relaxing the choice of the WZ gauge such divergences are cancelled. Different difficulties have been encountered in superspace computations: the supersymmetric generalization of the Fermi–Feynman gauge appears to be privileged since any different choice of gauge leads to IR divergences in off-shell Green functions.

These issues have been studied in a systematic way in chapter 3. The divergences found in one-loop calculations are expected to be gauge artifacts, and it would be interesting to obtain an explicit check of the cancellation of such divergences in gauge invariant Green functions. As briefly discussed in the final chapter in the context of the AdS/SCFT correspondence, superspace techniques can be applied to the calculation of correlation functions of gauge-invariant composite operators as well. $\mathcal{N}$=1 superfields should prove a powerful tool in this computations like in the case of elementary fields.

At the perturbative level the effect of the inclusion of mass terms for the (anti) chiral superfields has also been discussed. The considered mass terms break super-

symmetry down to $\mathcal{N}$=1, but it is explicitly proved, at the one loop-level, that the ultraviolet properties of the model are not affected by this modification. The same result had been obtained long ago by means of dimensional arguments and is reinforced by the analysis presented here.

The possibility of employing this mass deformed $\mathcal{N}$=4 model as a supersymmetry preserving regularization scheme for $\mathcal{N}$=1 theories has been recently proposed. As observed in chapter 3 this approach should be generalizable to $\mathcal{N}$=2 theories.

Non perturbative effects in $\mathcal{N}$=4 supersymmetric Yang–Mills theory are studied in chapter 4 with the aid of instanton calculus. The literature on $\mathcal{N}$=4 super Yang–Mills contains examples of instanton computations of corrections to the effective action in the Abelian Coulomb phase. The calculations presented in chapter 4 have been performed in the superconformal phase, corresponding to the origin of the moduli space, which is related to type IIB superstring theory in $\mathrm{AdS}_5 \times S^5$ by the duality recently proposed by Maldacena. In particular an exact field theoretical computation of the one-instanton contribution (in the semiclassical approximation) to a four-point function of scalar composite operators belonging to the current supermultiplet has been reported.

The non-perturbative corrections to Green functions in $\mathcal{N}$=4 supersymmetric Yang–Mills theory have been subsequently reconsidered in the context of the AdS/CFT correspondence. Previous checks of the conjectured correspondence had been obtained for two- and three-point functions at the perturbative level. The results reviewed here strongly support the possibility of extending the correspondence at the non-perturbative level as well. The agreement found between the contributions to four- and higher-point functions due to two different sources of non-perturbative effects, instantons in $\mathcal{N}$=4 super Yang–Mills and D-instantons in type IIB superstring theory, represents a highly non trivial check of the conjecture. Furthermore these results have been obtained in the case of a SU(2) gauge group and therefore suggest that the correspondence, originally formulated for a SU(N) gauge group in the large N limit, may actually hold at finite N as well. Perturbative checks of this claim have already been proposed, but the non-perturbative results discussed here bring stronger evidence in this direction, since they concern four- and higher-point functions that are not fixed by superconformal symmetry. Despite these interesting results on the field theory side, there is still very little understanding of the correct way of dealing with string corrections to the supergravity approximation for finite N. This is a very interesting subject for future developments. In particular generalizations of the orig-

inal conjecture to theories with less supersymmetry and even to non-supersymmetric theories have been proposed. In this context computations of quantities of phenomenological interest, such as glueball spectra, have been performed. However the possibility of comparing this results with, for instance, the continuum extrapolations from lattice simulations requires the extension of the correspondence to weak 't Hooft coupling, *i.e.* to finite N including string effects.

The four-point correlation function computed in chapter 4 displays logarithmic singularities when any two insertion points are taken to coincide. This peculiar behaviour in the operator product expansion (OPE) has also been observed by other authors in supergravity calculations in AdS_5 and is a very interesting feature of the superconformal field theory at hand.

In understanding this phenomenon a crucial rôle is played by the fact that the chiral primary operators, $\mathcal{O}_q^\Delta$, that enter the Green functions under consideration are protected, *i.e.* their dimension Δ and charge q are related by $\Delta = q$. The first point to observe is the absence of pole-type singularities in the OPE. This is a direct consequence of the condition $\Delta = q$. If one considers the OPE for chiral primary operators with charges q_1 and q_2,

$$\mathcal{O}_{(q_1)}(x)\mathcal{O}_{(q_2)}(y) \qquad \text{for} \quad |x-y| \to 0\,,$$

there are two possibilities. Either the OPE contains only chiral primary operators and one has

$$\mathcal{O}_{(q_1)}(x)\mathcal{O}_{(q_2)}(y) \sim c\,\mathcal{O}^\Delta_{(q_1+q_2)}\left(\frac{x+y}{2}\right) + \text{ terms vanishing as } \quad |x-y| \to 0\,,$$

with $\Delta = q_1 + q_2$ and c a numerical constant, or there is a mixing with a non chiral primary operator, whose dimension is strictly larger than the charge, and then

$$\mathcal{O}_{(q_1)}(x)\mathcal{O}_{(q_2)}(y) \sim \mathcal{V}^\Delta_q\left(\frac{x+y}{2}\right) + \dots\,,$$

where $q = q_1 + q_2$ and $\Delta = q_1 + q_2 + \gamma > q_1 + q_2$. As a consequence in any case there cannot be a pole-type singularity. The second situation described here could be responsible for the logarithmic behaviour. The anomalous dimension γ is determined by the calculation of the three point function $\langle \mathcal{V}\mathcal{O}\mathcal{O}\rangle$, that is obtained by taking the limit of coincident points in a four-point function of chiral primary operators like the one that is computed in chapter 4. Notice that one cannot study the limit in a three-point function, since three-point functions of chiral primaries are not renormalized.

The logarithmically divergent coefficient in the OPE of $\mathcal{O}_{(q_1)}$ and $\mathcal{O}_{(q_2)}$ could originate from a perturbative expansion of $\gamma = \gamma(g)$. More precisely if the operator $\mathcal{V}$ has an anomalous dimension, γ, in perturbation theory the latter is expected to be small and to admit an expansion in the coupling constant g. Expanding $\mathcal{O}_{(q_1)}(x)\mathcal{O}_{(q_2)}(y)$ in powers of γ yields

$$\begin{aligned}\langle \mathcal{O}_{(q_1)}(x)\mathcal{O}_{(q_2)}(y)\rangle \quad \sim \quad & \frac{1}{(x-y)^{2\Delta}}\left[1-\gamma\log(\mu^2(x-y)^2)+\right. \\ & \left.+\frac{1}{2}\gamma^2(\log(\mu^2(x-y)^2))^2+\dots\right],\end{aligned}$$

where an arbitrary mass scale μ has been introduced for dimensional reasons. At each non-trivial order in g, the last equation seems to violate conformal invariance. The exact expression is however conformally invariant by construction. One can reconcile the two results by observing that because of the presence of an anomalous dimension, the operator has to be renormalized at each order in g and as a consequence it does not provide a linear representation of the conformal group.

Appendix A

Notations and spinor algebra

Minkowskian metric

$$\eta_{\mu\nu} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} . \tag{A.1}$$

σ-matrices

$$\sigma^\mu = (-\mathbb{1}_2, \sigma^i) , \qquad \overline{\sigma}^\mu = (-\mathbb{1}_2, -\sigma^i) , \qquad i = 1, 2, 3 , \tag{A.2}$$

where

$$\sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} , \quad \sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} , \quad \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} . \tag{A.3}$$

Weyl spinors are denoted as follows

$$\psi_\alpha , \quad \overline{\psi}^{\dot\alpha} , \quad \text{with} \quad \alpha, \dot\alpha = 1, 2 . \tag{A.4}$$

Spinor indices are raised and lowered by the antisymmetric tensors $\varepsilon^{\alpha\beta}$ ($\varepsilon^{12} = -\varepsilon^{21} = -\varepsilon_{12} = \varepsilon_{21} = 1$) and $\varepsilon^{\dot\alpha\dot\beta}$ ($\varepsilon^{\dot1\dot2} = -\varepsilon^{\dot2\dot1} = -\varepsilon_{\dot1\dot2} = \varepsilon_{\dot2\dot1} = 1$)

$$\begin{aligned} \psi^\alpha &= \varepsilon^{\alpha\beta}\psi_\beta , \quad \psi_\alpha = \varepsilon_{\alpha\beta}\psi^\beta , \\ \overline{\psi}^{\dot\alpha} &= \varepsilon^{\dot\alpha\dot\beta}\overline{\psi}_{\dot\beta} , \quad \overline{\psi}_{\dot\alpha} = \varepsilon_{\dot\alpha\dot\beta}\overline{\psi}^{\dot\beta} . \end{aligned} \tag{A.5}$$

The matrices σ^μ have the following index structure

$$\sigma^\mu = \sigma^\mu{}_{\alpha\dot\alpha} . \tag{A.6}$$

Indices of the σ^μ matrices are raised and lowered by the ε tensor as well

$$\overline{\sigma}^{\mu\,\dot{\alpha}\alpha} = \varepsilon^{\dot{\alpha}\dot{\beta}}\varepsilon^{\alpha\beta}\sigma^\mu{}_{\beta\dot{\beta}}\,. \tag{A.7}$$

Properties of the σ-matrices.

The following properties of the σ-matrices can be easily derived from their definition

$$\begin{aligned}(\sigma^\mu\overline{\sigma}^\nu + \sigma^\nu\overline{\sigma}^\mu)_\alpha{}^\beta &= -2\eta^{\mu\nu}\delta_\alpha{}^\beta \\ (\overline{\sigma}^\mu\sigma^\nu + \overline{\sigma}^\nu\sigma^\mu)^{\dot{\alpha}}{}_{\dot{\beta}} &= -2\eta^{\mu\nu}\delta^{\dot{\alpha}}{}_{\dot{\beta}}\,,\end{aligned} \tag{A.8}$$

as well as the completeness relations

$$\begin{aligned}\mathrm{tr}\,(\sigma^\mu\overline{\sigma}^\nu) &= -2\eta^{\mu\nu} \\ \sigma^\mu_{\alpha\dot{\alpha}}\overline{\sigma}_\mu^{\dot{\beta}\beta} &= -2\delta_\alpha{}^\beta\delta_{\dot{\alpha}}{}^{\dot{\beta}}\,.\end{aligned} \tag{A.9}$$

The generators of Lorentz transformations read

$$\begin{aligned}\sigma^{\mu\nu}{}_\alpha{}^\beta &= \frac{1}{4}\left(\sigma^\mu_{\alpha\dot{\alpha}}\overline{\sigma}^{\nu\,\dot{\alpha}\beta} - \sigma^\nu_{\alpha\dot{\alpha}}\overline{\sigma}^{\mu\,\dot{\alpha}\beta}\right) \\ \overline{\sigma}^{\mu\nu\,\dot{\alpha}}{}_{\dot{\beta}} &= \frac{1}{4}\left(\overline{\sigma}^{\mu\,\dot{\alpha}\alpha}\sigma^\nu_{\alpha\dot{\beta}} - \overline{\sigma}^{\nu\,\dot{\alpha}\alpha}\sigma^\mu_{\alpha\dot{\beta}}\right)\,.\end{aligned} \tag{A.10}$$

Euclidean σ-matrices.

They are defined as [1]

$$\sigma^\mu_{\mathrm{E}} = (\mathbb{1}_2, i\sigma^i)\,, \qquad \overline{\sigma}^\mu_{\mathrm{E}} = (\mathbb{1}_2, -i\sigma^i)\,, \qquad i = 1,2,3\,. \tag{A.11}$$

The Euclidean σ-matrices satisfy properties analogous to their Minkowskian couterparts. In particular

$$\begin{aligned}(\sigma^\mu_{\mathrm{E}}\overline{\sigma}^\nu_{\mathrm{E}} + \sigma^\nu_{\mathrm{E}}\overline{\sigma}^\mu_{\mathrm{E}})_\alpha{}^\beta &= 2\delta^{\mu\nu}\delta_\alpha{}^\beta \\ (\overline{\sigma}^\mu_{\mathrm{E}}\sigma^\nu_{\mathrm{E}} + \overline{\sigma}^\nu_{\mathrm{E}}\sigma^\mu_{\mathrm{E}})^{\dot{\alpha}}{}_{\dot{\beta}} &= 2\delta^{\mu\nu}\delta^{\dot{\alpha}}{}_{\dot{\beta}}\,,\end{aligned} \tag{A.12}$$

$$\begin{aligned}\mathrm{tr}\,(\sigma^\mu_{\mathrm{E}}\overline{\sigma}^\nu_{\mathrm{E}}) &= 2\delta^{\mu\nu} \\ \sigma^\mu_{\mathrm{E}\,\alpha\dot{\alpha}}\overline{\sigma}^{\dot{\beta}\beta}_{\mathrm{E}\,\mu} &= 2\delta_\alpha{}^\beta\delta_{\dot{\alpha}}{}^{\dot{\beta}}\,.\end{aligned} \tag{A.13}$$

[1] The subscript E is suppressed in the text.

The Euclidean Lorentz generators are

$$\begin{aligned}\sigma_{\mathrm{E}}^{\mu\nu}{}_{\alpha}{}^{\beta} &= \frac{1}{4}\left(\sigma^{\mu}_{\mathrm{E}\,\alpha\dot{\alpha}}\overline{\sigma}_{\mathrm{E}}^{\nu\,\dot{\alpha}\beta} - \sigma^{\nu}_{\mathrm{E}\,\alpha\dot{\alpha}}\overline{\sigma}_{\mathrm{E}}^{\mu\,\dot{\alpha}\beta}\right)\\ \overline{\sigma}_{\mathrm{E}}^{\mu\nu\dot{\alpha}}{}_{\dot{\beta}} &= \frac{1}{4}\left(\overline{\sigma}_{\mathrm{E}}^{\mu\,\dot{\alpha}\alpha}\sigma^{\nu}_{\mathrm{E}\,\alpha\dot{\beta}} - \overline{\sigma}_{\mathrm{E}}^{\nu\,\dot{\alpha}\alpha}\sigma^{\mu}_{\mathrm{E}\,\alpha\dot{\beta}}\right).\end{aligned} \tag{A.14}$$

The following realization of the Dirac γ-matrices is used

$$\gamma^{\mu} = \begin{pmatrix} 0 & \sigma^{\mu} \\ \overline{\sigma}^{\mu} & 0 \end{pmatrix}, \qquad \gamma^{5} = i\gamma^{0}\gamma^{1}\gamma^{2}\gamma^{3} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \tag{A.15}$$

Spinor algebra.

Contractions between spinor indices are taken according to the conventions

$$\begin{aligned}\psi\chi &= \psi^{\alpha}\chi_{\alpha} = -\psi_{\alpha}\chi^{\alpha} = \chi\psi\\ \overline{\psi}\overline{\chi} &= \overline{\psi}_{\dot{\alpha}}\overline{\chi}^{\dot{\alpha}} = -\overline{\psi}^{\dot{\alpha}}\overline{\chi}_{\dot{\alpha}} = \overline{\chi}\overline{\psi}.\end{aligned} \tag{A.16}$$

Fierz rearrangements

$$\begin{aligned}\psi^{\alpha}\psi^{\beta} &= -\frac{1}{2}\varepsilon^{\alpha\beta}\psi\psi\,, \qquad & \psi_{\alpha}\psi_{\beta} &= \frac{1}{2}\varepsilon_{\alpha\beta}\psi\psi\\ \overline{\psi}^{\dot{\alpha}}\overline{\psi}^{\dot{\beta}} &= \frac{1}{2}\varepsilon^{\dot{\alpha}\dot{\beta}}\overline{\psi\psi}\,, \qquad & \overline{\psi}_{\dot{\alpha}}\overline{\psi}_{\dot{\beta}} &= -\frac{1}{2}\varepsilon_{\dot{\alpha}\dot{\beta}}\overline{\psi\psi}\,.\end{aligned} \tag{A.17}$$

Miscellaneous relations

$$\begin{aligned}\theta\sigma^{\mu}\overline{\theta}\theta\sigma^{\nu}\overline{\theta} &= -\frac{1}{2}\theta\theta\overline{\theta}\overline{\theta}\eta^{\mu\nu}\\ (\theta\psi)(\theta\chi) = -\frac{1}{2}(\theta\theta)(\psi\chi) \quad &, \qquad (\overline{\theta\psi})(\overline{\theta}\overline{\chi}) = -\frac{1}{2}(\overline{\theta\theta})(\overline{\psi}\overline{\chi})\\ \chi\sigma^{\mu}\overline{\psi} = -\psi\sigma^{\mu}\overline{\chi} \quad &, \qquad (\chi\sigma^{\mu}\overline{\psi})^{\dagger} = \psi\sigma^{\mu}\overline{\chi}\\ \chi\sigma^{\mu}\overline{\sigma}^{\nu}\overline{\psi} = \psi\sigma^{\nu}\overline{\sigma}^{\mu}\overline{\chi} \quad &, \qquad (\chi\sigma^{\mu}\overline{\sigma}^{\nu}\overline{\psi})^{\dagger} = \overline{\psi}\overline{\sigma}^{\nu}\sigma^{\mu}\overline{\chi}\\ (\psi\chi)\overline{\lambda}_{\dot{\alpha}} = -\frac{1}{2}(\chi\sigma^{\mu}\overline{\lambda})(\psi\sigma_{\mu})_{\dot{\alpha}}\,.\end{aligned} \tag{A.18}$$

Superspace covariant derivatives.

Covariant derivatives in Minkowskian superspace are defined as

$$D_{\alpha} = \frac{\partial}{\partial\theta^{\alpha}} + i\sigma^{\mu}_{\alpha\dot{\alpha}}\overline{\theta}^{\dot{\alpha}}\partial_{\mu}\,, \qquad \overline{D}_{\dot{\alpha}} = -\frac{\partial}{\partial\overline{\theta}^{\dot{\alpha}}} - i\theta^{\alpha}\sigma^{\mu}_{\alpha\dot{\alpha}}\partial_{\mu} \tag{A.19}$$

and satisfy

$$\begin{aligned} \{D_\alpha, \overline{D}_{\dot\alpha}\} &= -2i\sigma^\mu_{\alpha\dot\alpha}\partial_\mu \\ \{D_\alpha, D_\beta\} &= \{\overline{D}_{\dot\alpha}, \overline{D}_{\dot\beta}\} = 0\,. \end{aligned} \tag{A.20}$$

Moreover

$$\varepsilon^{\alpha\beta}\frac{\partial}{\partial\theta^\alpha} = -\frac{\partial}{\partial\theta_\alpha}\,, \qquad \varepsilon^{\alpha\beta}\frac{\partial}{\partial\theta^\alpha}\frac{\partial}{\partial\theta^\beta}\theta\theta = \varepsilon^{\dot\alpha\dot\beta}\frac{\partial}{\partial\overline{\theta}^{\dot\alpha}}\frac{\partial}{\partial\overline{\theta}^{\dot\beta}}\overline{\theta\theta} = 4\,. \tag{A.21}$$

Other properties

$$\begin{aligned} &[\overline{D}_{\dot\alpha}, \{\overline{D}_{\dot\beta}, D_\gamma\}] = 0\,, \qquad [D_\alpha, \overline{D}_{\dot\beta}\overline{D}^{\dot\beta}] = -4i\sigma^\mu_{\alpha\dot\alpha}\partial_\mu\overline{D}^{\dot\alpha} \\ &[D^2, \overline{D}^2] = -8iD^\alpha\sigma^\mu_{\alpha\dot\alpha}\overline{D}^{\dot\alpha}\partial_\mu - 16\Box = 8i\overline{D}^{\dot\alpha}\sigma^\mu_{\alpha\dot\alpha}D^\alpha\partial_\mu + 16\Box\,. \end{aligned} \tag{A.22}$$

The corresponding Euclidean superspace derivatives read

$$D^{\rm E}_\alpha = \frac{\partial}{\partial\theta^\alpha} + \sigma^\mu_{\alpha\dot\alpha}\overline{\theta}^{\dot\alpha}\partial_\mu\,, \qquad \overline{D}^{\rm E}_{\dot\alpha} = -\frac{\partial}{\partial\overline{\theta}^{\dot\alpha}} - \theta^\alpha\sigma^\mu_{\alpha\dot\alpha}\partial_\mu \tag{A.23}$$

and possess analogous properties

$$\begin{aligned} \{D^{\rm E}_\alpha, \overline{D}^{\rm E}_{\dot\alpha}\} &= -2\sigma^\mu_{\alpha\dot\alpha}\partial_\mu \\ \{D^{\rm E}_\alpha, D^{\rm E}_\beta\} &= \{\overline{D}^{\rm E}_{\dot\alpha}, \overline{D}^{\rm E}_{\dot\beta}\} = 0\,. \end{aligned} \tag{A.24}$$

$$\begin{aligned} &[\overline{D}^{\rm E}_{\dot\alpha}, \{\overline{D}^{\rm E}_{\dot\beta}, D^{\rm E}_\gamma\}] = 0 \\ &[D^{\rm E}_\alpha, \overline{D}^{\rm E}_{\dot\beta}\overline{D}^{{\rm E}\,\dot\beta}] = -4\sigma^\mu_{{\rm E}\,\alpha\dot\alpha}\partial_\mu\overline{D}^{{\rm E}\,\dot\alpha}\,, \qquad [D^\alpha_{\rm E}D^{\rm E}_\alpha, \overline{D}^{\rm E}_{\dot\alpha}] = -4\sigma^\mu_{{\rm E}\,\alpha\dot\alpha}\partial_\mu D^\alpha_{\rm E} \\ &[D^2_{\rm E}, \overline{D}^2_{\rm E}] = -8D^\alpha_{\rm E}\sigma^\mu_{{\rm E}\,\alpha\dot\alpha}\overline{D}^{\dot\alpha}_{\rm E}\partial_\mu - 16\Box = 8\overline{D}^{\dot\alpha}_{\rm E}\sigma^\mu_{{\rm E}\,\alpha\dot\alpha}D^\alpha_{\rm E}\partial_\mu + 16\Box\,. \end{aligned} \tag{A.25}$$

The following relations are useful in superspace calculations

$$\begin{aligned} &D^2_{\rm E}\overline{D}^2_{\rm E}D^2_{\rm E} = 16\Box D^2_{\rm E}\,, \qquad \overline{D}^2_{\rm E}D^2_{\rm E}\overline{D}^2_{\rm E} = 16\Box\overline{D}^2_{\rm E} \\ &\overline{D}^2_{\rm E}D^{\rm E}_\alpha D^{\rm E}_\beta\overline{D}^2_{\rm E} = -16\sigma^\mu_{{\rm E}\,\alpha\dot\alpha}\sigma^\nu_{{\rm E}\,\beta\dot\beta}\partial_\mu\partial_\nu\overline{D}^{\dot\alpha}_{\rm E}\overline{D}^{\dot\beta}_{\rm E} \\ &D^2_{\rm E}\overline{D}^{\rm E}_{\dot\alpha}\overline{D}^{\rm E}_{\dot\beta}D^2_{\rm E} = -16\sigma^\mu_{{\rm E}\,\alpha\dot\alpha}\sigma^\nu_{{\rm E}\,\beta\dot\beta}\partial_\mu\partial_\nu D^\alpha_{\rm E}D^\beta_{\rm E}\,. \end{aligned} \tag{A.26}$$

Berezin integration.

Integration over a Grassmannian variable η is defined as

$$\int d\eta = 0\,, \qquad \int d\eta\,\eta = 1\,. \tag{A.27}$$

The power expansion of a function $f(\eta)$ is simply

$$f(\eta) = a + b\eta\,, \tag{A.28}$$

so that

$$\int d\eta\, f(\eta) = b\,, \qquad \int d\eta\, \eta f(\eta) = a\,. \tag{A.29}$$

Integration by parts in Berezin integrals is allowed since

$$\int d\eta\, \frac{\partial}{\partial\eta} f(\eta) = 0\,. \tag{A.30}$$

The δ-function is defined by

$$\int d\eta\, f(\eta)\delta(\eta) = f(0)\,, \tag{A.31}$$

which implies $\delta(\eta) = \eta$. The integration measure over the Grassmannian variables of $\mathcal{N}$=1 superspace is defined as follows

$$\begin{aligned} &d^4\theta = d^2\theta d^2\overline{\theta} \\ &d^2\theta = \frac{1}{4}\varepsilon_{\alpha\beta} d\theta^\alpha d\theta^\beta\,, \qquad d^2\overline{\theta} = \frac{1}{4}\varepsilon^{\dot\alpha\dot\beta} d\overline{\theta}_{\dot\alpha} d\overline{\theta}_{\dot\beta}\,, \end{aligned} \tag{A.32}$$

so that

$$\int d^2\theta\, \theta\theta = \int d^2\overline{\theta}\, \overline{\theta\theta} = 1\,. \tag{A.33}$$

Other useful formulas can be found in the appendices of [4, 3]

Appendix B

Electric-magnetic duality

B.1 Electric-magnetic duality and the Dirac monopole

The free Maxwell equations

$$\begin{array}{ll} \vec{\nabla}\cdot\vec{E}=0 & \vec{\nabla}\cdot\vec{B}=0 \\ \vec{\nabla}\times\vec{B}-\dfrac{\partial E}{\partial t}=0 & \vec{\nabla}\times\vec{E}+\dfrac{\partial B}{\partial t}=0\,, \end{array} \tag{B.1}$$

are invariant under a duality transformation D, exchanging electric and magnetic fields according to

$$D: \qquad \vec{E}\longrightarrow\vec{B}\,, \qquad \vec{B}\longrightarrow-\vec{E}\,.$$

More generally the equations are invariant under a continuous transformation

$$\begin{pmatrix} \vec{E} \\ \vec{B} \end{pmatrix} \longrightarrow \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix} \begin{pmatrix} \vec{E} \\ \vec{B} \end{pmatrix}\,. \tag{B.2}$$

A covariant formulation can be given using the language of differential forms. The free equations can be written

$$d\,F=0\,, \qquad *^{-1}d*F=0\,,$$

where $(*F)_{\mu\nu}=\frac{1}{2}\varepsilon_{\mu\nu\rho\sigma}F^{\rho\sigma}$ and d denotes the external derivative. The duality transformation simply becomes $F\rightarrow *F$. In the presence of electric sources the equations become

$$dF=0\,, \qquad *^{-1}d*F=j\,,$$

where j is a one-form of components $j_e^\mu = (\rho_e, \vec{j}_e)$, so that the symmetry is broken. To recover the duality symmetry magnetic source terms must be included symmetrizing the equations as follows

$$dF = j_m \,, \qquad *^{-1} d * F = j_e \,.$$

The system is then symmetric under the simultaneous exchange of F with $*F$ and of j_e with j_m.

Even if the inclusion of magnetic point-like particles in classical electromagnetism seems to be ruled out by the experimental results, it allows to derive a consequence that is of fundamental importance in the more interesting case of non Abelian theories, the *Dirac quantization condition*. A consistent quantization [1] of a theory with both electric and magnetic charges q_i and g_j requires that they satisfy the condition

$$q_i \cdot g_j = 2\pi\hbar n_{ij} \,, \tag{B.3}$$

where n_{ij} are arbitrary integers.

The quantum mechanical description of a charged particle in an electromagnetic field requires the introduction of the vector potential A_μ, related to the field strength by $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, through the substitution $\vec{\nabla} \to (\vec{\nabla} - \frac{iq}{\hbar}\vec{A})$. This leads to the minimal coupling in the Schroedinger equation

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m}(\vec{\nabla} - \frac{iq}{\hbar}\vec{A})^2\psi + V\psi \,.$$

Gauge transformations for this system read

$$\psi \longrightarrow e^{-i\frac{q}{\hbar}\chi(\theta)}\psi \,, \qquad \vec{A} \longrightarrow \vec{A} - \vec{\nabla}\chi(\theta) \,. \tag{B.4}$$

In the presence of a point like magnetic charge it is not possible to associate a globally defined vector potential to the magnetic field through $\vec{B} = \vec{\nabla} \times \vec{A}$. However it is possible to construct a well defined vector potential in the northern hemisfere ($\vec{A}_{_N}$) and in the southern hemisfere ($\vec{A}_{_S}$). This is physically consistent if the fields $\vec{A}_{_N}$ and $\vec{A}_{_S}$ differ by a gauge transformation on the overlapping region (the plane of the equator)

$$\vec{A}_{_N}\Big|_{\rm eq} = \vec{A}_{_S}\Big|_{\rm eq} + \vec{\nabla}\chi$$

[1] Strikingly the Dirac quantization condition can be already derived in the context of classical electromagnetism.

and analogously for the wave function

$$\psi_{_N}|_{\rm eq} = e^{-i\frac{q}{\hbar}\chi}\,\psi_{_S}|_{\rm eq}\ .$$

By means of Stokes' theorem one obtains

$$g = \int_{S^3} d\vec{\sigma}\cdot\vec{B} = \int_{\rm eq} d\vec{\ell}\cdot\vec{A}_{_N} - \int_{\rm eq} d\vec{\ell}\cdot\vec{A}_{_S} = \chi(2\pi) - \chi(0)\,.$$

But the wave function ψ must be single valued along the equator requiring from (B.4) $\chi(2\pi) - \chi(0) = \frac{2\pi\hbar}{q}n$ which leads to the Dirac quantization condition, $qg = 2\pi\hbar n$, with $n \in \mathbb{Z}$. The relation (B.3) is not invariant under the continuous duality transformation (B.2), but it can be generalized to the so called Dirac-Schwinger-Zwanziger condition [57] that is invariant. The latter applies to a system containing two dyonic states (*i.e.* states possessing both electric and magnetic charge) of charges (q_i, g_i) and (q_j, g_j) and reads

$$q_i g_j - q_j g_i = 2\pi\hbar n_{ij}\,.$$

B.2 Non Abelian case: the 't Hooft–Polyakov monopole

As discussed in the preceding section magnetic monopoles must be included at hand in a purely electromagnetic model and there exist no experimental evidence justifying this construction. In non Abelian gauge theories on the contrary monopoles and dyons emerge naturally in association with particular field configurations. Indeed non Abelian gauge theories admit solitonic solutions of the classical equations of motion that can be interpreted as describing monopole (and in general dyon) states in the quantum theory. The case of the *'t Hooft–Polyakov monopole* [180, 181] in the Georgi–Glashow model with gauge group SU(2) will be considered here. The results can be generalized to a larger gauge group G by, for example, embedding a SU(2) factor in G.

The Georgi–Glashow model is defined by the Lagrangian

$$\mathcal{L} = -\frac{1}{4}F^a_{\mu\nu}F^{a\mu\nu} + \frac{1}{2}D^\mu\Phi^a D_\mu\Phi_a - V(\Phi)\,, \tag{B.5}$$

where $F^a_{\mu\nu} = \partial_\mu A^a_\nu - \partial_\nu A^a_\mu + e\varepsilon^{abc}A_{b\mu}A_{c\nu}$ is the non Abelian field strength and the scalar fields Φ^a are in the adjoint representation of the gauge group SU(2) with covariant

derivative [2] $D_\mu \Phi^a = \partial_\mu \Phi^a + e\varepsilon^{abc} A_{b\mu} \Phi_c$. The potential $V(\Phi)$ is

$$V(\Phi) = \frac{\lambda}{4} \left(\Phi^a \Phi_a - v^2 \right)^2 ,$$

so that the classical equations of motion are

$$\begin{aligned} D_\mu F^{a\mu\nu} &= \varepsilon^{abc} \Phi_b D^\nu \Phi_c \\ (D^\mu D_\mu \Phi)^a &= -\lambda \Phi^a (\Phi^b \Phi_b - v^2) . \end{aligned} \tag{B.6}$$

Furthermore $F_{\mu\nu}$ satisfies the Bianchi identity $D_\mu (*F)^{\mu\nu} = 0$.

The total energy for a given field configuration is

$$E = \int d^3x \, \Theta_{00} = \int d^3x \, \left\{ \left[\frac{1}{2} (B_i^a)^2 + (E_i^a)^2 + (\Pi^a)^2 + [(D_i \Phi)^a]^2 \right] + V(\Phi) \right\} , \tag{B.7}$$

where Π^a is the momentum conjugate to Φ^a and E_i^a and B_i^a are the 'non Abelian' electric and magnetic fields

$$F_{0i}^a = E_i^a , \qquad F_{ij}^a = -\varepsilon_{ijk} B^{ak} .$$

The energy density Θ_{00} is semi-positive definite, $\Theta_{00} \geq 0$, with equality if and only if $F_{\mu\nu}^a = D_\mu \Phi^a = V(\Phi) = 0$, implying that the vacuum of the theory corresponds to configurations with vanishing A_μ^a and constant Φ^a satisfying $\Phi^a \Phi_a = v^2$. The set of configurations corresponding to the Higgs vacuum (moduli space) will be denoted by $\mathcal{M}_H$

$$\mathcal{M}_H = \{ \Phi \, : D_\mu \Phi^a = 0 \, , \; V(\Phi) = 0 \}$$

In the case at hand $\mathcal{M}_H$ is the sphere S^2 parametrized by the 'coordinates' Φ^a constrained by $\Phi^a \Phi_a = v^2$.

For $v \neq 0$ the gauge symmetry is broken to U(1) and the Higgs mechanism produces a perturbative spectrum which consists of a massless 'photon', massive bosons $W^\pm$ with mass $m_{_W} = ev$ and a massive Higgs scalar with mass $m_{_H} = \sqrt{2\lambda} v$.

In addition to the constant vacuum solution the classical equations of motion admit solitonic solutions corresponding to static field configurations with finite (and non-vanishing) energy. In order to have finite energy the fields Φ^a must approach (sufficiently rapidly) values in $\mathcal{M}_H$ for large values of the radial coordinate r, so that they define a map $\Phi \, : S^2_\infty \to \mathcal{M}_H$. For the Georgi–Glashow model one has $\mathcal{M}_H = S^2$

[2] From now on natural units will be used so that $\hbar = c = 1$.

and the corresponding solitonic configurations are characterized by an integer winding number $n \in \Pi_2(S^2) \equiv \mathbb{Z}$. It can be shown [182] that the most general solution for the gauge field is

$$A^a_\mu = \frac{1}{v^2 e}\varepsilon^{abc}\Phi_b\partial_\mu\Phi_c + \frac{1}{v}\Phi^a K_\mu\,,$$

with an arbitrary K_μ. The resulting field strength is

$$F^a_{\mu\nu} = \frac{1}{v}\Phi^a F_{\mu\nu}\,,$$

where $F_{\mu\nu}$ is an 'Abelian' field strength

$$F_{\mu\nu} = \frac{1}{v^3 e}\varepsilon_{abc}\Phi^a\partial_\mu\Phi^b\partial_\nu\Phi^c + \partial_\mu K_\nu - \partial_\nu K_\mu\,,$$

satisfying the Maxwell equations. The corresponding magnetic field $B_i = \frac{1}{2}\varepsilon_{ijk}F^{jk}$ allows to define a magnetic charge obeying the Dirac quantization condition

$$g = \int_{S^2_\infty} d\vec{\sigma}\cdot\vec{B} = \frac{1}{2ev^3}\int_{S^2_\infty} d^2\sigma^i\,\varepsilon_{ijk}\varepsilon^{abc}\Phi_a\partial^j\Phi_b\partial^k\Phi_c = \frac{4\pi n}{e}\,, \tag{B.8}$$

where n is the winding number.

The configuration (A^a_μ, Φ^a) is interpreted as describing a magnetic monopole. In fact the energy density decreases to zero outside a finite region as is expected for a 'localized' object. Furthermore computing the non Abelian magnetic field

$$B^a_i = \frac{1}{2}\varepsilon_{ijk}F^{ajk} = \frac{1}{2v^4 e}\Phi^a\varepsilon_{ijk}\varepsilon^{bcd}\Phi_b\partial^j\Phi_c\partial^k\Phi_d = \frac{1}{v}\Phi^a B_i$$

allows to associate a magnetic charge to the configuration using the fact that for large r Φ^a must approach values in $\mathcal{M}_H$, *i.e.* $\Phi^a\Phi_a = v^2$ on S^2_∞, so that

$$g = \int_{S^2_\infty} d\vec{\sigma}\cdot\frac{1}{v}\Phi^a\vec{B}_a = \frac{1}{v^2}\int_{S^2_\infty} d\vec{\sigma}\cdot\Phi_a\Phi^a\vec{B} = \int_{S^2_\infty} d\vec{\sigma}\cdot\vec{B} = \frac{4\pi n}{e}\,. \tag{B.9}$$

Equation (B.9) shows that the "magnetic particle" satisfies the Dirac quantization condition, $eg = 4\pi n$. Notice that the non Abelian field strength is aligned along the Higgs field outside the core of the monopole. The magnetic charge g can be written in the form

$$g = \frac{1}{v}\int_{S^2_\infty} d\vec{\sigma}\cdot\vec{B}^a\Phi_a = \frac{1}{v}\int d^3x\,\vec{B}^a(\vec{D}\Phi)^a\,,$$

which follows from Stokes' theorem and the Bianchi identity, which implies $D_iB^i = 0$.

For a static configuration with vanishing electric field one then finds that the 'mass' given by (B.7) satisfies the *Bogomol'nyi bound*

$$m_{_M} \geq v|g|\,. \tag{B.10}$$

In (B.10) the equality holds if and only if $V(\Phi) = 0$ identically (*i.e.* $\lambda = 0$ in the Georgi–Glashow model) and the first order Bogomol'nyi equation

$$B_i^a = D_i\Phi^a \tag{B.11}$$

is satisfied.

An explicit solution for the monopole field configuration can be found in the special case of spherical symmetry and if the Bogomol'nyi bound is saturated. A field configuration corresponding to a non trivial vacuum cannot be invariant either under the group of spatial rotations $\mathrm{SO(3)_R}$ or under the $\mathrm{SU(2)_G}$ group of global gauge transformations [3]. However a solution invariant under the diagonal SO(3) subgroup of $\mathrm{SO(3)_R}\times\mathrm{SU(2)_G}$ can be obtained through the ansatz

$$\Phi^a = \frac{x^a}{er^2}H(\xi)\,, \quad A_0^a = 0\,, \quad A_i^a = -\varepsilon^a{}_{ij}\frac{x^j}{er^2}[1-K(\xi)]\,,$$

where $r = |x|$, $\xi = ver$ and the functions H and K satisfy $K(\xi) \to 1$, $H(\xi) \to 0$ for $r \to 0$ and $K(\xi) \to 0$, $H(\xi)/\xi \to 1$ for $r \to \infty$. In the limit in which the Bogomol'nyi bound is saturated an explicit solution was derived by Prasad and Sommerfield [183]. The exact form of the solution will not be discussed here since it will not be relevant for future considerations.

In general a non trivial vacuum in the limit $V(\Phi) = 0$ (BPS limit [4]) is rather unnatural and moreover the condition of vanishing potential is expected to be broken by quantum effects. However in supersymmetric theories this situation is perfectly consistent because of the presence of flat directions that are protected by supersymmetry. The construction leading to the monopoles considered so far can be generalized to the case of non vanishing electric field giving rise to configurations with both electric and magnetic charge (dyons) [184] [5]; for such states the Bogomol'nyi bound becomes

$$m_{_D} \geq v\sqrt{e^2+g^2}\,.$$

[3] Gauge transformations that do not reduce to the identity at spatial infinity are referred to as "global" or "large" gauge transformations.

[4] Field configurations saturating the Bogomol'nyi bound are called "BPS saturated".

[5] Static dyon solutions corresponding to finite energy configurations cannot be constructed in the temporal gauge $A_0 = 0$

The Lagrangian (B.5) can be modified by the introduction of a θ-term

$$\mathcal{L}_\theta = -\frac{\theta e^2}{32\pi^2} F^a_{\mu\nu} * F^{\mu\nu}_a \tag{B.12}$$

without spoiling renormalizability. The addition of $\mathcal{L}_\theta$ amounts to consider a complexified coupling constant

$$\tau = \frac{\theta}{2\pi} + \frac{4i\pi}{e^2} \tag{B.13}$$

and write the Lagrangian as

$$\mathcal{L} = -\frac{1}{32\pi} \mathrm{Im}\,\tau \left(F^{\mu\nu} + i * F^{\mu\nu}\right)\left(F_{\mu\nu} + i * F_{\mu\nu}\right) - \frac{1}{2} D^\mu \Phi D_\mu \Phi \,. \tag{B.14}$$

In the presence of a θ term the dyons acquire an additional electric charge [185] (*Witten effect*). More precisely the electric and magnetic charge operators are

$$\begin{aligned} Q_m &= \frac{1}{v}\int d^3x\, D_i\Phi^a B^i_a \\ Q_e &= \frac{1}{v}\int d^3x\, D_i\Phi^a E^i_a \end{aligned} \tag{B.15}$$

and one finds for the electric charge of a dyon

$$q = n_e e - \frac{e\theta n_m}{2\pi}\,,$$

with $n_e \in \mathbb{Z}$ and $n_m = \dfrac{eg}{4\pi} \in \mathbb{Z}$.

B.3 Monopoles and fermions

In realistic models the gauge fields are coupled to fermions and one is then led to consider monopole configurations in the presence of fermions. The coupling to spinors in a monopole background has important consequences that will be briefly discussed here in the simplest case of gauge group SU(2).

Dirac fermions can be coupled to the Georgi–Glashow model by adding to (B.5) the term

$$\mathcal{L}_{\mathrm{F}} = \sum_{k=1}^{N_f} i\overline{\psi}^k \gamma^\mu D_\mu \psi^k - i f \overline{\psi}^k \Phi \psi^k \,. \tag{B.16}$$

The Dirac equation reads

$$(i\gamma^\mu D_\mu - \Phi)\psi^k = 0$$

and a suitable choice for the γ-matrices is

$$\gamma^0 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \qquad \gamma^i = \begin{pmatrix} -i\sigma^i & 0 \\ 0 & i\sigma^i \end{pmatrix}.$$

In the background of a monopole one can look for a solution for ψ^k in the form

$$\psi^k(x,t) = e^{iEt}\psi^k(x)\,,$$

with $\psi^k(x) = \begin{pmatrix} \chi_+^k(x) \\ \chi_-^k(x) \end{pmatrix}$. The Weyl spinors $\chi_\pm^k$ then satisfy the equations

$$\begin{aligned} \not{D}\chi_+^k &= (i\sigma^i D_i + \Phi)\chi_-^k = E\chi_+^k \\ \not{D}\chi_-^k &= (i\sigma^i D_i - \Phi)\chi_+^k = E\chi_-^k\,, \end{aligned}$$

which in general have both positive energy ($E>0$) and zero energy (E=0) solutions. The monopole ground state is then constructed as a combination of the zero-energy eigenfunctions that are interpreted as 'fermionic collective coordinates' describing a Grassmannian deformation of the original monopole. The generic solution of the Dirac equation can be written in the form

$$\psi^k = \sum_{i=1}^{n} a_0^i \psi_{0\,i}^k + \sum_{j=1}^{\infty} a^j \psi_j^k\,, \tag{B.17}$$

where $\psi_{0\,i}^k$ are the zero-energy eigenfunctions. The coefficients of the zero-energy eigenfunctions in the expansion (B.17) become creation and annihilation operators and zero-energy solitonic states are obtained by acting with such operators on the "bosonic" monopole ground state. The number of fermionic zero-modes is determined by an index theorem [71].

In the case of fermions in the fundamental representation it can be proved that there is one zero-mode for each flavor for a charge one monopole. Rewriting the N_f Dirac fermions in terms of $2N_f$ Weyl fermions one finds that the creation and annihilation operators generate a Clifford algebra of dimension $2^{2N_f/2} = 2^{N_f}$, that splits into two irreducible representations of dimension 2^{N_f-1}. As a result the monopole ground state is a spinor of SO($2N_f$). The angular momentum generator for a symmetric monopole is $\vec{K} = \vec{L} + \vec{S} + \vec{T}$, where $\vec{T}$ are the SU(2)$_{\rm G}$ generators of global gauge transformations. Fermions in the fundamental can have 'spin' zero since they transform under SU(2)$_{\rm G}\times$SO(3)$_{\rm R}$ in the representation $\mathbf{2}\times\mathbf{2} = \mathbf{1}+\mathbf{3}$ which contains

the singlet. Hence the fermionic zero modes carry zero total angular momentum and the resulting monopole configurations have spin zero.

If the fermions are in the adjoint representation there are two zero-modes for each flavor in the one-monopole background [71]. Furthermore these zero modes necessarily have $\vec{K} \neq 0$ since the adjoint fermions transform in the $\mathbf{3} \times \mathbf{2} = \mathbf{2} + \mathbf{4}$ of $\mathrm{SU}(2)_{\mathrm{G}} \times \mathrm{SO}(3)_{\mathrm{R}}$. Since the zero modes are two-fold degenerate they must carry spin $\pm\frac{1}{2}$. Consequently in the simplest case of N_f=1 the corresponding creation and annihilation operators (that will be denoted by $a_{\pm}^{\dagger}$ and $a_{\pm}$) can be used to generate the following multiplet of zero-energy states acting on the charge-one Clifford vacuum $|\Omega\rangle$

State	Spin
$\|\Omega\rangle$	0
$a_{+}^{\dagger}\|\Omega\rangle$	$\frac{1}{2}$
$a_{-}^{\dagger}\|\Omega\rangle$	$-\frac{1}{2}$
$a_{+}^{\dagger}a_{-}^{\dagger}\|\Omega\rangle$	0

In conclusion the coupling to fermions in the adjoint representation gives a non-vanishing spin to the monopole.

Bibliography

[1] Yu.A. Gol'fand and E.P. Likhtman, "Extension of the algebra of Poincaré group generators and violation of P invariance", *JETP Lett.* **13** (1971) 323.

[2] J. Wess and B. Zumino, "Supergauge transformations in four dimensions", *Nucl. Phys.* **B70** (1974) 39.

[3] P. West, *Introduction to supersymmetry and supergravity*, 2nd Edition, World Scientific (1990).

[4] J. Wess and J. Bagger, *Supersymmetry and supergravity*, 2nd Edition, Princeton University Press (1992).

[5] M. Sohnius, "Introducing supersymmetry", *Phys. Rept.* **128** (1985) 39.

[6] J. Lykken, "Introduction to supersymmetry", `hep-th/9612114`.

[7] S.J. Gates, M.T. Grisaru, M. Roček and W. Siegel, *Superspace or one thousand and one lessons in supersymmetry*, Benjamin/Cummings London (1983).

[8] H.P. Nilles, "Supersymmetry, supergravity and particle physics", *Phys. Rept.* **110** (1984) 1.

[9] M.B. Green, J.H. Schwarz and E. Witten, *Superstring theory*, Voll. 1-2, Cambridge University Press (1987).

[10] S. Coleman and J. Mandula, "All possible symmetries of the S matrix", *Phys. Rev.* **159** (1967) 1251.

[11] R. Haag, J.T. Łopuszański and M. Sohnius, "All possible generators of supersymmetries of the S–matrix", *Nucl. Phys.* **B88** (1975) 257.

[12] M. Shifman,"Non-perturbative gauge dynamics in supersymmetric theories. A primer", in *Confinement, duality and nonperturbative aspects of QCD*, ed. by P. van Baal, Plenum Press, NATO ASI Series.

[13] S. Ferrara and M. Porrati, "AdS superalgebras with brane charges", *Phys. Lett.* **B458** (1999) 43, `hep-th/9903241`.

[14] A. Salam and J. Strathdee, "Supergauge transformations", *Nucl. Phys.* **B76** (1974) 477.

[15] S. Ferrara, J. Wess and B. Zumino, "Supergauge multiplet and superfields", *Phys. Lett.* **51B** (1974) 239.

[16] J. Wess and B. Zumino, "Supergauge invariant extension of quantum electrodynamics", *Nucl. Phys.* **B78** (1974) 1.

[17] A. Salam and J. Strathdee, "On superfields and Fermi–Bose symmetry", *Phys. Rev.* **D11** (1975) 1521.

[18] J. Wess and B. Zumino, "A lagrangian model invariant under supergauge transformations", *Phys. Lett.* **49B** (1974) 52.

[19] S. Ferrara and B. Zumino, "Supergauge invariant Yang–Mills theories", *Nucl. Phys.* **B79** (1974) 413.

[20] A. Salam and J. Strathdee, "Supersymmetry and non Abelian gauges", *Phys. Lett.* **51B** (1974) 353.

[21] G.G. Hartwell and P.S. Howe, "A superspace survey", *Class. Quant. Grav.* **12** (1995) 1823.

[22] A. Galperin, E. Ivanov, S. Kalitzin, V. Ogievetsky and E. Sokatchev, "Unconstrained N=2 matter, Yang–Mills and supergravity theories in harmonic superspace", *Class. Quant. Grav.* **1** (1984) 469.

[23] M.T. Grisaru, M. Roček and W. Siegel, "Improved methods for supergraphs", *Nucl. Phys.* **B159** (1979) 429.

[24] S. Ferrara and O. Piguet, "Perturbation theory and renormalization of supersymmetric Yang–Mills theories", *Nucl. Phys.* **B93** (1975) 261; D. Capper and

G. Leibbrandt, "On the degree of divergence of Feynman diagrams in superfield theories", *Nucl. Phys.* **B85** (1975) 492; K. Fujiikawa and W. Lang, "Perturbation calculations for the scalar multiplet in a superfield formulation", *Nucl. Phys.* **B88** (1975) 61.

[25] J. Wess and B. Zumino, "A Lagrangian model invariant under supergauge transformations", *Phys. Lett.* **49B** (1974) 52; J. Iliopoulos and B. Zumino, "Broken supergauge symmetry and renormalization", *Nucl. Phys.* **B76** (1974) 310; S. Ferrara, J. Iliopoulos and B. Zumino, "Supergauge invariance and the Gell-Mann Low eigenvalue", *Nucl. Phys.* **B77** (1974) 41.

[26] P. Howe and P.C. West, "Chiral correlators in Landau–Ginzburg theories and N=2 superconformal models", *Phys. Lett.* **B227** (1989) 397; P.C. West, "A comment on the non-renormalization theorem in supersymmetric gauge theories" ,*Phys. Lett.* **B258** (1991) 369.

[27] I. Jack, T. Jones and P.C. West, "Not the no-renormalization theorem", *Phys. Lett.* **B258** (1991) 375.

[28] P.C. West, "Quantum corrections to the supersymmetric effective superpotential and resulting modifications of patterns of symmetry breaking", *Phys. Lett.* **B261** (1991) 396.

[29] M. Shifman and A. Vainshtein, "On holomorphic dependence and infrared effects in supersymmetric gauge theories", *Nucl. Phys.* **B359** (1991) 571.

[30] L.J. Dixon, V. Klapunovski and J. Louis, "Moduli dependence of string loop corrections to gauge coupling constants", *Nucl. Phys.* **B355** (1991) 649.

[31] M.E. Peskin, "Duality in supersymmetric Yang–Mills theory", `hep-th/9702094`.

[32] M. Shifman, "Non–perturbative dynamics in supersymmetric gauge theories", `hep-th/9704114`.

[33] V. Novikov, M. Shifman, A. Vainshtein and V. Zakharov, "Exact Gell-Mann–Low function of supersymmetric Yang–Mills theories from instanton calculus", *Nucl. Phys.* **B229** (1983) 381; "Beta function in supersymmetric gauge theories: instantons versus traditional approach", *Phys. Lett.* **B166** (1986) 334.

[34] N. Seiberg and E. Witten, "Electric-magnetic duality, monopole condensation, and confinement in N=2 supersymmetric Yang–Mills theory", *Nucl. Phys.* **B426** (1994) 19; Erratum-ibid. **B430** (1994) 485; "Monopoles, duality and chiral symmetry breaking in N=2 supersymmetric QCD", *Nucl. Phys.* **B431** (1994) 484.

[35] L. Brink, J. Scherk and J.H. Schwarz, "Supersymmetric Yang-Mills theories", *Nucl. Phys.* **B121** (1977) 77.

[36] F. Gliozzi, D.I. Olive and J. Scherk, "Supersymmetry, supergravity and the dual spinor model", *Nucl. Phys.* **B122** (1977) 253.

[37] K. Montonen and D.I. Olive, "Magnetic monopoles as gauge particles", *Phys. Lett.* **66B** (1977) 61.

[38] J. Maldacena, "The large N limit of superconformal field theories and supergravity", *Adv. Theor. Math. Phys.* **2** (1998) 231,`hep-th/9711200`.

[39] M. Roček and W. Siegel, "On off-shell supermultiplets", *Phys. Lett.* **105B** (1981) 275.

[40] V. Rivelles and J.G. Taylor, "Field redefinition rules for auxiliary field searches", *J. Phys.: Math. Gen.* **A15** (1982) 163.

[41] P. Howe, K.S. Stelle and P.K. Townsend, "Supercurrents", *Nucl. Phys.* **B192** (1981) 332.

[42] P.L. White, "Analysis of the superconformal cohomology structure of N=4 super Yang–Mills", *Class. Quant. Grav.* **9** (1992) 413.

[43] S. Ferrara and B. Zumino, "Transformation properties of the supercurrent", *Nucl. Phys.* **B87** (1975) 207.

[44] E. Bergshoeff, M. de Roo and B. de Wit, "Extended conformal supergravity", *Nucl. Phys.* **B182** (1981) 173.

[45] D.T.R. Jones, "Charge renormalization in a supersymmetric Yang–Mills theory", *Phys Lett.* **72B** (1977) 199; E. Poggio and H. Pendleton, "Vanishing of charge renormalization and anomalies in a supersymmetric gauge theory", *Phys Lett.* **72B** (1977) 200.

[46] A.A. Vladimirov and O.V. Tarasov, "Vanishing of the three-loop charge renormalization function in a supersymmetric gauge theory", *Phys. Lett.* **96B** (1980) 94.

[47] M. Grisaru, M. Roček and W. Siegel, "Zero value of the three-loop β function in N=4 supersymmetric Yang–Mills theory", *Phys. Rev. Lett.* **45** (1980) 1063.

[48] W.E. Caswell and D. Zanon, "Zero three-loop beta function in the N=4 supersymmetric Yang–Mills theory", *Nucl. Phys.* **B182** (1981) 125.

[49] P.S. Howe, K.S. Stelle and P.K. Townsend, "The relaxed hypermultiplet: an unconstrained N=2 superfield theory", *Nucl. Phys.* **B214** (1983) 519; P.S. Howe, K.S. Stelle and P.C. West, "A class of finite four-dimensional supersymmetric field theories", *Phys. Lett.* **124B** (1983) 55; P.S. Howe, K.S. Stelle and P.K. Townsend, "Miraculous ultraviolet cancellations in supersymmetry made manifest", *Nucl. Phys.* **B236** (1984) 125.

[50] S. Mandelstam, "Light-cone superspace and the ultraviolet finiteness of the N=4 model", *Nucl. Phys.* **B213** (1983) 149.

[51] M.F. Sohnius and P.C. West, "Conformal invariance in N=4 supersymmetric Yang–Mills theory", *Phys. Lett.* **100B** (1981) 245.

[52] S.L. Adler, J.C. Collins and A. Duncan, "Energy-momentum–tensor trace anomaly in spin 1/2 quantum electrodynamics", *Phys Rev.* **D15** (1977) 1712.

[53] H.A. Kramers and G.H. Wannier, "Statistics of the two-dimensional ferromagnet", *Phys. Rev.* **60** (1941) 252.

[54] S. Coleman, "Quantum sine-Gordon equation as the massive Thirring model", *Phys. Rev.* **D11** (1975) 2088.

[55] S. Mandelstam, "Soliton operators for the quantized sine-Gordon model", *Phys. Rev.* **D11** (1975) 3026.

[56] P.A.M. Dirac, "Quantized singularities in the electromagnetic field", *Proc. Roy. Soc.* **A33** (1931) 60.

[57] J. Schwinger, "A magnetic model of matter", *Science* **165** (1969) 757; D. Zwanziger, "Quantum field theories of particles with both electric and magnetic charges", *Phys. Rev.* **176** (1968) 1489.

[58] D. Olive, "Exact electromagnetic duality", *Nucl. Phys. Proc. Suppl.* **45A** (1996) 88, `hep-th/9508089`.

[59] J.A. Harvey, "Magnetic monopoles, duality and supersymmetry", `hep-th/9603083`.

[60] P. Di Vecchia, "Duality in N=2,4 supersymmetric gauge theories", `hep-th/9803026`.

[61] P. Goddard and D. Olive, "Magnetic monopoles in unified gauge theories", *Rep. on Prog. in Phys.* **41** (1978) 1357.

[62] E.B. Bogomol'nyi, "Stability of classical solutions", *Sov. J. Nucl. Phys.* **24** (1976) 449.

[63] N.S. Manton, "A remark on the scattering of BPS monopoles", *Phys. Lett.* **110B** (1982) 54.

[64] H. Osborn, "Topological charges for N=4 supersymmetric gauge theories and monopoles of spin one", *Phys. Lett.* **83B** 321.

[65] E. Witten and D. Olive, "Supersymmetry algebras that include topological charges", *Phys. Lett.* **78B** (1977) 97.

[66] H. Osborn, "Semiclassical methods for quantizing monopole field configurations" in *Monopoles in quantum field theory*, ed. by N.S. Craigie, P. Goddard and W. Nahm, World Scientific, Singapore, 1982.

[67] J.P. Gauntlett, "Duality and supersymmetric monopoles", `hep-th/9705025`.

[68] M.F. Atiyah and N.J. Hitchin, *The geometry and dynamics of magnetic monopoles*, Princeton University Press, Princeton, NJ, 1988.

[69] J.P. Gauntlett, "Low energy dynamics of N=2 supersymmetric monopoles", *Nucl. Phys.* **B411** (1994) 443, `hep-th/9305068`; J.D. Blum, "Supersymmetric quantum mechanics of monopoles in N=4 Yang–Mills theory", *Phys. Lett.* **B333** (1994) 92, `hep-th/9401133`.

[70] E. Witten, "Constraints on supersymmetry breaking", *Nucl. Phys.* **B202** (1982) 253.

[71] C. Callias, "Index theorems on open spaces", *Comm. Math. Phys.* **62** (1978) 213.

[72] A. Sen, "Dyon-monopole bound states, self dual harmonic forms on the multi-monopole moduli space", *Phys. Lett.* **329B** (1994) 217, `hep-th/9402032`.

[73] G. Segal and A. Selby, "The cohomology of the space of magnetic monopoles", *Comm. Math Phys.* **177** (1996) 775; M. Porrati, "How to find H monopoles in brane dynamics", *Phys. Lett.* **B387** (1996) 492, `hep-th/9607082`.

[74] O. Piguet and A. Rouet, "Supersymmetric BPHZ renormalization 2: supersymmetric extension of pure Yang–Mills model", *Nucl. Phys.* **B108** (1976) 265.

[75] D. Storey, "General gauge calculations in N=4 super Yang–Mills theory", *Phys. Lett.* **105B** (1981) 171.

[76] A.J. Parkes and P.C. West, "N=1 supersymmetric mass terms in the N=4 supersymmetric Yang–Mills theory", *Phys. Lett.* **122B** (1983) 365.

[77] N. Arkani-Hamed and H. Murayama, "Holomorphy, Rescaling Anomalies and exact β functions in supersymmetric gauge theories", `hep-th/9707133`.

[78] S. Arnone, C. Fusi and K. Yoshida, "Exact renormalization group equation in presence of rescaling anomaly", `hep-th/9812022`.

[79] S. Kovacs, "A perturbative re-analysis of N=4 supersymmetric Yang–Mills theory", `hep-th/9902047`.

[80] J.W. Juer and D. Storey, "One-loop renormalization of superfield Yang–Mills theories", *Nucl. Phys.* **B216** (1983) 185.

[81] M.A. Namazie, A. Salam and J. Strathdee, "Finiteness of broken N=4 super Yang–Mills theory", *Phys.Rev.* **D28** (1983) 1481.

[82] J.W. Juer and D. Storey, "Nonlinear renormalization in superfield gauge theories", *Phys. Lett.* **119B** (1982) 125.

[83] T. Kinoshita, "Mass singularities of Feynman amplitudes", *J. Math. Phys.* **3** (1962) 650; T.D. Lee and M. Nauenberg, "Degenerate systems and mass singularities", *Phys. Rev.* **133** (1964) B1562.

[84] T. Kinoshita and A. Ukawa, "New approach to the singularities of Feynman amplitudes in the zero-mass limit", *Phys. Rev.* **D13** (1976) 1573; E.C. Poggio and H.R. Quinn, "The infrared behavior of zero-mass Green's functions and the absence of quark confinement in perturbation theory", *Phys. Rev.* **D14** (1976) 578; G. Sterman, "Kinoshita's theorem in Yang–Mills theories", *Phys. Rev.* **D14** (1976) 2123.

[85] A.A. Belavin, A.M. Polyakov, A.S. Schwartz and Yu.S. Tyupkin, "Pseudoparticle solutions of the Yang–Mills equations", *Phys. Lett.* **59B** (1975) 85.

[86] G. 't Hooft, "Computation of the quantum effects due to a four-dimensional pseudoparticle", *Phys. Rev.* **D14** (1976) 3432.

[87] D. Amati, K. Konishi, Y. Meurice, G.C. Rossi and G.Veneziano, "Nonperturbative aspects in supersymmetric gauge theories", *Phys. Rep.* **C162** (1988) 169.

[88] *Instantons in gauge theories*, ed. by M. Shifman, World Scientific, 1994.

[89] M. Bianchi, M.B. Green, S. Kovacs and G.C. Rossi, "Instantons in supersymmetric Yang–Mills and D-instantons in IIB superstring theory", *JHEP* **9808** (1998) 013, `hep-th/9807033`.

[90] N. Dorey, V.V. Khoze, M.P. Mattis and S. Vandoren, "Yang–Mills instantons in the large-N limit and the AdS/CFT correspondence", `hep-th/9808157`.

[91] N. Dorey, T.J. Hollowood, V.V. Khoze, M.P. Mattis and S. Vandoren, "Multi-instantons and Maldacena's conjecture", `hep-th/9810243`.

[92] N. Dorey, T.J. Hollowood, V.V. Khoze, M.P. Mattis and S. Vandoren, "Multi-instanton calculus and the AdS/CFT correspondence in N=4 superconformal field theory", `hep-th/9901128`.

[93] R. Jackiw and C. Rebbi, "Conformal properties of a Yang–Mills pseudoparticle", *Phys. Rev.* **D14** (1976) 517.

[94] M. Atiyah and A. Ward, "Instantons and algebraic geometry", *Comm. Math. Phys.* **55** (1977) 117.

[95] L.G. Yaffe, "Quantizing gauge theories: nonclassical field configurations, broken symmetries, and gauge copies", *Nucl. Phys.* **B151** (1979) 247.

[96] C.W. Bernard, N.H. Christ, A.H. Guth and E.J. Weinberg, "Pseudoparticle parameters for arbitrary gauge groups", *Phys. Rev.* **D16** (1977) 2967.

[97] C.W. Bernard, "Gauge zero modes, instanton determinants and quantum-chromodynamic calculations", *Phys. Rev.* **D19** (1979) 3013.

[98] S. Coleman, "Uses of instantons", in *The whys of subnuclear physics*, ed. by A. Zichichi, Plenum Press, New York, 1979.

[99] M. Atiyah and I. Singer, "The index of elliptic operators. 1", *Ann. Math.* **87** (1968) 484.

[100] D. Amati, G.C. Rossi and G. Veneziano, "Instanton effects in supersymmetric gauge theories", *Nucl. Phys.* **B249** (1985) 1.

[101] M. Atiyah, V. Drinfeld, N. Hitchin and Yu. Manin, "Construction of instantons", *Phys. Lett.* **A65** (1978) 186.

[102] E. Corrigan, D.B. Fairlie, P. Goddard and S. Tampleton, "A Green's function for the general selfdual gauge field", *Nucl. Phys.* **B140** (1978) 31.

[103] E. Corrigan, P. Goddard and S. Tampleton, "Instanton Green's functions and tensor products", *Nucl. Phys.* **B151** (1979) 93.

[104] N.H. Christ, E.J. Weinberg and N.S. Stanton, "General selfdual Yang–Mills solutions", *Phys. Rev.* **D18** (1978) 2013.

[105] H. Osborn, "Semiclassical functional integrals for selfdual gauge fields", *Ann. Phys.* **135** (1981) 373.

[106] D.Z. Freedman, S.D. Mathur, A. Matusis and L. Rastelli, "Comments on 4-point functions in the CFT/AdS correspondence", `hep-th/9808006`.

[107] J.H. Brodie and M. Gutperle, "String corrections to four-point functions in the AdS/CFT correspondence", `hep-th/9809067`.

[108] M. Bianchi and S. Kovacs, "Yang–Mills instantons vs. type IIB D-instantons", `hep-th/9811060`.

[109] H. Liu, "Scattering in anti-de Sitter space and operator product expansion", hep-th/9811152.

[110] M. Bianchi, S. Kovacs, G.C. Rossi and Ya.S. Stanev, in preparation.

[111] P. Howe and P. West, "Non-perturbative Green functions in theories with extended superconformal symmetry", hep-th/9509140.

[112] Z. Bern and G. Chalmers, "Factorization in one-loop gauge theory", *Nucl. Phys.* **B447** (1995) 466.

[113] V.V. Khoze, M.P. Mattis and J.Slater, "The instanton hunter's guide to supersymmetric gauge theories", hep-th/9804009.

[114] M.B. Green and M. Gutperle, "Effects of D-instantons", *Nucl. Phys.* **B498** (1997) 195, hep-th/9701093

[115] M.B. Green and M. Gutperle,"D-particle bound states and the D-instanton measure", *JHEP* **9801** (1998) 005, hep-th/9711107.

[116] G. Moore, N. Nekrasov and S. Shatashvili, "D-particle bound states and generalized instantons", hep-th/9803265.

[117] W. Krauth, H. Nicolai and M. Staudacher, "Monte Carlo approach to M theory", *Phys. Lett.* **B431** (1998) 31, hep-th/9803117; W. Krauth and M. Staudacher, "Finite Yang–Mills integrals", *Phys. Lett.* **B435** (1998) 350, hep-th/9804199.

[118] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, "Gauge theory correlators from non-critical string theory", *Phys. Lett.* **B428** (1998) 10, hep-th/9802109.

[119] E. Witten, "Anti de Sitter space and holography", *Adv. Theor. Math. Phys.* **2** (1998) 253, hep-th/9802150.

[120] G. 't Hooft, "Dimensional reduction in quantum gravity", in *Salamfest 1993*, gr-qc/9310026.

[121] L. Susskind, "The world as a hologram", *J. Math. Phys.* **36** (1995) 6377.

[122] G. 't Hooft, "A planar diagram theory for strong interactions", *Nucl. Phys.* **B72** (1974) 461.

[123] I.R. Klebanov, "From three-branes to large N gauge theories", hep-th/9901018; M.R. Douglas and S. Randjbar-Daemi, "Two lectures on AdS/CFT correspondence", hep-th/9902022; J.L. Petersen, "Introduction to the Maldacena conjecture on AdS/CFT", hep-h/9902131; P. Di Vecchia, "Introduction to AdS/CFT equivalence", hep-th/9903007.

[124] L. Brink, P. Di Vecchia and P. Howe, "A locally supersymmetric and reparametrization invariant action for the spinning string", *Phys. Lett.* **65B** (1976) 471.

[125] A.M. Polyakov, "Quantum geometry of bosonic strings", *Phys. Lett.* **103B** (1981) 207.

[126] J.H. Schwarz, "Covariant field equations of chiral N=2 D=10 supergravity", *Nucl. Phys.* **B226** (1983) 269.

[127] P. Pasti, D. Sorokin and M. Tonin, "On Lorentz invariant actions for chiral p-forms", *Phys. Rev.* **D55** (1997) 6292, hep-th/9611100.

[128] J.H. Schwarz, "An SL(2,Z) multiplet of type IIB superstrings", *Phys.Lett.* **B360** (1995) 13, Erratum-ibid. **B364** (1995), 252.

[129] M.B. Green and P. Vanhove, "D-instantons, Strings and M-theory", *Phys. Lett.* **B408** (1997) 122, hep-th/9704145.

[130] M.B. Green, M. Gutperle and H. Kwon, "Sixteen-fermion and related terms in M theory on T^2", *Phys. Lett.* **B421** (1998) 149, hep-th/9710151.

[131] L.J. Dixon, V. Kaplunovsky and C. Vafa, "On four-dimensional gauge theories from type II superstrings", *Nucl. Phys.* **B294** (1987) 43.

[132] G.T. Horowitz and A. Strominger, "Black strings and p-branes", *Nucl. Phys.* **B360** (1991) 197; M.J. Duff and J.X. Lu, "The self-dual type IIB superthree-brane", *Phys. Lett.* **B273** (1991) 409; M.J. Duff, R. Khuri and J.X. Lu, "String solitons", *Phys. Rep.* **C259** (1995) 213.

[133] J. Polchinski, "Dirichlet branes and Ramond-Ramond charges", *Phys. Rev. Lett.* **75** (1995) 4724, hep-th/9510017.

[134] J. Polchinski, "TASI lectures on D-branes", hep-th/9611050.

[135] E. Witten, "Bound states of strings and D-branes", *Nucl. Phys.* **B460** (1996) 335.

[136] J.M. Maldacena and A. Strominger, "Black hole grey body factors and D-brane spectroscopy" *Phys. Rev.* **D55** (1997) 861.

[137] S. Gubser, I.R. Klebanov and A. Peet, "Entropy and temperature of black 3-brane", *Phys. Rev.* **D54** (1996) 3915, `hep-th/9602135`; I.R. Klebanov and A.A. Tseytlin, "Entropy of near-extremal black p-branes", *Nucl. Phys.* **B475** (1996) 164, `hep-th/9604089`; I.R. Klebanov and A.A. Tseytlin, "Intersecting M-branes as four dimensional black holes", *Nucl. Phys.* **B475** (1996) 179, `hep-th/9604166`; I.R. Klebanov, "World volume approach to absorption by non-dilatonic branes", *Nucl. Phys.* **B496** (1997) 231, `hep-th/9702076`; S. Gubser, I.R. Klebanov and A.A. Tseytlin, "String theory and classical absorption by branes", *Nucl. Phys.* **B499** (1997) 217, `hep-th/9703040`; S. Gubser and I. Klebanov, "Absorption by branes and Schwinger terms in the world volume theory", *Phys. Lett.* **B413** (1997) 41, `hep-th/9708005`; G.T. Horowitz and A. Strominger, "Black strings and p-branes", *Nucl. Phys.* **B360** (1991) 197.

[138] M. Gunaydin and N. Marcus, "The unitarity supermultiplet of N=8 conformal superalgebra involving fields of spin ≤ 2", *Class. Quant. Grav.* **2** (1985) L11-17.

[139] H.J. Kim, L.J. Romans, and P. van Nieuwenhuizen, "Mass spectrum of chiral ten-dimensional N=2 supergravity on S^5", *Phys. Rev.* **D32** (1985) 389.

[140] S. Ferrara and C. Frønsdal, "Conformal Maxwell Theory as a Singleton Field Theory on AdS_5, IIB Three-Branes and Duality", *Class. Quant. Grav.* **15** (1998) 2153, `hep-th/9712239`.

[141] L. Andrianopoli and S. Ferrara, "K–K excitations on $AdS_5 \times S^5$ as N=4 "primary" superfields", *Phys. Lett.* **B430** (1998) 248, `hep-th/9803171`.

[142] S. Ferrara, C. Frønsdal and A. Zaffaroni, "On N=8 supergravity on AdS_5 and N=4 superconformal Yang–Mills theory", `hep-th/9802203`.

[143] L. Andrianopoli and S. Ferrara, "'Non chiral' primary superfields in the AdS_{d+1}/CFT_d correspondence", `hep-th/9807150`.

[144] R.R. Metsaev and A.A. Tseytlin, "Supersymmetric D3 brane action in $AdS_5 \times S^5$", *Phys. Lett.* **B436** (1998) 281, `hep-th/9806095`; R. Kallosh, "Superconformal Actions in Killing Gauge", `hep-th/9807206`; R. Kallosh and J. Rahmfeld, "The GS String Action on $AdS_5 \times S^5$", `hep-th/9808038`; R. Kallosh and A.A. Tseytlin, "Simplifying superstring action on $AdS_5 \times S^5$", `hep-th/9808088`; I. Pesando, "A κ Gauge Fixed Type IIB Superstring Action on $AdS_5 \times S^5$", `hep-th/9808020`; I. Pesando, "All roads lead to Rome: Supersolvables and Supercosets", `hep-th/9808146`; J. Park and S.-J. Rey, "Dual D-Brane Actions on $AdS_5 \times S^5$", `hep-th/9810154`.

[145] D.Z. Freedman, S.D. Mathur, A. Matusis and L. Rastelli, "Correlation Functions in the CFT_d/AdS_{d+1} Correspondence", `hep-th/9804058`.

[146] M. Henningson and K. Sfetsos, "Spinors in the AdS/CFT correspondence", *Phys. Lett.* **B431** (1998) 63, `hep-th/9803251`; A.M. Ghezelbash, K. Kaviani, S. Parvizi and A.H. Fatollahi, "Interacting spinors-scalars and AdS/CFT correspondence", *Phys. Lett.* **B435** (1998) 291, `hep-th/9805162`.

[147] D.Z. Freedman, E. D'Hoker and W. Skiba, "Field theory tests for correlators in the AdS/CFT correspondence", `hep-th/9807098`.

[148] S. Lee, S. Minwalla, M. Rangamani and N. Seiberg, "Three-point functions of chiral operators in $D = 4$ N=4 SYM at large N_c", `hep-th/9806074`.

[149] S. Kachru and E. Silverstein, "$4d$ conformal field theories and strings on orbifolds", `hep-th/9802183`.

[150] A. Lawrence, N. Nekrasov and C. Vafa, "On conformal field theories in four dimensions", `hep-th/9803015`.

[151] E. Witten, "Anti de Sitter space, termal phase transition and confinement in gauge theories", *Adv. Theor. Math. Phys.* **2** (1998), `hep-th/9803131`.

[152] S.-J. Rey, S. Theisen and J. Yee, "Wilson–Polyakov loop at finite temperature in large N gauge theories and anti de Sitter supergravity", `hep-th/9803135`; A. Brandhuber, N. Itzhaki, J. Sonnenschein and S. Yankielowicz, "Wilson loops in the large N limit at finite temperature", `hep-th/9803137`; E. Witten, "Baryons and branes in anti-de Sitter space", *JHEP* **9807** (1998) 006,

hep-th/9805112; D.J. Gross and H. Ooguri, "Aspects of large N gauge theory dynamics as seen by string theory", *Phys.Rev.* **D58** (1998) 106002, hep-th/9805129.

[153] C. Csaki, H. Ooguri, Y. Oz, J. Terning, "Glueball mass spectrum from supergravity", hep-th/9806021; H. Ooguri, H. Robins and J. Tannenhauser, "Glueballs and their Kaluza–Klein cousins", 9806171.

[154] A. Polyakov, "The wall of the cave", hep-th/9809057.

[155] L. Dixon and J. Harvey, "String theories in ten dimensions without space-time supersymmetry, *Nucl. Phys.* **B274** (1986) 93; N. Seiberg and E. Witten, "Spin structures in string theory", *Nucl. Phys.* **B276** (1986) 272.

[156] M. Bianchi and A. Sagnotti, "On the systematics of open string theories", *Phys. Lett.* **B247** (1990) 517.

[157] O. Bergman and M. Gaberdiel, "A non supersymmetric open string theory and S-duality", *Nucl. Phys.* **B499** (1997) 183, hep-th/9701137.

[158] I.R. Klebanov and A.A. Tseytlin, "D-branes and dual gauge theories in type 0 strings", hep-th/9811035.

[159] J. Minahan, "Glueball mass spectra and other issues for supergravity duals of QCD models", hep-th/9811156.

[160] I.R. Klebanov and A.A. Tseytlin, "Asymptotic freedom and infrared behaviour in the type 0 string approach to gauge theory", hep-th/9812089.

[161] H. Liu, A.A. Tseytlin, "d=4 super Yang–Mills, d=5 gauged supergravity, and d=4 conformal supergravity", *Nucl. Phys.* **B533** (1998) 88, hep-th/9804083.

[162] M. Henneaux, "Boundary terms in the AdS/CFT correspondence for spinor fields", hep-th/9902137.

[163] S. Ferrara and A. Zaffaroni, "Bulk gauge fields in AdS supergravity and supersingletons", hep-th/9807090.

[164] K. Konishi, "Anomalous supersymmetry transformation of some composite operators in SQCD", *Phys. Lett.* **B135** (1984) 439.

[165] D. Anselmi, D.Z. Freedman, M. Grisaru and A. Johansen, "Nonperturbative formulas for central functions of supersymmetric gauge theories", hep-th/9708042.

[166] H. Liu and A.A. Tseytlin, "On four-point functions in the CFT/AdS correspondence", hep-th/9807097.

[167] E. D'Hoker and D.Z. Freedman, "General scalar exchange in AdS_{d+1}", hep-th/9811257.

[168] E. D'Hoker and D.Z. Freedman, "Gauge boson exchange in AdS_{d+1}", hep-th/9809179.

[169] E. D'Hoker, D.Z. Freedman, S.D. Mathur, A. Matusis and L. Rastelli, "Graviton and gauge boson propagators in AdS_{d+1}", hep-th/9902042.

[170] T. Banks and M.B. Green, "Nonperturbative effects in $AdS_5 \times S^5$ string theory and d=4 SUSY Yang–Mills", *JHEP* **9805** (1998) 002, hep-th/9804170.

[171] F. Gonzalez-Rey, I. Park, K. Schalm, "A note on four point functions of conformal operators in N=4 super Yang–Mills", hep-th/9811155.

[172] B. Eden, P.S. Howe, C. Schubert, E. Sokatchev, P.C. West, "Four point functions in N=4 supersymmetric Yang–Mills theory at two loops", hep-th/9811172.

[173] G. Chalmers and K. Schalm, "The large N limit of four-point functions in N=4 super Yang–Mills theory from anti-de Sitter supergravity", hep-th/9810051.

[174] G. Chalmers and K. Schalm, "Holographic normal ordering and multi-particle states in the AdS/CFT correspondence", hep-th/9901144.

[175] G.W. Gibbons, M.B. Green and M.V. Perry, "Instantons and seven-branes in type IIB superstring theory", *Phys. Lett.* **B370** (1996) 37, hep-th/9511080.

[176] R.I. Nepomechie, "Magnetic monopoles from antisymmetric tensor gauge fields", *Phys. Rev.* **D31** (1985) 1921; C. Teitelboim, "Gauge invariance for extended objects, *Phys. Lett.* **167B** (1986) 63; C. Teitelboim, "Monopoles of higher rank", *Phys. Lett.* **B167** (1986) 69.

[177] P.S. Howe and P.C. West, "The complete N=2 D=10 supergravity", *Nucl. Phys.* **B238** (1984) 181.

[178] A. Kehagias and H. Partuche, "The exact quartic effective action for the IIB superstring, *Phys. Lett.* **B422** (1998) 109, `hep-th/9710023`; B. Pioline, "A note on non-perturbative R^4 couplings", *Phys.Lett.* **B431** (1998) 73, `hep-th/9804023`.

[179] M.B. Green and S. Sethi, "Supersymmetry constraints on type IIB supergravity", `hep-th/9808061`.

[180] G. 't Hooft, "Magnetic monopoles in unified gauge theories", *Nucl. Phys.* **B79** (1974) 276.

[181] A. Polyakov, "Particle spectrum in quantum field theory", *JETP Lett.* **20** (1974) 194.

[182] E. Corrigan, D. Olive, D.B. Fairlie and J. Nuyts, "Magnetic monopoles in SU(3) gauge theories" *Nucl. Phys.* **B106** (1976) 475.

[183] M.K. Prasad and C.M. Sommerfield, "An exact classical solution for the 't Hooft monopole and the Julia–Zee dyon", *Phys. Rev. Lett.* **35** (1975) 760.

[184] B. Julia and Zee, "Poles with both magnetic and electric charges in non Abelian gauge theory", *Phys. Rev.* **D11** (1975) 2227.

[185] E. Witten, "Dyons of charge $\frac{e\theta}{2\pi}$", *Phys. Lett.* **86B** (1979) 283.